

# Risk Factors for Health Care-Associated Bloodstream Infections in a Neonatal Intensive Care Unit

Sabahattin Ertugrul,<sup>1</sup> Fesih Aktar,<sup>2,\*</sup> Ilyas Yolbas,<sup>3</sup> Ahmet Yilmaz,<sup>4</sup> Bilal Elbey,<sup>5</sup> Ahmet Yildirim,<sup>3</sup> Kamil Yilmaz,<sup>3</sup> and Recep Tekin<sup>6</sup>

<sup>1</sup>Department of Neonatology, Medical Faculty, Dicle University, Diyarbakir, Turkey

<sup>2</sup>Department of Pediatric Infectious Disease, Medical Faculty, Dicle University, Diyarbakir, Turkey

<sup>3</sup>Department of Pediatrics, Medical Faculty, Dicle University, Diyarbakir, Turkey

<sup>4</sup>Department of Family Medicine, Medical Faculty, Dicle University, Diyarbakir, Turkey

<sup>5</sup>Department of Immunology, Medical Faculty, Dicle University, Diyarbakir, Turkey

<sup>6</sup>Clinical Microbiology and Infectious Disease, Medical Faculty, Dicle University, Diyarbakir, Turkey

\*Corresponding author: Fesih Aktar, Department of Pediatric Infectious Disease, Medical Faculty, Dicle University, Diyarbakir, Turkey. Tel: +90-5056690064, Fax: +90-4122488523, E-mail: fesihaktar@yahoo.com

Received 2015 December 30; Revised 2016 April 18; Accepted 2016 May 28.

## Abstract

**Background:** Healthcare-associated bloodstream infections (HCA-BSI) are a major cause of morbidity and mortality in neonatal intensive care units (NICUs).

**Objectives:** We aimed to determine the causative organisms and risk factors of HCA-BSIs in NICUs.

**Methods:** This study was performed between January 2011 and December 2014 in the neonatal intensive care unit of Dicle university, Turkey. The study consisted of 126 patients (infected group) with positive blood culture and 126 randomly selected patients (uninfected control group) with negative blood culture after four days of hospitalization.

**Results:** We found that the most common causative agents isolated from nosocomial infections (NIs) were 20.7% *Staphylococcus epidermidis*, 26.7% *Klebsiella* spp., and 13.3% *Acinetobacter* spp. Incidences of low gestational age, low birth weight, vaginal birth type, and long length of hospitalization were higher in the infected neonates than in the uninfected neonates. In the univariate analysis, surgical operation, ventriculoperitoneal shunt, use of umbilical catheter, nasogastric or orogastric tube, urinary catheter, mechanical ventilation, surfactant treatment, erythrocyte transfusion, plasma transfusion, thrombocyte transfusion, total parenteral nutrition infusion, intracranial hemorrhage, length of hospital stay, fifth-minute Apgar score, and total parenteral nutrition time were significantly associated with NIs. In the multiple logistic regression analysis, fifth-minute Apgar, use of erythrocyte transfusion and surgical operation were found as the independent risk factors for HCA-BSI.

**Conclusions:** This study determined the causative organisms and risk factors of HCA-BSIs in NICUs.

**Keywords:** Healthcare-Associated Infections, Bloodstream Infections, Neonates, Risk Factors

## 1. Background

Healthcare-associated bloodstream infections (HCA-BSI) rated in neonatal intensive care units (NICUs) between 1.8% and 74.3% in several reports are a major cause of morbidity and mortality in NICUs and affect the cost of medical care by increasing resource consumption and duration of hospitalization (1-3). The most common type of neonatal healthcare-associated infection (HC-AI) is bloodstream infection (BSI). A neonate may be at risk of infection through one or many intrinsic and extrinsic factors, such as gestational age and presence of a single or multiple invasive devices (4). In developed and developing countries, most nosocomial infections (NIs) in NICUs are related to a longer duration of hospitalization, low birth weight and gestational age, respiratory diseases, invasive interventions, and medical treatments (5). HCA-BSIs, device-associated HC-AI in particular, are generally common in newborns because of their insufficient immune system and mechanical barriers as well as the lack of protective flora (6). Therefore, among hospitalized patients, neonates

belong to those at the highest risk of HCA-BSIs. In addition, neonatal HCA-BSIs not only have high mortality but also increased risk of subsequent adverse outcomes, including white matter injury to the preterm brain and attendant neurodisability (7).

A few studies have been conducted on the risk factors of HCA-BSIs in NICUs in Turkey (8, 9). In this study, we described the demographic features of the patients, the causative organisms, and the risk factors of HCA-BSIs in NICUs.

## 2. Objectives

The purpose of the present study was to identify the risk factors associated with the development of HCA-BSIs and the most frequent causative agents of the same in NICUs. Knowledge of the modifiable risk factors for HCA-BSIs would enable developing countries to implement interventions, thereby decreasing HCA-BSIs and the associated complications.

### 3. Methods

This retrospective study was conducted in the clinic of neonatology in Dicle university between January 2011 and December 2014. The neonatal unit has 34 incubators, 20 mechanical ventilators, and 2 open intensive care cots. This study included 126 patients (infected group) with positive blood cultures and 126 randomly selected patients (uninfected control group) with negative blood cultures after four days of hospitalization. Infection control doctors and nurses conducted hospital infection surveillance actively and prospectively. During the period of hospitalization, the infection control committees recorded patient information every day using the patients' follow-up forms. The diagnosis of HCA-BSIs was made according to the criteria of the US center for disease control and prevention (10).

Healthcare-associated bloodstream infection (HCA-BSI): Patients with no sepsis on admission and who had microorganisms isolated from blood cultures taken 48 - 72 hours after birth on suspicion of sepsis according to the clinical signs and/or laboratory findings were diagnosed as HCA-BSI. Clinical findings, including apnea, bradycardia, hypothermia, hyperthermia, circulatory disorder, lethargy, hypotonia, and feeding difficulty, and laboratory findings, including leukocytosis, leucopenia, thrombocytopenia, a ratio of immature/mature neutrophils > 0.25, and a C-reactive protein value of > 0,05 mg/dL, were considered significant. Patients with signs of sepsis and showed no growth in culture and patients with duration of hospitalization of less than 48 - 72 hours were excluded from the study.

Strains of *Staphylococcus epidermidis* were considered involved in the infection if they represented the only microorganisms isolated from the blood specimen in the presence of clinical signs or symptoms of infection. Most authorities recommend obtaining two independent cultures to fulfill two workups of an episode of suspected bloodstream infection (11).

The blood samples taken from patients with suspected bacteremia and/or sepsis were inoculated into blood culture bottles and incubated in BACTEC 9120 and 9240 (Becton Dickinson, MD, USA) blood culture systems for 7 - 10 days at 37°C. The blood samples in which bacterial growth was detected were inoculated into 5% sheep blood agar, EMB agar, and chocolate agar media. These media were incubated at 35 ± 2°C for 20 - 24 hours. Wound swabs and other clinical specimens were directly inoculated into 5% sheep blood agar, EMB agar, and chocolate agar and incubated at 35 ± 2°C for 20 - 24 hours. All the strains isolated from the clinical specimens were identified using conventional methods and BD Phoenix™ 100 (Becton Dickinson, MD, USA).

Data were obtained from the database of the hospital infection control committee and patients' medical records. Gender, gestational age, birth weight, birth presentation, type of delivery, length of hospitalization, results of blood cultures, surgical operation, ventriculoperitoneal shunt, tracheostomy procedure, diagnosis with intracranial hemorrhage, phototherapy, tracheostomy procedure, exchange transfusion, (four days of mechanical ventilation and neonatologist's prediction of prolonged ventilation were the common indication), Apgar score point (fifth minute), use of umbilical catheter, nasogastric or orogastric tube, urinary catheter, mechanical ventilation, use of surfactant, erythrocyte transfusion, plasma transfusion, thrombocyte transfusion, intravenous immunoglobulin, and total parenteral nutrition infusion were recorded prospectively.

The cases not appropriate for hospital acquired infections (HAIs) because of clinical and laboratory findings and those who had a single blood culture with isolated central nervous system considered as contamination were excluded from the study.

The study protocol was approved by the Dicle university faculty of medicine ethic committee.

#### 3.1. Statistical Analysis

The two-sided statistical tests were performed using SPSS for Windows (version 18.0; SPSS, Inc., Chicago, IL). A P value of < 0.05 was considered statistically significant. For the comparative analysis between groups (case vs. control), the  $\chi^2$  test was used for categorical variables, and either the student t test or the Mann-Whitney test was used for the continuous variables. In the univariate analysis, differences were considered significant at P < 0.05. For the identification of independent factors of NI that could influence disposition, multivariable logistic regression analyses (polytomous responses) were performed to calculate the odds ratio and the corresponding 95% confidence intervals.

### 4. Results

A total of 126 positive blood cultures that were proven HCA-BSI attacks were found in 268 cases of suspected HCA-BSI. The study included 126 neonates (47.6% females and 52.4% males) with positive blood cultures and 126 neonates (44.4% females and 55.6% males) with negative blood cultures. Incidence of low gestational age, low birth weight, vaginal birth type, and length of hospitalization were higher in the infected neonates than in the uninfected neonates (Table 1).

A total of 135 pathogens were isolated from 126 blood cultures. The most common organisms isolated from

**Table 1.** Demographic Features of the Infected and Uninfected Groups

	Infected Group, No. (%)	Uninfected Group, No. (%)	P Value
<b>Gender</b>			0.352
Female	60 (47.6)	56 (44.4)	
Male	66 (52.4)	70 (55.6)	
<b>Gestational age, wk</b>			< 0.001
≤ 32	48 (38.1)	26 (20.7)	
33 - 37	22 (17.5)	54 (42.9)	
≥ 38	56 (44.4)	46 (36.4)	
<b>Birth weight, g</b>			< 0.001
< 1000	24 (19)	0 (0)	
1001 - 1500	17 (13.5)	18 (14.3)	
15001 - 2000	16 (12.7)	38 (30.2)	
2001 - 2500	12 (9.5)	22 (17.5)	
> 2500	57 (45.2)	48 (38.1)	
<b>Delivery presentation</b>			0.184
Head Presentation	121 (96)	124 (98.4)	
Breech Presentation	5 (4)	2 (1.6)	
<b>Type of delivery</b>			0.005
Vaginal birth type	33 (26.2)	16 (12.7)	
Caesarean section	93 (73.8)	110 (87.3)	
<b>Length of hospitalization, d</b>			< 0.001
≤ 15	20 (15.9)	79 (62.7)	
16 - 30	28 (22.2)	37 (29.4)	
31 - 60	52 (41.3)	6 (4.8)	
≥ 61	26 (20.6)	4 (3.2)	

blood culture were *S. epidermidis* (20.7%), *Klebsiella* spp. (26.7%), and *Acinetobacter* spp. (13.3%) (Table 2).

The results of the univariate analysis of the risk factors for HAIs in neonates are shown in Table 3.

We selected 10 variables with a P value < 0.05 for the logistic regression model. In the multiple logistic regression analysis, fifth-minute Apgar score, use of erythrocyte transfusion, and surgical operation were found to be the independent risk factors for NIs in patients with a diagnosis of hospital infection (Table 4).

## 5. Discussion

The development of new treatment options and life support techniques in neonatology over the last few decades has increased the survival rate of neonates with low birth weight and preterm infants. However, NIs have

become a major problem in the NICUs because of the prolonged length of hospitalization (12-14). The hospitalization of infants in NICUs has been reported in the literature as being longer, and the susceptibility to infection has been greater than in other pediatric intensive care units (15, 16). Unsurprisingly, we found that low gestational age and lower birth weight placed infants at a higher risk of developing bacterial BSIs. As indicated in other studies, this finding could be due to the patients' immature immune system, long duration of hospitalization, and use of invasive procedures (17-19).

The variety of pathogens most commonly isolated from BSIs may vary from clinic to clinic or region to region (20, 21). Generally, gram-positive organisms are more common than gram-negative organisms in NICUs. The reason may be that gram-positive organisms, such as skin flora, are more likely to colonize the catheter during catheter insertion and use, whereas infections with gram-negative or-

ganisms are more commonly associated with the translocation of flora from the gut or respiratory or urinary tract (17). Djordjevic et al. reported that the most common causative agents isolated from NIs were *Klebsiella-Enterobacter* (39.3%), *Escherichia coli* (25.0%), coagulase-negative *Staphylococci* (16.1%), *Acinetobacter* spp. (10.7%), *Candida albicans* (5.4%), and *Stenotrophomonas maltophilia* (3.6%) (22). Pellizzer et al. reported that the most common causative agents isolated from HCA-BSIs were *Staphylococcus aureus* (17.3%), *Pseudomonas aeruginosa* (15.5%), *Enterococcus* spp. (15.2%), *E. coli* (12.1%), coagulase-negative *Staphylococci* (10.8%), and *Candida* spp. (8.5%) (23). Auriti et al. reported that the most common causative agents isolated from HCA-BSIs were coagulase-negative *staphylococci* (25%), *Klebsiella pneumoniae* (21.7%), *C. albicans* (25%), and *P. aeruginosa* (13%) (24). From Turkey, Tekin et al. found that the most common causative agents isolated from HCA-BSIs were *Acinetobacter baumannii* (48%), *P. aeruginosa* (32%), and *Klebsiella* spp. (6%) (1). We found that the most common causative agents isolated from HCA-BSIs were *S. epidermidis* (20.7%), *Klebsiella* spp. (26.7%), and *Acinetobacter* spp. (13.3%) (Table 2). However, these results indicate that gram-positive organisms, still the most common causative agents isolated from HCA-BSIs, also show that the recently increased incidence of *Acinetobacter* spp. and resistance of *Acinetobacter* spp. may cause serious health problems if the necessary measures are not taken (25, 26).

**Table 2.** Distribution of Nosocomial Pathogens<sup>a</sup>

Microorganism	No. (%)
<b>Gram-Positive Organisms</b>	
<i>Staphylococcus epidermidis</i>	28 (20.7)
<i>Staphylococcus hominis</i>	5 (3.7)
<i>Staphylococcus haemolyticus</i>	5 (3.7)
<i>Staphylococcus aureus</i>	3 (2.2)
Other coagulase-negative <i>staphylococci</i>	8 (5.9)
<i>Enterococcus</i> spp.	3 (2.2)
Other gram-positive organisms	11 (8.1)
<b>Gram-negative organisms</b>	
<i>Klebsiella</i> spp.	36 (26.7)
<i>Acinetobacter</i> spp.	18 (13.3)
<i>Pseudomonas aeruginosa</i>	7 (5.2)
Other gram-negative organisms	2 (1.5)
<i>Candida</i>	9 (6.7)
<b>Total number of isolates</b>	135 (100)

<sup>a</sup>No. = 135 isolates in 126 neonates.

Recent studies have identified several risk factors for NIs in NICUs (9, 27). Pellizzer et al. reported that surgery, antibiotic exposure, antacid medications, leucopenia, chemotherapy, use of steroids, length of hospitalization (> 5 days), ICU admission, tracheostomy, and exposure to indwelling devices were associated with a significantly increased risk of HCA-BSIs by univariate analysis, and that risk factors such as antibiotic exposure, surgical intervention, trauma, central venous catheter, urinary catheter, intubation, tracheostomy, length of hospitalization (15 days), Charlson index, and McCabe classification were independently associated ( $P < 0.05$ ) with HCA-BSIs (23). Consistent with our study, Kishk et al. found that HCA-BSI was more frequent among low birth weight infants than high birth infants (28).

Length of hospitalization, gestational age, birth weight, mechanical ventilation, total parenteral nutrition, umbilical catheter, use of antibiotics, and intubation at birth were associated with a significantly increased risk of HCA-BSIs. Some studies (15, 17, 29) reported that surgery, orogastric tube, central venous catheter, and nasal continuous positive airway pressure were not associated with a significantly increased risk of HCA-BSIs. Djordjevic and Auriti found that low birth weight (< 1500 g), low gestational age at birth (< 32 weeks), first-minute Apgar score, endotracheal intubation at birth, mechanical ventilation and assisted ventilation by continuous positive pressure through nasal prongs, central venous catheter, parenteral nutrition, continuous enteral nutrition, premature rupture of membranes, cesarean section, male sex, aspiration, congenital anomalies, intracranial hemorrhage, and surgical intervention were associated with a significantly increased risk of HCA-BSIs; in the multivariate logistic regression analyses, only low gestational age (< 32 weeks) and central venous catheter were independent risk factors for HAIs (22, 24). In the present study we found that surgical operation, ventriculoperitoneal shunt, use of umbilical catheter, use of nasogastric or orogastric tube, use of urinary catheter, use of mechanical ventilation, surfactant, erythrocyte transfusion, plasma transfusion, thrombocyte transfusion, total parenteral nutrition infusion, intracranial hemorrhage, length of stay, fifth-minute Apgar score, and duration of total parenteral nutrition were all associated with a significantly increased risk of HCA-BSIs using univariate analysis. In the multivariate logistic regression analyses, only the fifth-minute Apgar score, use of erythrocyte transfusion, and surgical operation were independent risk factors for HAIs.

The main limitation of our study is the fact that it is a retrospective study with a relatively small sample size. Studies with more comprehensive analyses and a larger number of patients can provide further data on these vari-

**Table 3.** Comparison of Risk Factors Between the Infected and Uninfected Groups<sup>a</sup>

Risk Factors	Infected group	Uninfected group	P Value <sup>b</sup>
Surgical operation	46 (36.5)	2 (1.6)	< 0.001
Ventriculoperitoneal shunt	21 (16.7)	2 (1.6)	< 0.001
Umbilical catheter	49 (38.9)	20 (15.9)	< 0.001
Nasogastric or orogastric tube	122 (96.8)	113 (89.7)	0.021
Tracheostomy	3 (2.4)	0 (0)	0.124
Urinary catheter	5 (4)	0 (0)	0.030
Mechanical ventilation	97 (77)	32 (25.4)	< 0.001
Use of surfactant	40 (31.7)	20 (15.9)	0.002
Phototherapy	11 (8.7)	10 (7.9)	0.500
Exchange transfusion	3 (2.4)	4 (3.2)	0.491
Erythrocyte transfusion	102 (81)	19 (15.1)	< 0.001
Plasma transfusion	49 (38.9)	9 (7.1)	< 0.001
Thrombocyte transfusion	24 (19)	2 (1.6)	< 0.001
Intravenous immunoglobulin	1 (0.8)	3 (2.4)	0.311
Total parenteral nutrition infusion	105 (83.3)	44 (34.9)	< 0.001
Intracranial hemorrhage	15 (11.9)	4 (3.2)	0.008
Length of hospitalization, d, mean ± SD	43.9 ± 28.9	16.9 ± 14.3	< 0.001
Apgar score (5th minute), mean ± SD	6.7 ± 1.5	7.9 ± 1.3	< 0.001
Total parenteral nutrition, d, mean ± SD	18.5 ± 18.5	2.7 ± 5.8	< 0.001

<sup>a</sup>Values are expressed as No. (%) unless otherwise indicated.  
<sup>b</sup> $\chi^2$  and t distribution.

**Table 4.** Multivariate Logistic Regression Analysis of the Prediction of Independent Risk Factors for Nosocomial Infections

	Unadjusted		Adjusted	
	OR (95% CI)	P Value	OR (95% CI)	P Value
Apgar score, 5th minute	1.16 (0.79 - 1.53)	< 0.001	1.42 (1.03 - 1.96)	0.030
Mechanical ventilation	0.10 (0.05 - 0.18)	< 0.001	0.42 (0.14 - 1.24)	0.119
Erythrocyte transfusion	0.04 (0.02 - 0.08)	< 0.001	0.07 (0.31 - 0.19)	< 0.001
Plasma transfusion	0.12 (0.05 - 0.26)	< 0.001	1.50 (0.47 - 4.71)	0.487
Thrombocyte transfusion	0.06 (0.01 - 0.29)	< 0.001	0.17 (0.02 - 1.00)	0.050
Use of surfactant	0.40 (0.22 - 0.74)	0.003	2.84 (0.66 - 12.0)	0.157
Ventriculoperitoneal shunt	0.08 (0.18 - 0.35)	< 0.001	4.79 (0.50 - 45.7)	0.173
Umbilical catheter	0.29 (0.16 - 0.53)	< 0.001	0.40 (0.11 - 1.47)	0.173
Surgical operation	0.02 (0.00 - 0.11)	< 0.001	0.01 (0.02 - 0.07)	< 0.001
Nasogastric or orogastric tube	0.28 (0.09 - 0.90)	0.024	1.31 (0.22 - 7.72)	0.760

ables.

### 5.1. Conclusion

This study identified the risk factors associated with HAIs in our NICUs that are necessary for HCA-BSIs prevention programs and found that infection control is a signif-



icant problem in NICUs. We suggest that preventive strategies for BSI in neonates in NICUs should continue to focus on limiting the use of invasive devices to reduce HAIs.

## Footnotes

**Authors' Contribution:** Sabahattin Ertugrul and Fesih Aktar developed the study concepts and design. Ilyas Yolbas, Ahmet Yildirim, and Kamil Yilmaz collected the data. Ahmet Yilmaz and Recep Tekin conducted the statistical analysis and interpreted the data. Sabahattin Ertugrul and Fesih Aktar wrote the manuscript. Bilal Elbey revised and edited the manuscript. All authors read and approved the final manuscript.

**Conflict of Interest:** The authors declare no conflict of interest.

**Funding/Support:** This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

## References

1. Tekin R, Dal T, Pirinccioglu H, Oygucu SE. A 4-year surveillance of device-associated nosocomial infections in a neonatal intensive care unit. *Pediatr Neonatol*. 2013;54(5):303-8. doi: [10.1016/j.pedneo.2013.03.011](https://doi.org/10.1016/j.pedneo.2013.03.011). [PubMed: 23643153].
2. Correa L, Pittet D. Problems and solutions in hospital-acquired bacteraemia. *J Hosp Infect*. 2000;46(2):89-95. doi: [10.1053/jhin.2000.0803](https://doi.org/10.1053/jhin.2000.0803). [PubMed: 11049700].
3. Hufnagel M, Burger A, Bartelt S, Henneke P, Berner R. Secular trends in pediatric bloodstream infections over a 20-year period at a tertiary care hospital in Germany. *Eur J Pediatr*. 2008;167(10):1149-59. doi: [10.1007/s00431-007-0651-4](https://doi.org/10.1007/s00431-007-0651-4). [PubMed: 18231812].
4. Kaufman D, Fairchild KD. Clinical microbiology of bacterial and fungal sepsis in very-low-birth-weight infants. *Clin Microbiol Rev*. 2004;17(3):638-80. doi: [10.1128/CMR.17.3.638-680.2004](https://doi.org/10.1128/CMR.17.3.638-680.2004). [PubMed: 15258097].
5. Kawagoe JY, Segre CA, Pereira CR, Cardoso MF, Silva CV, Fukushima JT. Risk factors for nosocomial infections in critically ill newborns: a 5-year prospective cohort study. *Am J Infect Control*. 2001;29(2):109-14. doi: [10.1067/mic.2001.114162](https://doi.org/10.1067/mic.2001.114162). [PubMed: 11287879].
6. Mehta Y, Gupta A, Todi S, Myatra S, Samaddar DP, Patil V, et al. Guidelines for prevention of hospital acquired infections. *Indian J Crit Care Med*. 2014;18(3):149-63. doi: [10.4103/0972-5229.128705](https://doi.org/10.4103/0972-5229.128705). [PubMed: 24701065].
7. Stoll BJ, Hansen NI, Adams-Chapman I, Fanaroff AA, Hintz SR, Vohr B, et al. Neurodevelopmental and growth impairment among extremely low-birth-weight infants with neonatal infection. *JAMA*. 2004;292(19):2357-65. doi: [10.1001/jama.292.19.2357](https://doi.org/10.1001/jama.292.19.2357). [PubMed: 15547163].
8. Balkhy HH, Alsaif S, El-Saed A, Khawajah M, Dichinee R, Memish ZA. Neonatal rates and risk factors of device-associated bloodstream infection in a tertiary care center in Saudi Arabia. *Am J Infect Control*. 2010;38(2):159-61. doi: [10.1016/j.ajic.2009.09.004](https://doi.org/10.1016/j.ajic.2009.09.004). [PubMed: 19900739].
9. Couto RC, Pedrosa TM, Tofani Cde P, Pedroso ER. Risk factors for nosocomial infection in a neonatal intensive care unit. *Infect Control Hosp Epidemiol*. 2006;27(6):571-5. doi: [10.1086/504931](https://doi.org/10.1086/504931). [PubMed: 16755475].
10. Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. *Am J Infect Control*. 1988;16(3):128-40. [PubMed: 2841893].
11. Gray JW. Surveillance of infection in neonatal intensive care units. *Early Hum Dev*. 2007;83(3):157-63. doi: [10.1016/j.earlhumdev.2007.01.006](https://doi.org/10.1016/j.earlhumdev.2007.01.006). [PubMed: 17289308].
12. Jeong IS, Jeong JS, Choi EO. Nosocomial infection in a newborn intensive care unit (NICU), South Korea. *BMC Infect Dis*. 2006;6:103. doi: [10.1186/1471-2334-6-103](https://doi.org/10.1186/1471-2334-6-103). [PubMed: 16796741].
13. Brito DV, Brito CS, Resende DS, Moreira do OJ, Abdallah VO, Gontijo Filho PP. Nosocomial infections in a Brazilian neonatal intensive care unit: a 4-year surveillance study. *Rev Soc Bras Med Trop*. 2010;43(6):633-7. [PubMed: 21181013].
14. Pessoa-Silva CL, Richtmann R, Calil R, Santos RM, Costa ML, Frota AC, et al. Healthcare-associated infections among neonates in Brazil. *Infect Control Hosp Epidemiol*. 2004;25(9):772-7. doi: [10.1086/502475](https://doi.org/10.1086/502475). [PubMed: 15484803].
15. Nagata E, Brito AS, Matsuo T. Nosocomial infections in a neonatal intensive care unit: incidence and risk factors. *Am J Infect Control*. 2002;30(1):26-31. [PubMed: 11852413].
16. National Nosocomial Infections Surveillance System. National Nosocomial Infections Surveillance (NNIS) System Report, data summary from January 1992 through June 2004, issued October 2004. *Am J Infect Control*. 2004;32(8):470-85. doi: [10.1016/S0196655304005425](https://doi.org/10.1016/S0196655304005425). [PubMed: 15573054].
17. Perlman SE, Saiman L, Larson EL. Risk factors for late-onset health care-associated bloodstream infections in patients in neonatal intensive care units. *Am J Infect Control*. 2007;35(3):177-82. doi: [10.1016/j.ajic.2006.01.002](https://doi.org/10.1016/j.ajic.2006.01.002). [PubMed: 17433941].
18. Rojas MA, Efirid MM, Lozano JM, Bose CL, Rojas MX, Rondon MA, et al. Risk factors for nosocomial infections in selected neonatal intensive care units in Colombia, South America. *J Perinatol*. 2005;25(8):537-41. doi: [10.1038/sj.jp.7211353](https://doi.org/10.1038/sj.jp.7211353). [PubMed: 16047032].
19. Contreras-Cuellar GA, Leal-Castro AL, Prieto R, Carvajal-Hermida AL. Device-associated infections in a Colombian neonatal intensive care unit. *Rev Salud Publica (Bogota)*. 2007;9(3):439-47. [PubMed: 18026608].
20. Couto RC, Carvalho EA, Pedrosa TM, Pedroso ER, Neto MC, Biscione FM. A 10-year prospective surveillance of nosocomial infections in neonatal intensive care units. *Am J Infect Control*. 2007;35(3):183-9. doi: [10.1016/j.ajic.2006.06.013](https://doi.org/10.1016/j.ajic.2006.06.013). [PubMed: 17433942].
21. Cimiotti JP, Haas J, Saiman L, Larson EL. Impact of staffing on bloodstream infections in the neonatal intensive care unit. *Arch Pediatr Adolesc Med*. 2006;160(8):832-6. doi: [10.1001/archpedi.160.8.832](https://doi.org/10.1001/archpedi.160.8.832). [PubMed: 16894083].
22. Djordjevic ZM, Markovic-Denic L, Folic MM, Igrutinovic Z, Jankovic SM. Health care-acquired infections in neonatal intensive care units: risk factors and etiology. *Am J Infect Control*. 2015;43(1):86-8. doi: [10.1016/j.ajic.2014.10.005](https://doi.org/10.1016/j.ajic.2014.10.005). [PubMed: 25564130].
23. Pellizzer G, Mantoan P, Timillero L, Allegranzi B, Fedeli U, Schievano E, et al. Prevalence and risk factors for nosocomial infections in hospitals of the Veneto region, north-eastern Italy. *Infection*. 2008;36(2):112-9. doi: [10.1007/s15010-007-7092-x](https://doi.org/10.1007/s15010-007-7092-x). [PubMed: 18327681].
24. Auriti C, Maccallini A, Di Liso G, Di Ciommo V, Ronchetti MP, Orzalesi M. Risk factors for nosocomial infections in a neonatal intensive-care unit. *J Hosp Infect*. 2003;53(1):25-30. [PubMed: 12495682].
25. El-Ageery SM, Abo-Shadi MA, Alghaithy AA, Ahmad MA, Alsharif NH, Alharbi SA. Epidemiological investigation of nosocomial infection with multidrug-resistant *Acinetobacter baumannii*. *Eur Rev Med Pharmacol Sci*. 2012;16(13):1834-9. [PubMed: 23208968].
26. Tekin R, Dal T, Bozkurt F, Devenci O, Palanc Y, Arslan E, et al. Risk factors for nosocomial burn wound infection caused by multidrug resistant *Acinetobacter baumannii*. *J Burn Care Res*. 2014;35(1):73-80. doi: [10.1097/BCR.0b013e31828a493f](https://doi.org/10.1097/BCR.0b013e31828a493f). [PubMed: 23799478].
27. Duenas L, Bran de Casares A, Rosenthal VD, Jesus Machuca L. Device-associated infections rates in pediatrics and neonatal intensive

- care units in El Salvador: findings of the INICC. *J Infect Dev Ctries.* 2011;**5**(6):445-51. [PubMed: [21727643](#)].
28. Kishk RM, Mandour MF, Farghaly RM, Ibrahim A, Nemr NA. Pattern of Blood Stream Infections within Neonatal Intensive Care Unit, Suez Canal University Hospital, Ismailia, Egypt. *Int J Microbiol.* 2014;**2014**:276873. doi: [10.1155/2014/276873](#). [PubMed: [25389439](#)].
29. Aktar F, Tekin R, Gunes A, Ulgen C, Tan I, Ertugrul S, et al. Determining the Independent Risk Factors and Mortality Rate of Nosocomial Infections in Pediatric Patients. *Biomed Res Int.* 2016;**2016**:7240864. doi: [10.1155/2016/7240864](#). [PubMed: [26981536](#)].