











- before and after lymphocyte immunotherapy. *Am J Reprod Immunol.* 2012;68(2):164-74.
- 42- Liu YS, Wu L, Tong XH, Wu LM, He GP, Zhou GX, et al. Study on the relationship between Th17 cells and unexplained recurrent spontaneous abortion. *Am J Reprod Immunol.* 2011;65(5):503-11.
- 43- Messaoudi S, Al-Khateeb GM, Dendana M, Sater MS, Jazia KB, Nourira M, et al. Genetic variations in the interleukin-21 gene and the risk of recurrent idiopathic spontaneous miscarriage. *Eur Cytokine Netw.* 2011;22(2):123-6.
- 44- Saifi B, Rezaee SA, Tajik N, Ahmadpour ME, Ashrafi M, Vakili R, et al. Th17 cells and related cytokines in unexplained recurrent spontaneous miscarriage at the implantation window. *Reprod Biomed Online.* 2014;29(4):481-9.
- Intravenous immunoglobulin G modulates peripheral blood Th17 and Foxp3(+) regulatory T cells in pregnant women with recurrent pregnancy loss. *Am J Reprod Immunol.* 2014;71(5):441-50.
- 39- Bansal AS, Bajardeen B, Thum MY. The basis and value of currently used immunomodulatory therapies in recurrent miscarriage. *J Reprod Immunol.* 2012;93(1):41-51.
- 40- Gharesi Fard B, Zolghadri J, Kamali Sarvestani E. Effect of leukocyte therapy on tumor necrosis factor-alpha and interferon-gamma production in patients with recurrent spontaneous abortion. *Am J Reprod Immunol.* 2008;59(3):242-50.
- 41- Liang P, Mo M, Li GG, Yin B, Cai J, Wu T, et al. Comprehensive analysis of peripheral blood lymphocytes in 76 women with recurrent miscarriage