Mucor	
Mucor	
Mucor circinelloides f.janssenii	Mucor hiemalis f.silvaticus
Mucor plumbeus	Mucor circinelloides f.lusitanicus
•	
	•
	Mucor:

. . . .( )
Mucor

PDA

.( )

.

.( ).

%

.( )

. Mucor

Mucor hiemalis f.silvaticus

| BSA .()

Mucor circinelloides f.janssenii
|
| SAS (V.9.1)

Mucor circinelloides f.lusitanicus
|
| Mucor plumbeus

.()

(g/lit) (mg/lit) Mucor circinelloides f.circinelloides Е Mucor circinelloides f.janssenii EF AB Mucor circinelloides f.lusitanicus CDA Mucor hiemalis f.corticola В EF Mucor hiemalis f.silvaticus CD Α

C

Е

BC

.( )

Mucor plumbeus

Mucor racemosus f.chibinensis


Mucor strictus

F


C

CD

.( )



Mucor circinelloides f.lusitanicus Mucor hiemalis f.silvaticus



Mucor circinelloides f.lusitanicus Mucor hiemalis f.silvaticus

.....

## Archive of SID

.  Mucor hiemalis f.silvaticus	рН
	.( )
. Mucor circinelloides f.lusitanicus	.( )
	.( )
.( )	
	pH)

1. Bradford, M.M. 1976. A rapid and sensitive method for quantification of microgram quantities of protein of utilizing the principle dye binding. Anal. Biochemistry. 72: 680-685.

- 2. Gupta, S., M. Kapoor, K. Kant Sharma, L.M. Nair and R. Chander Kuhad. 2008. Production and recovery of an alkaline exo-polygalacturonase from *Bacillus subtilis* RCK under solid-state fermentation using statistical approach. Bioresource Technology. 99: 937-945.
- 3. Hadj-Taieb, N., M. Ayadi, M. Khlif, K. Mrad and I. Hassairi. 2006. Fermentor production of pectinases on gruel, a local by-product and their use in olive oil extraction. Enzyme and Microbial Technology. 39: 1072-1076.
- 4. Jayani, R.S., S. Saxena and R. Gupta. 2005. Microbial pectinolytic enzymes: A review. Process Biochemistry. 40: 2931-2944.
- 5. Kossem, A. and P. Nannipieri. 1995. Soil cellulase activity methods. In: Kossem A, Nannipieri P (eds) Applied Soil Microbial and Biochnology. Academic Press: San Diego, pp: 345-350.
- 6. Patil, S.R. and A. Dayanand. 2006. Exploration of Regional Agrowastes for the Production of Pectinase by *Aspergillus niger*. Food Technology. Biotechnology. 44(2): 289-292.
- 7. Silva, D., E.M. Martin, R. Silva and E. Gomes. 2002. Pectinase production by *penicillium viridicatum* RFC3 by solid state fermentation using agricultural wastes and agro-industrial by-products. Brazil Journal of Microbiology. 33: 318-324.
- 8. Soares. M., R. Silva and E. Gomes. 1999. Screening of bacterial strains for pectinolytic activity: characterization of the polygalacturonase producted by *bacillus* sp. Revista de Microbiology. 30: 299-303.
- 9. Souza. J., E. Silva, M. Maia and M. Teixeira. 2003. Screening of fungal strains for pectinolytic activity: endopolygalacturonase production by Peacilomyces clav isporus 2A.UMIDA.1. Process Biochemistry. 39: 455-458.
- 10. Tari. C., N. Dogan and N. Gogus. 2008. Biochemical and thermal characterization of crude exo-polygalacturonase produced by *Aspergillus sojae*. Food Chemistry. 111: 824-829.

.....

Archive of SID

# A Study on Released Sugars From Pectin and Related Proteins By Several Mocur

M.H. Maleki<sup>1</sup>, G.A. Ranjbar<sup>2</sup>, M.A. Tajick<sup>2</sup>, A. Asgharzadeh<sup>3</sup> and A. Lotfi<sup>4</sup>

#### **Abstract**

The present study was aimed to screen the pectinase activity from some genera of *Mocur* in laboratory conditions. Pectin was used as the only carbon source in a minimal culture medium. Tow days after inoculation released proteins and sugars were assayed with related reagents and repeated each 2 days up to 21<sup>th</sup> day. Statistical analysis showed significant variation in released sugars and released proteins among tested genera. *Mucor hiemalis silvaticus* had highest and *Mucor circinelloides janssenii* had lowest sugar levels. Also *Mucor circinelloides* f. *lusitanicus* had highest and *Mucor plumbeus* had lowest protein levels. Glucose and protein levels for superior Species have increased until 15 and 13 days, respectively for glucose and protein after inoculation, then decreased until 25<sup>th</sup> day, but had no variation until 30<sup>th</sup> day. These results showed that isolates belong to the same forms had no significance difference in pectinase activity.

Keywords: Mocur, Pectin, Spectrophotometer, Pectinase activity

<sup>1-</sup> Former M. Sc. Student, Sari Agricultural Sciences and Natural Resources University

<sup>2-</sup> Assistant Professor, Sari Agricultural Sciences and Natural Resources University

<sup>3-</sup> Associate Professor, Soil and Water Research Institute, Tehran

<sup>4-</sup> Instructor, Sari Agricultural Sciences and Natural Resources University