
PCA

(// : // :)

()

(*)

/

(

) TWINSPAN

(

)

...

-

-

.()

.()

()

()

()

() ()

.()

.()

.()

() ()

.()

)

(

°

"

°

"

.()

°

"

°

"

Ecological Species Group

Michigan

Wisconsin

Spies and Barnes

Barnes, et al.

.()

PCA () () () *

()

TWINSpan

PCA Windows PC-ORD

() ()

()

TWINSpan (PCA)

() ()

() ()


()

)

Menta aquatica (

PCA (Principal Component Analysis)
 Guadal
 Pearson

Carex grioleti ()
Oplismenus undulatifolius ()
 A,B,C,D,E
Brachypodium pinnatum ()
 () *Hedera pastuchovii* ()
Ruscus () *Euphorbia amygdaloides*
hyrcanus
Rubus ()
Viola odorata ()
caesius



TWINSPAN

...

<i>Carpinus betulus</i> L.	<i>Menta aquatica</i> L.	A
<i>Parrotia persica</i> (DC.) C. A. Mey. - <i>Ruscus hyrcanus</i> L.	<i>Hedera pastuchovii</i> L.- <i>Oplismenus undulatifolius</i> (AC.)	B
<i>Parrotia persica</i> (DC.) C. A. Mey. - <i>Ruscus hyrcanus</i> L.	L. <i>Hedera Carex grioletia</i> L. <i>Pastuchivii</i>	C
<i>Parrotia persica</i> (DC.) C. A. Mey. - <i>Cratagus</i> SP.	<i>Brachypodium pinnatum</i> L. - <i>Violaodorata</i> L.	D
<i>Parrotia persica</i> (DC.) C. A. Mey. - <i>Cratagus</i> SP. - <i>Quercus castaneifolia</i> C.	<i>Brachypodium pinnatum</i> L. - <i>Rubus caesius</i> L.	E

PCA

PCA

()

PCA

Clayb CECc Na Ca

Clayc

PCA

()

Bec Nec

(

Sandc Sandb Nlit Wc Wb pHc pHb pHa


Claya Pa

()


)

PCA

a,b,c			
pHa, pHb, pHc			
Wa, Wb, Wc			
Spa, Spb, Spc			
Eca, Ecb, Ecc	Ds/m		
Ca, Cb, Cc			
Na, Nb, Nc			
C/Na, C/Nb, C/Nc			
CECa, CECb, CECc	p.p.m		
Pa, Pb, Pc	p.p.m		
Sanda, Sandb, Sandc			
Silta, Siltb, Siltc			
Claya, Clayb, Clayc			
La, Lb, Lc			
Nea, Neb, Nec			
Bea, Beb, Bec	gr		
Clit			
Nlit			
C/Nlit			
Alt	m		
Slope			
Aspect			
Sloasp			



PCA



PCA

...

()

()

()

()

()

() ()

()

()

()

()

()

Doweling

()

(PCA)

)

)

(

(

Menta aquatica

()

pHc pHb pHa Bec Nec

Sandc Sandb Nlit Wc Wb

Hedera pastuchovii

Ruscus hyrcanus Euphorbia amygdaloides

Oplismenus undulatifolius

Claya Pa

Euphorbia Hedera pastuchovii

Carex Ruscus hyrcanus amygdaloides

grioleti

Clayb Na Ca

Clayc

Viola Brachypodium pinnatum

odorata

Clayb Na Ca

CECc Clayc

CECc

Brachypodium

Rubus caesius pinnatum

()

()

()

()

Archive of SID

-
- 18-Abdi, H., 2003. Multivariate analysis, Encyclopedia of social science, The University of Texas Dalas. 1-11.
- 19-Archambault I. 1990. Landscape ecosystems of disturbed oak forest of southern Michigan, USA. *Can J. For Res*, 20: 12790-1582.
- 20-Baldock, J. A. and Oades, L. 1992. Aspects of the chemical structure of soil organic materials as revealed by solid – state. *Soil Biology and Biochemistry*. 16: 1 - 42.
- 21-Barnes B. V. 1998. *Forest Ecology* (4Th). John Wiley & Sons, Inc. 774pp.
- 22-Barnes, B.V., Pregitzer K.S., Spies T.A. 1982. Ecological forest site classification, *J.For.* 80:493-498.
- 23-Burton, V., Barnes, D. E., Shirley, K. R., Stephen, H. and Sandspur, H. 1998. *Forest Ecology*. John Willey and Sons, 609 - 610pp.
- 24-Cain, P. D., 1959. *Manual of vegetation analysis*. Harper and Row, New York. 325pp.
- 25-Coker, P. D., 2000. *Vegetation mapping: From patch to planet*, edited by: Alexander R., and Millington A.C., John Wiley and Sons publication, 135:158.
- 26-Crowley, W., Harrison, S.S.C., Coroi, M., and Sacre, V.M., 2003. An ecological assessment of the plant communities at Port Ban nature reserve in south-western Ireland. *Biology and Environmental: Proceeding of Royal Irish Academy*, 1038 (2): 69-82.
- 27-Dombois, D.M., and Ellenberg, H., 1974. *Aims and methods of vegetation ecology*. John Wiley and Sons Published New York, London, Sydney, Toronto, 547pp.
- 28-Doweling, A. Y., Webb, A. A. and Scanlan, Y. C. 1986. Surface soil chemical and physical patterns in a brig low – Dawson gom forest central careens land. *Journal of Ecology*, 12: 155 -182.
- 29-Ewald, J. 1999. Soziologie und standortbindung subalpiner fichtenwalder in den bayerischen Alpen. *Tuexenia* 19: 107-125.
- 30-Ewald, J. 2000. The influence of coniferous canopies on under story vegetation and soils in mountain forests of the northern calcareous Alps. *Appl. Veg. Sci.* 3: 123-134.
- 31-Fisher, R., Binkley, D. 2000. *Ecology and Management of forest soils* (Third edition), John Wiley & Sons, INC, 489 pp.
- 32-Grant, C. D. and Loneragan, W.A.2001. The effects of burning on the under story composition of rehabilitated bauxite mines bin Western Australia: community changes vegetation succession. *Ecology and Management*. 145: 255-277.
- 33-Greiege-Smith, P., 1983. *Quantitative plant ecology*. University of California press. Berkeley, Calif., 359pp.
- 34-Hedman, C.W., Grace, S.L. and Ling, S.E., 2000. Vegetation composition and structure of southern coastal plain pine forests: An ecological comparison. *Forest ecology and Management*. 134: 233-247.
- 35-Hix, M.O. 1994. DECORANA and TWINSPLAN for ordination and classification of multivariate species data: a new edition, together with supporting programs, in FORTRAN 77. Institute of Terrestrial Ecology, Huntingdon, 58pp.
- 36-Kooch, Y., Jalilvand, H., Bahmanyar, M. A. and Pormajidian, M. R. 2007. Ecological distribution of Indicator species and effective edaphical factors on the northern Iran lowland forests, *Journal of Applied Science*. 7: 1475 -1483.
- 37-Oland, J.M., Loidi, J., Ander G. and Escudero, A.1998. Relating variation in the under story of beech forests to ecological factors. *Folia Geobot*. 33: 77-86.

...

38-Pregitzer, K.S. and Barnes, B.V., 1984. Classification and comparison of the upland hardwood and conifer ecosystems of the Cyrus H. McCormick Experimental Forest. Upper Peninsula, Michigan. Canadian journal of forestry Research, 14:362-375.

39-Schoenholtz, S.H, Van Miegroet, H, Burger, J.A. 2000. A review of chemical and physical properties as indicators of forest soil quality: challenges and properties. Forest ecology and management, 138: 335-356.

40-Zahedi Amiri, GH. 1998. Relation between ground vegetation and soil characteristics in a mixed hardwood stand, PhD. thesis, university of Ghent, begum academic press, 319 pp.

Archive of SID

Differentiation of ecosystem units of Caspian lowland forests and its relation with some soil characteristics

Y. Kooch^{*1}, H. Jalilvand², M. A. Bahmanyar³ and M. R. Pormajidian²

¹ Ph.D. Candidate of Forestry, Faculty of Natural Resources, Tarbiat Modarres University, Noor, I.R.Iran

² Assistant Prof., University of Natural Resources and Agriculture Sciences of Sari, Sari, I.R.Iran

³ Associate Prof., University of Natural Resources and Agriculture Sciences of Sari, Sari, I.R.Iran

(Received: 13 May 2008, Accepted: 27 November 2008)

Abstract

Due to differentiation ecosystem units and related soil physico - chemical and biological characteristics, 268.7 ha⁻¹ of lowland forests in Khanikan were studied. Vegetation data and information (trees, shrubs, and herbs) were collected from 60 sample plots with systematically random method (20m×20m) which were then analyzed by using TWINSpan program. The data were classified into five ecological groups. Some of physical, chemical and biological characteristics of soil, such as soil acidity (pH), bulk density, saturation moisture, electrical conductivity (EC), organic carbon, total nitrogen, cation exchange capacity, available phosphorous, soil texture, lime, biomass of earthworms, litter carbon, and litter nitrogen were measured. Principal component analysis (PCA) was used to determine correlation as well as distribution in each ecological group of environmental factors. It was noticed that 1st, 3rd, 4th ecological groups had the highest correlation with the 1st axes whereas 2nd and 5th ecological groups demonstrated the highest correlation with the 2nd axes. The results indicated that the ecological group's distribution pattern was mainly related to soil characteristics such as pH, bulk density, texture, phosphorous, organic carbon, nitrogen, and CEC.

Keywords: Ecological species group, Classification, PCA, Soil characteristics, Caspian forests, Iran

*Corresponding author: Tel: +98 911 2932313 , Fax: +98 152 4222982 , E-mail: yahya_kooch@yahoo.com