
*

(I I : II :)

(*Buxus hyrcana* Pojark.)

×

Hypericum Pteris cretica L.

Atropa belladonna L. *Athyrium flix-femina* (L.) Roth *Carex sylvatica* L. *Rubus hyrcanus* Woron. *androsaemum*
| | | *Diospyrus lotus* L.

(*Buxus hyrcana* Pojark)

(
(Esmailzadeh et al., 2011 b)

(Jalili &

Jamzad, 1999)

(Esmailzadeh et al., 2011 a)
Grime and Thompson (1979)

:

:

(Bakker et al.,

1996)

(Levin, 1990)

(Bakker, 1989; Baskin

& Baskin, 1998)

(Thompson, 2000)

(Thompson & Grime, 1979)

(Thompson et al., 1997)

(Baskin & Baskin, 1998)

(Esmailzadeh et al., 2010 a)

(Stark et

al., 2008)

² Transient
³ Persistent

¹ Soil seed bank

/

(Bossuyt & Honnay,
et al., Yan et al. (2009) .2008)
(2009) Chaideftou (2009)

(Anonymous,
.1998)

Jalili et al.,
Esmailzadeh et al., (2010 b) (2003)

(Barbour et al., 1987)

()

×)
(Barnes, 1998) (

)
(

° ° ° °

¹ Floristic-Physiognomic



×

(Baskin

.& Baskin, 1998)

(Leckie et al., 2000)

(Godefroid et al., 2006)

(Augusto et al., 2001)

(Leckie et al., 2000)

(Esmailzadeh et al., 2010 b)

(Fourie, 2008)

(Esmailzadeh et al., 2011

(Baskin &

.b)

.Baskin, 1998)

(Baskin & Baskin, 1998)

¹ Seedling emergence method

(Diaz-Villa et al., 2003)

(Raunkaier, 1934)

(Greig-Smith, 1983)

Archive of SID

Carex sylvatica (*Rubus hyrcanus* Woron.)

(*Athyrium flix- femina* (L.) Roth) L.

Atropa belladonna L.

(*Diospyrus lotus* L.)

/ / /


¹ Jaccard Similarity Index

...


		()
Adiantaceae			
<i>Adiantum capillus-veneris</i> L.	He	/	
Asteraceae			
<i>Conyza canadensis</i> (L.) Cronq.	Th	/	/
<i>Circaea lutetiana</i> L.	He	/	/
<i>Artemisia annua</i> L.	Th	/	/
<i>Sigesbeckia orientalis</i> L.	Th	/	/
<i>Willemetia tuberosa</i>		/	/
Athyriaceae			
<i>Athyrium flix-femina</i> (L.) Roth.	Cry	/	/
Betulaceae			
<i>Alnus subcordata</i> C. A. Mey.	Ph	/	/
Boraginaceae			
<i>Lithospermum officinale</i> L.	He	/	/
Caprifoliaceae			
<i>Sambucus ebulus</i> L.	Cry	/	/
Cruciferae			
<i>Capsella bursa-pastoris</i> (L.) Medicus	Th	/	/
<i>Cardamine impatiens</i> L.	He		/
<i>Cardamine hirsuta</i> L.	Th	/	/
Cyperaceae			
<i>Carex riparia</i> Curtis	Cry	/	/
<i>Carex sylvatica</i> L.	Cry	/	/
<i>Carex</i> sp.	Cry	/	/
Ebenaceae			
<i>Diospyrus lotus</i> L.	Ph		/
Ephorbiaceae			
<i>Euphorbia amygdaloides</i> L.	He		/
Equisetaceae			
<i>Equisetum maximum</i> Lam.	Cry	/	/
Mimosaceae			
<i>Albizia julibrissin</i> Durraz.	Ph	/	/
Moraceae			
<i>Ficus carica</i> L.	Ph	/	/
<i>Morus alba</i> L.	Ph	/	/
Poaceae			
<i>Poa bulbosa</i> L.	Cry	/	/
<i>Brachypodium pinnatum</i> (L.) P. Beauv.	He	/	/
<i>Oplismenus undulatifolius</i> (Ard.) P. Beauv.	He	/	/
Hypericaceae			
<i>Hypericum androsaemum</i> L.	Ch		/
<i>Hypericum perforatum</i> L.	He		/
Juncaceae			
<i>Juncus</i> sp.	Cry	/	/

			()
Lamiaceae				
<i>Clinopodium umbrosum</i> C. Koch	He	/	/	
<i>Lamium album</i> L.	He	/	/	
<i>Lycopus europaeus</i> L.	He	/	/	
<i>Mentha aquatica</i> L.	He		/	
Phytolaccaceae				
<i>Phytolacca americana</i> L.	He	/	/	
Polygonaceae				
<i>Rumex acetosella</i> L.	He	/	/	
Portulacaceae				
<i>Portulaca oleracea</i> L.	Th	/	/	
Primulaceae				
<i>Cyclamen coum</i> Miller	Cry	/	/	
<i>Primula heterochroma</i> Stapf	He	/	/	
Pteridaceae				
<i>Pteris cretica</i> L.	Cry			
<i>Pteridium aquilinum</i> (L.) Kuhn.	Cry		/	
Rosaceae				
<i>Rubus hyrcanus</i> Woron.	Ph		/	
Scrophulariaceae				
<i>Digitalis nervosa</i> Steud. & Hochst. ex	Th	/	/	
<i>Scrophularia vernalis</i> L.	He	/	/	
Solanaceae				
<i>Atropa belladonna</i> L.	He	/	/	
<i>Solanum kieseritzkii</i> C. A. Mey.	Ch	/	/	
<i>Solanum nigrum</i> L.	Th		/	
Typhaceae				
<i>Typha latifolia</i> L.	Cry	/	/	
Ulmaceae				
<i>Ulmus glabra</i> Hudson	Ph	/	/	
Urticaceae				
<i>Urtica dioica</i> L.	He	/	/	
Verbenaceae				
<i>Verbena officinalis</i> L.	He	/	/	
Violaceae				
<i>Viola alba</i> L.	He	/	/	
Vitaceae				
<i>Vitis sylvestris</i> Gmelin	Ph	/	/	

(=Th =He =Cry =Ch =Ph)



(=E =Th =He =Cry =Ch =Ph)



<i>Adiantum capillus-veneris</i> L.	<i>Alnus subcordata</i> C. A. May.	<i>Acer cappadocicum</i> Gled.
<i>Albizzia julibrissin</i> Durraz.	<i>Athyrium flix- femina</i> (L.) Roth.	<i>Acer velutinum</i> Boiss.
<i>Anonymous</i>	<i>Brachypodium pinnatum</i> (L.) P.	<i>Asplenium adiantum- nigrum</i>
<i>Artemisia annua</i> L.	<i>Cardamine impatiens</i> L.	<i>Buxus hyrcana</i> Pojark.
<i>Atropa belladonna</i> L.	<i>Carex riparia</i> Curtis	<i>Carex remota</i> L.
<i>Capsella bursa-pastoris</i> (L.) Medicus	<i>Carex sylvatica</i> L.	<i>Carex pendula</i> Huds.
<i>Cardamine</i> sp.	<i>Circaea lutetiana</i> L.	<i>Carpinus betulus</i> L.
<i>Carex</i> sp.	<i>Diospyrus lotus</i> L.	<i>Cephalanthera caucasica</i>
<i>Clinopodium umbrosum</i> C. Koch	<i>Equisetum maximum</i> Lam.	<i>Cerasus avium</i> (L.) Moench
<i>Conyza Canadensis</i> (L.) Cronq.	<i>Euphorbia amygdaloides</i> L.	<i>Crataegus microphylla</i> C.
<i>Cyclamen coum</i> Miller	<i>Ficus carica</i> L.	<i>Danae racemosa</i> (L.) Moench
<i>Digitalis nervosa</i> Steud. & Hochst. ex	<i>Hypericum androsaemum</i> L.	<i>Dryopetris borrieri</i> Newm.
<i>Hypericum perforatum</i> L.	<i>Lamium album</i> L.	<i>Dryopetris dilatata</i> (Hoffm.) A. Grav
<i>Juncus</i> sp.	<i>Oplismenus undolatifolius</i> (Ard.) P. Beauv.	<i>Epipactis helleborine</i> (L.) Crantz
<i>Lithospermum officinale</i> L.	<i>Primula heterochroma</i> Stapf.	<i>Evonymus latifolia</i> (L.) Mill.
<i>Lycopus europaeus</i> L.	<i>Pteris cretica</i> L.	<i>Fagus orientalis</i> Lipsky
<i>Mentha aquatica</i> L.	<i>Rubus hyrcanus</i> Woron.	<i>Frangula alnus</i> Miller
<i>Morus alba</i> L.	<i>Solanum kieseritzkii</i> C. A. May.	<i>Fragaria vesca</i> L.
<i>Poa bulbosa</i> L.	<i>Ulmus glabra</i> Hudson	<i>Galium oduratum</i> (L.). Scop.
<i>Phytolacca Americana</i> L.	<i>Viola alba</i> L.	<i>Hedera pastuchovii</i> Woron. Ex Grossh.
<i>Portulaca oleracea</i> L.		<i>Ilex spinigera</i> (Loes) Loes
<i>Pteridium aquilinum</i> (L.) Kuhn.		<i>Laurocerasus officinalis</i>
<i>Rumex acetosella</i> L.		<i>Mespilus germanica</i> L.
<i>Sambucus ebulus</i> L.		<i>Parotia persica</i> (DC.) C. A.
<i>Scrophularia vernalis</i> L.		<i>Petasites hybridus</i> (L.) P.
<i>Sigesbeckia orientalis</i> L.		<i>Polypodium vulgare</i> L.
<i>Solanum nigrum</i> L.		<i>Polystichum aculeatum</i> (L.)
<i>Typha latifolia</i> L.		<i>Polystichum woronowii</i>
<i>Urtica dioica</i> L.		<i>Quercus qastaneaefolia</i> C. A.
<i>Verbena officinalis</i> L.		<i>Ruscus hyrcanus</i> Woron.
<i>Vitis sylvestris</i> Gmelin		<i>Sanicula europaea</i>
<i>Wilmetica tuberosa</i>		<i>Solidago virga- aurea</i> L.
		<i>Tamus communis</i> L.
		<i>Tilia platyphyllos</i> Scop.

Conyza *Atropa belladonna* *Carex sylvatica*
(2010 Esmailzadeh et al., *canadensis*

b)

(*Carex* sp.) (*Rubus* spp)

(Kostel- Hughes et al., 1998;

Onaaindia & Amezaga, 2000; Bossuyt & Hermy,
2001; Diaz- Villa et al., 2003; Amrein et al., 2005;
Roovers et al., 2006; Zobel et al., 2007 and
.Chaideftou et al., 2009)

()

Jalili et al., (2003)

Esmailzadeh et al., (2011 b)

Esmailzadeh et al., (2010 b)

Bossuyt et al., (2002)

Rovers et al., (2006) Diaz-villa et al., (2003)

l) (*Rubus hyrcanus*)

l) (*Hypericum androsaemum*) (

Carex sylvatica (l) *Atropa belladonna* (

(l) *Conyza canadensis* (l)

Portulaca oleracea (l) (*Diospyrus lotus* L.)

(l)

() (*Pteris cretica*)

(l) (*Athyrium flix- femina*)

	(m ²)	(Cm)	(Cm ³)
Ashton <i>et al.</i> , 1998			
Halpern <i>et al.</i> , 1999			
Leckie <i>et al.</i> , 2000			
Bossuyt <i>et al.</i> , 2002			(<i>F. sylvatica</i>) (<i>Q. robur</i>)
Diaz- villa <i>et al.</i> , 2003			(<i>Q. canariensis</i>)
Diaz- villa <i>et al.</i> , 2003			(<i>Q. suber</i>)
Jalili <i>et al.</i> , 2003			
Arriaga and Mercado, 2004			
Amrein <i>et al.</i> , 2005			(<i>F. sylvatica</i>) Basel
Godefroid <i>et al.</i> , 2006			
Roovers <i>et al.</i> , 2006			(<i>Q. robur</i>) (<i>F. sylvatica</i>)
Zobel <i>et al.</i> , 2007			
Allen and Nowak, 2008			(<i>P. monophylla</i>) (<i>J. osteosperma</i>)
Esmailzadeh <i>et al.</i> , 2011			(<i>F. orientalis</i>) ()

(Esmailzadeh et

.al., 2011 b)
Esmailzadeh et al., (2010 b)

(Gaps)

(Sabeti,

.1994)

(Bossuyt

.& Hermy, 2003)
Esmailzadeh et al., (2010 b)
Chaideftou et al., (2009)
Allen & Nowak (2008)
et al., (2007)
et al.,
Pozas
Roovers (2006)

Archive of SID

Esmailzadeh et al., (2010 b)

(/)

)

(Augusto et al., 2001; Bossuyt et al., 2002; Diaz-
. Villa et al., 2003 & Amrein et al., 2005)

) ((Esmailzadeh et al., 2011 b) (

References

- Allen, E. A. and Nowak, R. 2008. Effect of tree cover on soil seed bank in pinyon- juniper, *Pinus monophylla* and *Juniperus osteosperma*. *Woodland. Rangeland Ecol Manage* 61: 63–73.
- Amrein, D., Rusterholz, HP. And Baur, B. 2005. Disturbance of suburban *Fagus* forests by recreational activities: Effects on soil characteristics, above-ground vegetation and seed bank. *Applied Vegetation Science*. 8: 175-182.
- Anonymous 1998. Esrak forest plane notebook. Forest Rangeland and Watershed Organization. Sari, 228 pp.
- Arriaga, L. and Mercado, C. 2004. Seed bank dynamics and tree- fall gaps in a northwestern Mexican *Quercus- Pinus* forest. *Journal of Vegetation Science*. 15: 661- 668.
- Ashton, P. M. S., Harris, P. G. and Thadani, R. 1998. Soil seed bank dynamics in relation to topographic position of a mixed- deciduous forest in southern New England, USA. *Forest Ecology and Management*. 111: 15- 22.
- Augusto, L., Dupouey, J.L., Picard, J.F. and Ranger, J. 2001. Potential contribution of the seed bank in coniferous plantations to the restoration of native deciduous forest vegetation. *Acta Oecologica*. 22: 87- 98.
- Bakker, H. G. 1989. Ecology of soil seed banks. Academic Press, Inc. San Diego. 462 pp.
- Bakker, J.P., Poschlod, P., Strykstra, R.J., Bekker, R.M. and Thompson, K. 1996. Seed banks and seed dispersal: important topic in restoration ecology. *Acta Botanica Neerlandica*. 45: 461- 490.
- Barbour, M. G., Bruk, J. H. and Pitts, W. D. 1987. *Terrestrial Plant Ecology*. Second edition. The Benjamin/Commings Publishing Company, Inc., Menlo Park. 649 pp.
- Barnes, B.V. 1998, *Forest ecology*, John Wiley and Sons. INC., 773 pp.
- Baskin, C.C. and Baskin, J.M. 1998. *Seeds: ecology, biogeography, and evolution of dormancy and germination*. Academic Press, San Diego. 666 pp.
- Bossuyt, B. and Hermy, M. 2001. Influence of Land Use History on Seed Banks in European Temperate Forest Ecosystems: A Review. *Ecography*. 24(2): 225-238.
- Bossuyt, B. and Hermy, M. 2003. The potential of soil seedbanks in the ecological restoration of grassland and heath land communities. *Belgian Journal of Botany*. 136:23–34.
- Bossuyt, B. and Honnay, O. 2008. Can the seed bank be used for ecological restoration? An overview of seed bank characteristic in European communities. *Journal of Vegetation Science*. 19, 875- 884.
- Bossuyt, B., Heyn, M. and Hermy M. 2002. Seed bank and vegetation composition of forest stands of varying age in central Belgium: consequences for regeneration of ancient forest vegetation. *Plant Ecology*. 162: 33–48.
- Chaideftou, E., Thanos, C.A., Bergmier, E., Kallimanis, A. and Dimopoulos, P. 2009. Seed bank composition and above- ground vegetation in response to grazing in sub- Mediterranean Oak forests (NW Greece). *Plant Ecology*. 201: 255- 265.
- Diaz-Villa, M.D., Maranon, T., Arroyo, J. and Garrido, B. 2003. Soil seed bank and floristic diversity in a forest-grassland mosaic in southern Spain. *Journal of Vegetation Science*. 14: 701- 709.

- Esmailzadeh, O., Hosseini, S. M., Mesdaghi, M., Tabari, M. and Mohammadi, J. 2010. Can soil seed bank floristic data describe the above ground vegetation plant communities? *Environmental Sciences*. 7 (2): 41-62.
- Esmailzadeh, O., Hoseeini, S. M., Tabari, M., Baskin, C. C. and Asadi, H. 2011 a. Persistent soil seed banks and floristic diversity in *Fagus orientalis* forest communities in the Hyrcanian Vegetation Region of Iran. *Flora*. In press.
- Esmailzadeh, O., Hosseini, S. M., Mesdaghi, M., Tabari, M. and Mohammadi, J. 2011 b. Persistent soil seed bank study of Darkola oriental Beech (*Fagus orientalis* Lipsky) forest. *Forest and Wood Products*. 7 (2): 41-62.
- Fourie, S. 2008. Composition of the soil seed bank in alien-invaded grassy fynbos: Potential for recovery after clearing. *Sout African Journal of Botany*. 74: 445- 453.
- Godefroid, S., Phartyal, S. S. and Koedam, N. 2006. Depth distribution and composition of seed banks under different tree layers in a managed temperate forest ecosystem. *Acta oecologica*. 5: 1437-1443.
- Greig-Smith, P. 1983. *Quantitative plant ecology*, University of California Press, Berkely. 359 pp.
- Halpern, C.B., Evans, A. Sh. and Nielson, S. 1999. Soil seed bank in young, closed- canopy forests of the Olympic Peninsula, Washington: potential contributions to understory reinitiation. *Canadian Journal of Botany*. 77: 922- 935.
- Jalili, A. and Jamzad, Z. 1999. *Red data book*. Research Institute of Forests and Rangelands. Tehran, 748 pp.
- Jalili, A., Hamzeh'ee, B., Asria, Y., Shirvanya, A., Yazdania, Sh., Khoshnevisa, M., Zarrinkamara, F., Ghahramania, M.A., Safavia, R., Shaw, S., Hodgson, J. G., Thompson, K., Akbarzadeh, M. and Pakparva, M. 2003. Soil seed banks in the Arasbaran protected area of Iran and their significance for conservation management. *Biological Conservation*. 109: 425-431.
- Kostel- Hughes, F., Young, T.P. and McDonnell, M. 1998. The soil seed bank and its relationship to the aboveground vegetation in deciduous forests in New York City. *Urban Ecosystems*. 2: 43- 59.
- Leckie, S., Vellend, M., Bell, G., Waterway, M.J. and Lechowicz, M.J. 2000. The seed bank in old-growth, temperate deciduous forest. *Canadian Journal of Botany*. 78, 181-192.
- Levin, D. 1990. The seed bank as a source of genetic novelty in plant. *American Naturalist*. 135: 563-572.
- Onaindia, M. and Amezaga, I. 2000. Seasonal variation in the seed banks of native woodland and coniferous plantations in Northern Spain. *Forest Ecology and Management*. 126: 163- 172.
- Roovers, P., Bossuyt, B., Igødt, B. and Hermy, M. 2006. May seed banks contribute to vegetation restoration on paths in temperate deciduous forest?. *Plant Ecology*. 187: 25- 38.
- Sabeti, H. 1994. *Forests, trees and shrubs of Iran*. Yazd University publication. Yazd, 876 pp.
- Stark, K. E., Arsenault, A. and Bradfield, G. E. 2008. Variation in soil seed bank species composition of a dry coniferous forest: spatial scale and sampling considerations. *Plant Ecology*. 197:173-181.
- Thompson, K. 2000. The functional ecology of seed banks. In: Fenner, M. (ed.) *Seeds: the ecology of regeneration in plant communities*, pp. 231-258. CAB International, Wallingford.
- Thompson, K., Bakker, J. and Bekker, R. 1997. *The soil seed banks of North West Europe: methodology, density and longevity*. Cambridge University Press. Cambridge, 288 pp.
- Thompson, K. and Grime, J. P. 1979. Seasonal variation in the seed banks of herbaceous species in ten contrasting habitats. *Journal of Ecology*. 67, 893-921.
- Yan, Q.L., Zhu, J.J., Zhang, J., Yu, L. and Hu, Zh. 2009. Spatial distribution pattern of soil seed bank in canopy gaps of various sizes in temperate secondary forests, Northeast China. *Plant Soil*. 329: 469-480.
- Zobel, M., Kalamees, R., Pussa, K., Roosaluuste, E. and Moora, M. 2007. Soil seed bank and vegetation in mixed coniferous forest stand with different disturbance regimes. *Forest Ecology and Management*. 250: 71- 76.

Persistent soil seed bank in *Khybus* protected area

H. Asadi¹, S. M. Hosseini^{*2} and O. Esmailzadeh³

¹ Ph.D Student of Forestry, Faculty of Natural Resources, University of Tarbiat Modares, I.R. Iran

² Associate Prof., Faculty of Natural Resources, University of Tarbiat Modares, I.R. Iran

³ Assistant Prof., Faculty of Natural Resources, University of Tarbiat Modares, I.R. Iran

(Received: 30 October 2010, Accepted: 14 March 2011)

Abstract

The objective of this study was to determine the floristic composition of the persistent soil seed bank and comparison of its seed bank composition with its above-ground vegetations in *Khybus* protected forest that is one of the best and most virgin Habitats of Box trees in *Hyrceanian* forest. The study of ground vegetation was carried out with releve method and the soil seed bank was estimated using the seedling emergence method. For this purpose 64 releves with 400 m² area were made by systematic-selective method and considering indicator stand concept. For sampling of soil seed bank within each releve, two soil samples were randomly collected by hammering a 20 cm × 20 cm square (400 cm²) hollow metal frame into the soil to a depth of 10 cm. 52 species with mean of 3880 seeds per m² emerged from soil seed banks. *Pteris cretica* L., *Hypericum androsaemum*, *Rubus hyrcanus* Woron., *Carex sylvatica* L., *Rubus hyrcanus* Woron, *Athyrium flix-femina* (L.) Roth, *Atropa belladonna* L. and *Diospyrus lotus* L. with the mean density of 3132, 213, 154, 125.4, 98.5, 34.4 and 17 seeds per m² were exhibited as the most abundant taxa respectively. Forbs, with the possession of the 64 percent seed bank flora, were the most numerous growth forms in soil seed bank. The comparison of seed bank and above vegetation revealed that the similarity between soil seed bank and vegetation is low. Finally, we are concluded that persistence soil seed bank of the study area like other areas, is not capable of restoring the ground vegetation of plant communities.

Keyword: Persistent soil seed bank, above ground vegetation, species similarity, Box tree (*Buxus hyrcanus* Pojark), *Khybus* protected area, *Hyrceanian* forests