

\*

( / / : // : )

± /  
/

(P<0.01)

Archive of SID  
)  
( ) NRC ( )  
( )  
( )  
( )

( ) .( )

/ /

.( )

.( )

( ) .( ) / )

( ) .( ) .( ) (

:

( ) .( ) ( )

( ) / ( )

.( ) .( )

.( )

( )

( ) .( )

( ) /)

.( ) ( )

( ) .( )

Archive of SID

---

Sampson, 1952  
 Stoddart & Smith, 1955  
 Society for Range Management

:  
 .  
 . ( )      . ( )      ( )

%

:( )

$$ME_m = \frac{1}{1 + \frac{1}{W}}$$

$$ME_m = \frac{1}{1 + \frac{1}{W}}$$

$$( ) = ME_m : = W$$

Archive of SID

/  
 o o / o / o  
 +  
 % /      % /

)  
(

)  
(

(AUE)  
( ) ( )

$$\frac{(LAW)^{0.75}}{Y^{0.75}} = AUE$$

) (LAW)<sup>0.75</sup> :  
Y ( /  
AUE

Archive of SID

NRC ( )  
( ) ( )  
:

$$ME = / + / W$$

( ) NRC  
( )

)  
(

---

γ - Live Animal Weight  
Yong & Corbett (1972)

( )	( )	( )				
$/ \pm /$	$/ \pm /$	$/ \pm /$	*			
		$/ \pm /$	*			
	$/ \pm /$	$/ \pm /$	$/ \pm /$			
		$/ \pm /$	$/ \pm /$			
	$/ \pm /$	$/ \pm /$	$/ \pm /$			
		$/ \pm /$	$/ \pm /$			
$/ \pm /$	$/ \pm /$	$/ \pm /$				
		$/ \pm /$				
	$/ \pm /$	$/ \pm /$	$/ \pm /$			
		$/ \pm /$	$/ \pm /$			
	$/ \pm /$	$/ \pm /$	$/ \pm /$			
		$/ \pm /$	$/ \pm /$			
$/ \pm /$	$/ \pm /$	$/ \pm /$				
		$/ \pm /$				
	$/ \pm /$	$/ \pm /$	$/ \pm /$			
		$/ \pm /$	$/ \pm /$			
	$/ \pm /$	$/ \pm /$	$/ \pm /$			
		$/ \pm /$	$/ \pm /$			
—	$/ \pm /$	$/ \pm /$		**		
		$/ \pm /$				
—	$/ \pm /$	$/ \pm /$		**		
		$/ \pm /$				

.\*  
.  
.\*\*

(P<0.01) ) ( )

(P<0.05)

/ ns	/	/		
/ **	/	/		
/ **	/	/		*
/ ns	/	/		
/ ns	/	/		*
/ **	/	/		*
/ ns	/	/		* *
/ ns	/	/		
/ ns	/	/		*
/ **	/	/		*
/ **	/	/		* *
/ ns	/	/		*
/ ns	/	/		* *
/ **	/	/		* *
/ **	/	/		* * *
	/	/		

Archive of SID

/ / /

( )

( )				
/ B	/ B	/ A	/ B	
/ B	/ B	/ A	/ B	
/ A	/ A	/ A	/ A	
/ A	/ A	/ A	/ A	

( )										
/	/	/	/	/	/	/	/	/	/	( )

( )

( ) NRC

(ME = / + / W) ( )

( ) NRC

/ /

/

( )

( )

( ) /

=

/

/

/

/

/ × / = /

( )

( )NRC

( )	( )	
( )NRC	( )	
/ aA	/ aA	/ a
/ bA	/ bA	/ b

(a, b)

(A, B)

(.)

( )NRC

بحث و نتیجه گیری

$$= / \times / = /$$



---

( ) ( ) ( )

( ) / )

/ ( )

/ ( )  
/ ( ) NRC

NRC ( ) ( )

NRC  
(range)

( ) NRC

( )

( )

( )

( ) ( )

( ) NRC

/

Archive of SID

---

				( )...
				(GIS)
				( )
	)			( )
				(
				( )
	( )			( )
				( )
(In Vivo)				( )
	(In Sito – In Vitro)	–	(In Situ)	(In Vitro)
				( )
				( ) .
				( )
	)			( )
				(
				( )
				( )

Archive of SID

15- AFRC (1993). Energy and protein Requirements of Ruminants. An Advisory Manual Prepared by the AFRC Technical Committee on Responses to Nutrients, CAB International, Wallingford, UK.

16- ARC (1984). The nutrient requirements of Ruminants livestock. Supplement No. 1, Report of the Protein Group of the ARC Working Party, Commonwealth Agricultural Bureau, Farnham Royal, UK.

17- Arzani, H. (1994). Some aspect of estimating short-term and long-term rangeland carrying capacity in the western division of New South Wales. PhD. thesis, University of New South Wales, Australia, p: 380

- 
- 18- Arzani, H., M. Zohdi, E. Fish, G.H. Zahedi Amiri, A. Nikkhah and D. Wester (2004). Phonological Effects on Forage Quality of Five Grass Species. University of Tehran, *J. of Range mgt*, 57(6): 624-629.
- 19- Biondini, M., R.D. Pettit and V. Jones (1986). Nutritive Value of Forage on Study Soils as Affected by Tebuthiurn. *Journal of Range Management*, 39(5): 396-399.
- 20- Garza, A.JR., T.E., Fulbright (1988). Comparative chemical composition of armed saltbush and fourwing saltbush. *Journal of Range Management*, 401-403.
- 21- George, Ruyle and Phil, Ogden (1993). What is an A.U.M.? Rangeland Management Specialists, School of Renewable Natural Resources, College of Agriculture, University of Arizona, p:3.
- 22- Geenti, K. G. and Ratry, P. V. (1987). Livestock Feeding on Pasture New Zealand Society of Animal Production. Occasional Publication No. 10, p:39.
- 23- Havstad, K. M. and John C. Malechek (1982). Energy Expenditure by Heifers Grazing Crested Wheatgrass of Diminishing Availability. *J. of Range mgt*. 35(4): 447-450 p.
- 24- Havstad, K. T., D. L. Beker, J. E. Elis, D. M. S. L. Kronberg, D. E. Doornbos and E. L. Ayers (1986). Forage Intake of Different Beef Cattle Biological Types under Range Conditions. *Mon. Ag. Res.* 3:2: 17-19.
- 25- Low, S.G. and C.L. Andrews (1987). A Service for Estimating the Nutritive Value of Forages. Department of Agriculturer, Nutrition and Feed Evaluation Unit, Glenfield, NSW 2167. 423-25.
- 26- Maff (1984). Energy Allowances and Feeding Systems for Ruminants. ADAS Reference Book 43, HMSO, London.
- 27- McDowel, L.R. (1985). Nutrition of Grazing Ruminants in Warm Climatics. Academic Press, New York.
- 28- Oddy, V.H., G.H. Robards and S.G. Low (1983). Prediction of In-vivo Dry Matter Digestibility from the Fiber and Nitrogen Content of a Feed. N. S. W. Department of Agriculture, Nuutritions and Feeds Evaluation Unit, Veterinary Research Station, Glenfield, N. S. W.
- 29- Osuji, P. O. (1974). The Physiology of Eating and the Energy Expenditure of the Ruminant at Pasture. *J. of Range Management*. 27(6): 437-443 p.
- 30- Sampson, W. (1952). *Range Management Principles and Practices*. Wiley, New York. 570 p.
- 31- Society for Range Management (1974). *A Glossary of terms used in range management*. 3th edition. compiled by P.W Jacoby, Denver, Colorado, USA.
- 32- Steven, H. Umberger (1996). *Feeding Sheep: Extension Animal Scientist*, Virginia Tech, Publication Number 410-853, p: 8.
- 33- Stodart Laurence A. and Arthure D. Smith (1955). *Range Management*. 2nd ed. McGraw-Hill, New York, 433 p.
- 34- Vallentine J.F, 2001, *Grazing management*, 2th edi, Academic Press, New York, 657 p.
- 35- Van Soest, P. J. (1982). *Nutritional Ecology of the Ruminant Books*. Ins. Corvallis, or 3-36.
- 36- Yong, B.A. and J.L. Corbett (1972). Maintenance Energy Requirement of Geazing Sheep in Relatin to Herbage Availability. *Icalori Metric Estimates*, Australian Journal of Agricultural, res 23:57.

## Determination of animal unit equivalent (AUE) and daily requirement energy for Sanjabi sheep breed

H. Arzani<sup>\*1</sup>, A. Esfandiari<sup>2</sup>, H. Nourozian<sup>3</sup>, M. Ghorbani<sup>4</sup> and J.Torkan<sup>4</sup>

<sup>1</sup> Professor, Faculty of Natural Resources, University of Tehran, Karaj, I.R.Iran

<sup>2</sup> Expert, Office of Pastoral Affairs Organization, Tehran, I.R.Iran

<sup>3</sup> Professor, Faculty of Agriculture, Islamic Azad University of Karaj, I.R.Iran

<sup>4</sup> Ph.D. Student, Faculty of Natural Resources, University of Tehran, Karaj, I.R.Iran

(Received: 16 October 2005, Accepted: 28 January 2009)

### Abstract

Knowledge on animal requirement, available forage and its quality is fundamental for successful livestock and range management. Since sheep is dominant grazing animal of rangelands in Iran and there are over 27 sheep breeds and each breed has special body size, physiological condition and consequently different daily nutrition requirement, animal unit should be determined for dominant breed of sheep in every region to be calculated the forage requirement based on body weight, daily energy requirement and quality of available forage. In this study, two herds of Kermanshah's Sanjabi breed have been weighted including 50 animals in each herd (15 heads three years old ewe, 15 heads four years old ewe, 5 heads three years old ram, 5 heads four years old ram, 5 heads three months old lamb and 5 heads six months old lamb) in three stages. The body weight of Sanjabi breed (based on average body weight of three and four years old ewe) was  $60.68 \pm 0.56$  kg. Animal unit equivalent (AUE) for ewe, ram and lamb were 1.29, 1.88 and 0.63 kg, respectively. The results of analysis of variance indicated that there are significant differences between weight of ewes and rams ( $P < 0.01$ ). The metabolizable energy requirement determined based on maintenance condition (with 40% extra energy because of regions, topographic condition and watering point distances) using two method of MAFF (1984) formula and NRC (1985) tables. In the first method, the daily energy requirement of animal unit was obtained 10.97 MJ while second method it was 12.28 MJ. Generally, information about daily requirement is essential for animal nutrition management in rangeland and it should be considered in range management designs.

**Keywords:** Animal unit, Sanjabi breed, Daily requirement, Forage quality

---

\*Corresponding author: Tel: +98 261 2249313 , Fax: +98 261 2249313 , E-mail: harzani@ut.ac.ir