

« »

*

(// : // :)

« »

Salsola lanchnantha *Hammada salicornia*

Pteropyrum noeatum *Convolvulus oxyphyllus*

(ADF)

(CP)

(ME)

(DMD)

Hammada

(CP)

()

salicornia

Mj/kg)

(ME)

(

(Torkan *et al.*, 2003)

(Arzani, 2009)

(NDF)

(ADF)

Arzani,)

(Ball *et al.*, 2001 2009

E A

(Arzani, 2009)

(Arzani, 2006)

(Licitra *et al.*, 1997)

(DM)

(NDF)

(ash)

(ADF)

Cook &) .

(Minson, 1987) (Stubbendieck, 1986

(Arzani *et al.*, 2006 Arzani, 1994)

(Arzani, 1994, 2004, 2005)

(Leen & Matin, 2004)

(DMD)

(CP)

(ME)

Crispim *et*) .

(*al.*, 2004

(ADF)

(N)

Brachiaria spp

(Uniyal *et al.*, 2005) .

CP¹ DM

(Heady & Denis, 1994) .

² Neutral detergent fiber(NDF)

³ Acid detergent fiber (ADF)

⁴ Dry matter digestible (DM)

⁵ Dry matter digestible (DMD)

⁶ Metabolizable energy (ME)

¹ Crude protein (CP)

Ritch *et al.*, 2006) .

,« »

NDF ADF

(

(Jamie, 1996) .

Hammada (Moq.) Iljin

Salsola (Botsch.)*Botsch salicornica*

Convolvulus oxyphyllus (Boiss.) *lanchnantha*

Pteropyrum noeanum (Jaub & Spach)

(Arzani & Naseri., 2005)

()

/ /

Jafari *et al.*,)

.(2008

« » (N)

() (Calora, 1998)

(CP)

(CP) % = / × N% (Jafari *et al.*, 2008)

(ADF)

Van Soest,) (AOAC, 1973)

(Soofi Siavash & Janmohamadi, 2004) (1963

) (Oddy *et al.*, 1983)

...

S. lanchnantha (CP) (N) (

H. (CP) (ADF)

salicornia

S. lanchnantha (CP) (

P. noeanum *C. oxyphyllus* $DMD\% = \frac{\text{DM}}{\text{DM}} - \frac{\text{ADF}}{\text{DM}} + \frac{\text{N}}{\text{DM}}$

(ME)

(DMD)

() (SCA)

(ME)

(ADF)

()

(CP)

(ADF)

$ME(Mj/kg) = \frac{\text{DM}}{\text{DM}} - \frac{\text{ADF}}{\text{DM}}$

(ME)

(DMD)

DMD%

ME

(CP)

(ME)

(DMD)

(ADF)

DMD

ADF

(CP)

(DMD)

(ME)

(ADF)

Hammada salicornia

(DMD)

(CP)

(ADF)

(ME)

¹ اندازه واحد دامی در ایران، میش بالغ غیر آبستن و خشک با میانگین وزن ۵۰ کیلوگرم گزارش شده است (Arzani, 2009)

¹ Standing Committee on Agriculture (SCA)

Salsola lanchnantha
(Mj/kg)


Hammada salicornia

F

(ADF)	(ME)	(DMD)	(CP)
**	**	**	**
**	**	**	**
**	**	**	**


×

**



(%))

ADF



(%))



DMD

...

	(CP)	(ADF)	(DMD)	(Mj/Kg DM)
<i>Salsola lanchmantha</i>	/ ± / a	± / a	/ ± / a	/ ± / a
	/ ± / b	± / b	/ ± / b	/ ± / b
	/ ± / c	/ ± / c	/ ± / c	/ ± / c
<i>Hammada salicornia</i>	/ ± / a	/ ± / a	/ ± / a	± / a
	/ ± / b	/ ± / b	/ ± / b	/ ± / b
	/ ± / c	/ ± / c	/ ± / c	/ ± / c
<i>Convolvulus oxyphyllus</i>	/ ± / a	± / a	/ ± / a	/ ± / a
	/ ± / b	± / b	/ ± / b	/ ± / b
	/ ± / c	/ ± / c	/ ± / c	/ ± / c
<i>Pteropyrum noeanum</i>	/ ± / a	/ ± / a	/ ± / a	/ ± / a
	/ ± / b	± / b	/ ± / b	/ ± / b
	/ ± / c	± / c	/ ± / c	/ ± / c

.()

... c b a



Pteropyrum noeanum

(ME)

/ / /

Convolvulus oxyphyllus

...

Salsola

Hammada

lanchnantha

/

(CP)

salicornia

/ /

(CP)

(CP)

(ME)

(ME)

(Arzani *et al.*, 2004)

Convolvulus

(CP)

oxyphyllus

(CP)

Hammada salicornia

(CP)

(ME)

()

Convolvulus oxyphyllus

(Rodny *et al.*, 1991)

(ME)

(Kempton *et al.*, 1979)

(ME)

(Nikkhah & Mohareri, 1996)

(ME)

ADF

ADF

(ME)


(ME)

(Arzani, 2005)

&) (Ghorbani & Asadi Alamuti, 2003)

(Soofi Siavash Janmohamadi, 2004)

(CP) *Pteropyrum* (CP) *noeanum*
 (ME)
 (CP) (ME) (ME)
 (DMD) (CP)
 (DMD) ADF (ME)
 (Arzani *et al.*, 2006) ()
 (Elizalde *et al.*, 1999) (Jafari *et al.*, 2008) .
 (DMD) (ME) (CP)
 (CP) (ME) (ME) (CP)
 ± / (Pepall, 2007) .
 /
 (Pepall, 2007)
 Barrett-) (Lennard *et al.*, 2003)
 ()
 ()



References

- Association of Official Analytical Chemists (AOAC). 1973. Official Methods of Analysis. Washington, D.C.
- Arzani, H. 2009. Forage Quality and Daily Requirement Grazing Animal, 1^{ed} Edition, University of Tehran Press, 354p.
- Arzani, H., Mahdavi, S.Kh., Nikkhah, A. and Azarnivand, H. 2007. Determination of Animal Unit weight and Animal Unit Requirement of Dalagh breed (Case Study: Agh Ghala Region), Iranian Journal of Range and Desert Research, 13, 236-247.
- Arzani, H. 2006. Analysis of measurement methods of the Rangelands. Faculty of Natural Resources, University of Tehran, 50p.
- Arzani, H. and Naseri, K. 2005. Livestock Feeding on Pasture. 1^{ed} Edition, University of Tehran Press (Translated), 300p.
- Arzani, H., Zohdi, M., Fish, E., Zahedi, GH., Nikkhah, A. and Wesrer, D. 2004. Phonological Effects on Forage Quality OF five Grass Species. Journal of Range Management, 57, 624- 629.
- Arzani, H. 1994. Some aspects of estimating short term and long term rangeland carrying capacity in the western division of New South Wals. Ph.D. thesis. University of New South Wals, Australia, 308pp.
- Arzani, H., Basiri, M., Khatibi, F. and Ghorbani, G. 2005. Nutritive value of some zagros mountain rangeland species, Small Ruminant Research, 65, 128-135.
- Ball, D.M., Collins, M., Lacefield, G.D., Martin, N.P., Mertens, D.A., Olson, K.E., Putnam, D.H., Undersander, D.J. and Wolf, M.W. 2001. Understanding Forage Quality. American Farm Bureau Federation Publication 1-01, Park Ridge, IL.
- Barrett-Lennard, E.G., Malcolm, C.V. and Bathgate, A. 2003. Saltland Pastures in Australia- a practical guide. 2nd Edition, Land and Water, Bradon, Australia, 123p.
- Cook, C.W. and Stubbendieck, L. 1986. Range Research Basic Problems and Techniques, Society for Range Management, Colorado, 317p.
- Crispim, S.M.A., Cardoso, E.L. and Frandes, F.A. 2001. Seasonal variation of *Brachiaria spp*, quality in the pantanal of mato Grosso do sul, Brazil. Grassland congress, 379- 380.
- Elizalde, J.C., Merchen, N.R. and Faulkner, D. B. 1999. Fraction of Fiber and Crude protein content In Fresh Forage During Spring Growth. Journal of Animal Science, 77, 476- 484
- Ghorbani, GH. and Assadi Alamuti, A. 2003. Advanced Management of Cow Milky, 1^{ed} Edition, University of Esfahan press, 428p.
- Heady, H.F., Denis, R.D. 1994. Rangeland and Ecology management. West view press. USA. 520p.
- Jafari, M., Javadi, M.R., Hamedanian, F. and Ghorbani, M. 2008. Saltland Pastures (Translated), 1^{ed} Edition, University of Tehran Press, 290p.
- Jamie, L. 1996. Plant materials center plant material Midesouth, 4, 12-19.
- Kempton, T.J., Smith, G.H. and Leng, R.A. 1979. Glucose Metabolism and Growth Cattle Given Molasses Urea Based Diets. Trop Animal Production, 4, 2-9.
- Khedri Gharibvand, H., Dianati, GH.A., Mesdaghi, M. and Shimardi, H. 2008. Comparison of Forage Quality of *Camphorosma monspeliaca* and *Echinoiphora platyloba* in Chaharmahal and Bakhtiari Province. Iranian Journal of Rangeland, 151-161.
- Leen, J. and Martin, N.P. 2004. Forage quality tests and interpretation. Agriculture, Food and Environment sciences, 6, 67-80.
- Licitra, G., Carpino, S., Schadt, I., Avondo, M. and Barresi, S. 1997. Forage quality of native pastures in a Mediterranean area. Animal Feed Science Technology, 69, 315-328.
- Minson, D.J. 1987. Estimation of the nutritive value of forage, in Temperate Pastures, their production, use and management, Eds. J.L. Wheeler, C.J. Pearson and G.E. Roberts, Australian Wool Corporation, 6, 415-422.
- Nikkhah, A. and Mohareri, A. 1996. Nutrient Protein in Ruminant, 1^{ed} Edition, University of Tehran Press, 231p.

-
- Pepall, J. 2007. A saltland grazing trial on their Great Southern property 'Edenia' has fitted in perfectly with the Pepall family's existing farming practices, 1^{ed} Edition, Department of Agriculture & Food, 234p.
 - Ritchie, J.C., Reeves, J.B., Krizek, D.T., Foy, C.D. and Gitz, D.C. 2006. Fiber composition of eastern gamagrass forage grown on a degraded, acid soil. *Field Crops Research*, 97,176–181.
 - Rodny, K.H., Schmidt, J. and Wostuth, M. 1991. *Grazing management and ecological perspective*, 2^{ed} Edition, 261p
 - Safaeyan, N. and Shokri, M. 1995. Using of phenological data in determined of palatability and nutrient value range plants in the Mazandaran Flat. *Iranian Journal of Natural Resources*, 49, 105-113.
 - Soofi Siavash, R. and Janmohamadi, H. 2004. *Animal Feeding (Translated)*, 1^{ed} Edition, Ayej press, 840p.
 - Torkan, J. and Arzani, H. 2003. Study of forage quality changes range species in divers climates. *Iranian Journal of Natural Resources*, 58, 459-471.
 - Uniyal, S.Kr., Awasthi, A. and Rawat, G.S. 2005. Biomass availability and forage quality of *Eurotia ceratoides* Mey IN The rangelands of Changthang, eastern Ladakh. *Current sciences*, 89, 23-36.

Archive of SID

Forage Quality of Range Species in the Steppe Rangelands of Changuleh, Ilam Province

H. Arzani¹, M. Moameri^{*2}, J. Motamedi³ and M. Mohammadpoor⁴

¹Professor, Faculty of Natural Resources, University of Tehran, Karaj, I.R. Iran

²Ph.D. Student, Faculty of Natural Resources, University of Tehran, Karaj, I.R. Iran

³ Assistant professor, Faculty of Natural Resources, University of Orumieh, Orumieh, I.R. Iran

⁴Scientific Member, Agriculture and Natural Resources Research Center of Ilam, Ilam, I.R. Iran

(Received: 2008/October/13, Accepted: 2010/June/2)

Abstract

One of the important factors to make a balance between number of grazing livestock and grazing capacity is whether forage contains sufficient energy and protein during grazing period or not. Therefore in this study, forage quality was evaluated for important species in steppe rangelands of Changuleh located in Ilam province. To do this, samples from four species of *Hammada salicornia*, *Salsola lanchmantha*, *Convolvulus oxyphyllus* and *Pteroporum noeanum*, those made considerable components of rangelands vegetation in the region, were collected during three phenological stages (active growth, flowering and maturing) over 2008. Then chemical compositions were determined including crude protein (CP), acid detergent fiber (ADF), dry matter digestibility (DMD) and metabolizable energy (ME). According to the results, maximum CP was obtained from *Hammada salicornia* in active growth stage. Crude protein component of this species was more than critical level (7%) for sheep in the maintenance condition. Also in this species, ME was in higher critical level (8 Mj/kg) compared with the other species.

Keywords: Forage quality, Crude protein, Dry matter digestibility, Metabolizable energy, Critical level, Phenological stage

*Corresponding author: Tel: +98 915 1895892 Fax: +98 26 32249313 E-mail: m_moameri16@yahoo.com