

(        )

**(*Oryza sativa L.*)**

\*

( / / : / / : )

%

$$\frac{MS(GCA)}{MS(SCA)}$$

**IR62871-175-1-10×** ×

.(Babaeian et al., 1999)

(Hari Ramakrishnan et al.,  
2006) .(Momeni, 1995)

(1953) Jinks & Hayman  
(1956a, b) Griffing              (1954) Hayman

(2003) Vanaja et al. (1985) Wright (1984) Pooni et al.

×

(2006) Malini et al.

(2006) Pradhan et al.

IR62871-175-1-10

$$\frac{MS(GCA)}{MS(SCA)}$$

(2008) Shukla & Pandey  
%

/

GCA

×

SCA

( )

GCA

(

)

( )

SCA

(            ×    )

(2004) Srivastava & Verma

%

$$\frac{MS(GCA)}{MS(SCA)}$$

(Malini et al., 2006; Narayana & Rangasamy, 1991; Shukla & Pandey, 2008; Vanaja et al., 2003)

B (1956a) Griffing / / (GCA)  
 ( / ) ( / ) (Griffing, 1956a) (SCA)  
 ( / ) ( .( ) )  
 GCA t SCA GCA  
 Fs

GCA Ft  
 SAS  
 Diallel Hayman

SCA  
 SCA  
 SCA  
 (Vejdani & Sepahvand, 1993) (Grami, 2000)  
 GCA (Logx)  
 %  
 GCA ( .( ) )  
 SCA

x  
 x IR62871-175-1-10 x %  
 .( ) x IR62871-175-1-10

$$\frac{MS(GCA)}{MS(SCA)}$$

GCA IR62871-175-1-10  
 .( )  
 IR62871-175-1-10 x  
 SCA  
 .( )  
 GCA

GCA

IR62871-175-1-10


MS							
(kg/ha)	(g)	(g)					
/	/	/	/	/	/	/	/
	/	/	/	/	/	/	/
/	/	/	/	/	/	/	/
/	/	/	/	/	/	/	GCA
/	/	/	/	/	/	/	SCA
/	/	/	/	/	/	/	CV
/	/	/	/	/	/	/	
/	/	/	/	/	/	/	$\frac{MS(GCA)}{MS(SCA)}$
							:

(kg/ha)	(g)	(g)					
/	/	/	/	/	/	/	/
/	/	/	/	/	/	/	/
/	/	/	/	/	/	/	/
/	/	/	/	/	/	/	IR 62871
/	/	/	/	/	/	/	
/	/	/	/	/	/	/	S.E(g)
							:

(kg/ha)	(g)	(g)						x
/	/	/	/	/	/	/	/	x
/	/	/	/	/	/	/	/	x
/	/	/	/	/	/	/	/	x
/	/	/	/	/	/	/	/	x
/	/	/	/	/	/	/	/	x
/	/	/	/	/	/	/	/	x
/	/	/	/	/	/	/	/	x
/	/	/	/	/	/	/	/	x
/	/	/	/	/	/	/	/	x
/	/	/	/	/	/	/	/	S.E(S <sub>ig</sub> )
								:

= IR62871-175-1-10 = = = =

× SCA .( )  
 .( ) GCA × × × IR62871-175-1-10



(Hari Ramakrishnan et al., 2006;  
Heidary et al., 2006; Izge et al., 2006)

( )

.(Joshi, 1979)

SCA

x

×IR62871-175-1-10

.(Jenson, 1970)

SCA GCA

SCA

x

×IR62871-175-1-10

x

IR62871-175-1-10

SCA

(kg/ha)							
	/	/	/	/	/	/	/
(g)	/	**	/	/	/	/	/
(g)	/		/	/	/	/	/
	/	**	/	/	/	**	/
	/		/	**	/	**	/
=	=	=	=	=	=	=	=
							**

## REFERENCES

- Babaeian, J. N., Nematzade, Gh. A., Karbalaei, M. T. & Taeb, M. (1999). Evaluation of rice agronomic traits variation in local cultivars of Mazandaran. *Quarterly Sci. & Res. of Shahed Uni*, 26, 15-26. (In Farsi).
- Gerami, A. (2000). Essential options in analysis of statistical data. *Iranian Agron J*, 2, 30-37. (In Farsi).
- Griffing, B. (1956a). A generalized treatment of the use of diallel crosses in quantitative inheritance. *Heredity*, 10, 31-50.
- Griffing, B. (1956b). Concept of general and specific combining ability in relation to diallel crossing systems. *Aust J Biol Sci*, 9, 463-493.
- Haji Amiri, M. (2007). *Estimation of genetic variation and correlations between characters in rice cultivars (Oryza sativa) using path analysis method*. M. Sc. Thesis, SANRU. p 110. (In Farsi).
- Hari Ramakrishnan, S., Anandakumar, C. R., Saravanan, S. & Malini, N. (2006). Association analysis of some yield traits in rice (*Oryza sativa* L.). *J Appl Sci Res*, 2, 402-404.
- Hayman, B. I. (1954). The analysis of variance of diallel tables. *Biometrics*, 10, 235-244.
- Heidary, B., Rezaei, A. M. & Mir Mohamadi Meibodi, M. (2006). Diallel analysis for estimation of genetic parameters of yield and yield components in bread wheat. *J Sci Tech Agric Natu*, 10, 121-139. (In Farsi).
- House, L. R. (1985). *A guide to sorghum breeding*. Second Edition ICRISAT Patanchru, A. P. India.

10. Izge, A. U., Kadams, A. M. & Gungula, D. T. (2006). Studies on character association and path analysis of certain quantitative characters among parental lines of pearl millet (*Pennisetum glaucum*) and their F<sub>1</sub> hybrids in a diallel cross. *Afric J Agric Res*, 1, 194-198.
11. Jenson, N. F. (1970). A diallel selective mating system for cereal breeding. *Crop Sci*, 10, 629-635.
12. Jinks, J. L. & Hayman, B. I. (1953). The analysis of diallel crosses. *Maize Genet Coop New*, 27, 48- 54.
13. Joshi, A. B. (1979). Breeding methodology for autogamous crops. *Indian. J Genet Plant Breed* 39, 567-578.
14. Malini, N., Sundaram, T., Hari Ramakrishnan, S. & Saravanan, S. (2006). Genetic Interpretation of yield related traits in rice (*Oryza sativa* L.). *Res J Agric and Biol Sci*, 2, 153-155.
15. Momeni, A. (1995). *Consideration of combining ability, gene action and correlations study for important agronomic traits in rice*. M. Sc. thesis, Agric. College of Tehran Uni. p 105. (In Farsi).
16. Narayana, K. K. & Rangasamy, S. R. (1991). *Genetic analysis for salt tolerance in rice*. Rice Genetics II. IRRI. Manila. Philippines. P. 167-173.
17. Pooni, S., Jinks, J. L. & Singh, R. K. (1984). Methods of analysis and the estimation of the genetic parameters in brown planthopper resistant varieties. *Indian Agriculturist*, 31, 257-265.
18. Pradhan, S. K., Bose, L. K. & Meher, J. (2006). Studies on gene action and combining ability ananlysis in basmati rice. *J Cent Europ Agric*, 7, 267-272.
19. Shukla, S. K. & Pandey, M. P. (2008). Combining ability and heterosis over environments for yield and yield componens in two-line hybrids involving thermosensitive genetic male sterile line in rice (*Oryza sativa* L.). *Plant Breed*, 127, 28-32.
20. Srivastava, H. K. & Verma, O. P. (2004). Genetic component and combining ability analyses in relation to heterosis for yield and associated traits using three diverse rice growing ecosystems. *Field Crops Res*, 88, 91- 102.
21. Vanaja, T. C., Luckins, C., Babu., V. Radhakrishnan, V. & Pushkaran, K. (2003). Combining ability analysis for yield and yield components in rice varieties of diverse origin. *J Trop Agric*, 41, 7-15.
22. Vejdani, P. & Sepahvand, N. A. (1993). Consideration of general and specific combining abilities of wheat cultivars using diallel cross technique. *Plant and Seed Agric Sci J*, 9(3), 18-22. (In Farsi).
23. Wright, A. J. (1985). Diallel design, analyses and reference population. *Heredity*, 54, 307-311.