

Distribution patterns of *Convolvulaceae* in Iran: priorities for conservation

Received: 09.08.2017 / Accepted: 28.11.2017

Sadaf Sayadi: MSc Student, Department of Plant Sciences and Biotechnology, Faculty of Life Sciences and Biotechnology, Shahid Beheshti University, GC, Tehran, Iran

Ahmadreza Mehrabian✉: Assistant Prof., Department of Plant Sciences and Biotechnology, Faculty of Life Sciences and Biotechnology, Shahid Beheshti University, GC, Tehran, Iran (a_mehrabian@sbu.ac.ir)

Abstract

Assessing priorities is the most important action in conservation programs. Rudimentary, mapping of diversity, and distribution patterns have been used to set priorities on diverse scales. The distribution patterns of plant taxa, largely reflect an actual ecological niche. *Convolvulaceae* s.str exhibits cosmopolitan dispersal of diverse life forms and includes 50–60 genera comprising 1600–1700 species including valuable ornamental, medicinal and food crops as well as weedy taxa. Little attention has been paid to the patterns and diversity centers of this family in scale of Iran and Asia. The present study, explains in as much detail as possible distribution patterns and the priorities for conservation of *Convolvulaceae* in the context of species and habitats in Iran. Threatened species of Iranian *Convolvulaceae* include 20.4% distributed in the Irano-Turanian phytochorion as a priority region for conservation planning. Moreover, this family represents the highest richness in the Alborz and central Zagros mountains of Iran. A meaningful proportion of priority taxa and habitats are located in protected areas of Iran. Regardless, some face several threatening factors (e.g. land use change, deforestation, and overgrazing) outside protected areas; therefore, emergent conservation actions (*in situ* and *ex situ* methods) appear necessary to protect these taxa.

Keyword: Conservation management, distribution map, phytogeography, rare species, species richness

الگوی پراکنش تیره پیچک در ایران: الویت‌های حفاظتی*

دریافت: ۱۳۹۶/۰۵/۱۸ / پذیرش: ۱۳۹۶/۰۹/۰۷

صدف صیادی: دانشجوی کارشناسی ارشد دانشکده علوم و فناوری زیستی، دانشگاه شهید بهشتی، تهران، ایران
احمدرضا محرابیان✉: استادیار دانشکده علوم و فناوری زیستی، دانشگاه شهید بهشتی، صندوق پستی ۱۹۸۳۹۶۹۴۱۱، تهران، ایران
(a_mehrabian@sbu.ac.ir)

خلاصه

ارزیابی الویت‌ها مهم‌ترین اقدام در برنامه‌های حفاظتی محسوب می‌شود. تهیه نقشه تنوع و الگوهای انتشار به صورت متداولی در پایه‌ریزی الویت‌ها در مقیاس‌های متنوع به کار گرفته شده است. تیره پیچک در مفهوم محدود شده نمایانگر یک الگوی انتشار جهان‌وطن متشکل از اشکال متنوع زیستی و مشتمل بر ۵۰-۶۰ جنس از ۱۷۰۰-۱۶۰۰ گونه می‌باشند. تا کنون، مراکز تنوع و انتشار این تیره در مقیاس ایران و آسیا کمتر مورد توجه قرار گرفته است. این مطالعه بیانگر دقیق الگوهای انتشار و الویت‌های حفاظتی در قالب زیستگاهی و گونه‌ای در تیره مذکور در کشور می‌باشد. گونه‌های در معرض تهدید این تیره در ایران ۲۰/۴ درصد می‌باشد که غالباً در مناطق ایران-تورانی اکوسیستم البرز و زاگرس انتشار یافته‌اند. این در حالی است که بیش‌ترین تنوع تیره مورد نظر نیز در این مناطق استقرار یافته است. به علاوه، تعداد زیادی از آرایه‌ها در محدوده مناطق تحت حفاظت قرار می‌گیرند. علی‌رغم این، در خارج از مناطق حفاظت شده با چالش‌های متعددی چون تغییرات کاربری اراضی، پاک‌تراشی جنگل و چرای بی‌رویه در زیستگاه‌هایشان مواجه هستند. بنابراین، اقدامات حفاظتی سریع (مشتمل بر حفاظت داخل و خارج از زیستگاه) جهت حفاظت از آن‌ها بسیار ضروری به نظر می‌رسد.

واژه‌های کلیدی: جغرافیای گیاهی، غنای گونه‌ای، گونه‌های نادر، مدیریت حفاظتی، نقشه انتشار

* بخشی از پایان‌نامه کارشناسی نگارنده اول به راهنمایی دکتر احمدرضا محرابیان ارائه شده به دانشکده علوم و فناوری‌های زیستی دانشگاه شهید بهشتی، تهران

Introduction

Assessing priorities is the most important action in conservation programs (Lovett *et al.* 2000, Bottrill *et al.* 2008). Rudimentary, mapping of diversity, and distribution patterns have been used to set priorities on diverse scales (Wagnetiz 1986, Barthlott *et al.* 1996, Mutke & Barthlott 2005, Olmstead 2013, Mehrabian 2012). The distribution patterns of plant taxa largely reflect an actual ecological niche (MacArthur 1972, Toledo *et al.* 2012), and can be used as a bioindicator of ecosystems (Dieckmann 2003). Accordingly, distribution patterns are considered to be the most important targets of biogeography and ecology (Vetaas & Ferrer-Castán 2008).

The family *Convolvulaceae* s.str. exhibits cosmopolitan dispersal of diverse life forms (Zohary 1973), and includes 50–60 genera comprising 1600–1700 species (Mabberley 1987) that are mainly distributed in tropical and subtropical regions of the world. This huge diversity is the result of a complex interaction between different climatic zones and biogeographical regions in a particular orographic context (Zohary 1973).

Mandaville (1990) has provided valuable data on *Convolvulaceae* in scale of Saudi Arabia besides Sa'ad (1967), and Wood *et al.* (2015) who published well-known monographs on *Convolvulus*. Moreover, phytogeography of some taxa of *Convolvulaceae* (Austin 1998) and distribution of southern *Convolvulaceae* (Austin 2006) are another principal references on this family in the region.

Little attention has been paid to the patterns and diversity centers of scale in Iran and Asia. The present study explains in as much detail as possible distribution patterns and the priorities for conservation in the context of species and habitats. Besides, it aims to determine important plant areas for conservation and phytodiversity of *Convolvulaceae* in Iran.

Materials and Methods

- Study area

The geographical location of the study area is within the borders of Iran. Iran has a total surface area of 1.6 million km² located at 24°–40° N longitude and 44°–64° E latitude and include a limited zone in the orogenic belt (Zagros, Alborz and other mountain chains) that spans the Arabian-African unit and the Asian block (Berberian & King 1981).

Iranian habitats are surrounded by several orographic structures. The Alborz mountain system is composed of an active fold and a segment of the Alpine-Himalayan belt (Stocklin 1974). It appears with a gently sinuous east-west orientation along the northern coastlines of the Caspian Sea (Stöcklin 1974) with an average elevation of over 2000 m containing the highest peak in Iran at Damavand (5670 m). The Zagros is the longest mountain range system of the country and forms a natural massif between the Iranian plateau and the Mesopotamian and Persian Gulf basins at an average elevation of 1305 m (Homke 2007). The Kopet-Dagh in the northeastern (Afshar 1979), Jebal Barez in the central and Makran in southeastern part of Iran are other prominent geomorphological structures (Fischer 1968). Iranian habitats range from 27 m below sea level in the Caspian basin to 5671 m above sea level at Mt. Damavand (Ghorbani 2013), and is categorized into eight zones: Alborz, Zagros, Kopet-Dagh, Sanandaj-Sirjan, southeastern Iran, central Iran, Azerbaijan and eastern Iran (Fig. 1).

Iran receives an annual rainfall of less than one-third of the global average (Shakur *et al.* 2010) and is considered to be an arid region. Because of its topographical diversity, it experiences diverse climatic conditions. Iranian habitats are situated in zonobiom III in the southern zones and zonobiom VII (rIII) in the northern zones (Breckle 2002). Using the new method of world classification by Rivas-Martínez *et al.* (1997); Iranian habitats comprise the macro-bioclimates of the Mediterranean (western, northwestern Iran), temperate (northern Iran) and tropical (southern coast zones of the Persian Gulf and Gulf of Oman) regions.

The distribution zones of the taxa were compiled from 3123 herbarium specimen records from the HSB, W and WU (herbarium abbreviations according to Thiers 2008), and from scientific studies on Iranian flora. The climate data was based on data from the Iranian Climatological Organization [50-year average (1955–2005)] for precipitation and temperature. The bioclimatic units was based on Rivas-Martínez *et al.* (*l.c.*), and the conservation status was assessed on the basis of the criteria of the IUCN Red List at regional scale (IUCN 2011) calculated by Kew Geocad (<http://geocat.kew.org/>; Buchman *et al.* 2011).

The distribution points were marked using ArcView (version 3.2) (ESRI 2000) on geo-referenced geological and geomorphological maps ($1/4^6$ to $1/10^6$) of Iran. These points were mapped per $1^\circ \times 1^\circ$ universal transverse Mercator grid cells (100 km^2 with the exception of boundary area) to analysis the pattern of dispersal and the priorities for conservation management. The Flora Iranica (Rechinger 1963), Flora of Iran (Norouzi 2001) as well as Mozaffarian (2010) were the main references for this assessment. Regardless, several literatures on Iranian flora were assessed as the supplementary data.

The criteria for calculation were the index of species rarity (RI) based on Williams *et al.* (1991), the index of species distribution (SDI) by Selvi (1997), and Solymos & Feher (2005). The output data was scored from zero to one (0–1) for each index with the higher numbers showing higher vulnerability. The RI was computed as the inverse of the cell numbers, including the target area, as $RI = 1/C_i$ in which C_i is the number of grid cells and l is the number of taxa present. The SDI is calculated as $SDI = l - C_i/C$, where C is the total number of grid cells. To calculate the conservation value (CV), the RI and SDI of each species were summed up (the higher scores represent lower CVs). The lowest RI as well as the lowest CV, therefore, showing highest vulnerability and priority to conservation. The biological forms were assessed based on Raunkaier (1934).

Results

- Diversity and distribution

Convolvulaceae is widely distributed in Iran and includes 43 species belonging to the following four genera: *Convolvulus* L. (39), *Calystegia* R.Br. (2), *Ipomoea* L. (1), and *Cressa* L. (1) (Norouzi 2001). Table 1 shows that, 22.2% (10) of *Convolvulaceae* (only *Convolvulus* L.) are confined to the borders of Iran (endemic) (Fig. 3). The greatest abundance occurs on the southern slopes of the central Alborz and Zagros mountains. The greatest abundance belongs to *Convolvulus* L. (up to 20 grid cells) in the central Alborz, Kopet Dag, and the northern and southern Zagros, *Calystegia* R.Br. (up to 3 grid cells) in the central Alborz, *Ipomoea* L. (up to 3 grid cells) in the central Alborz and central Zagros (Figs 2 & 3), *Cressa* L. (up to 8 grid cells) in the northern and southern Zagros, central Alborz and Taftan. The family (up to 65 grid cells) showed the greatest abundance in the central Alborz, Kopet Dag, and the northern and southern Zagros (Fig. 1).

The southern slopes of central Alborz, shows the highest richness of species and genera on the scale of Iran (Fig. 2). The peak species richness is at 35° – 36° N latitude and the highest genera richness was observed at 27° – 32° N latitude (Fig. 3). Altitudinal patterns (Kapos *et al.* 2000) of the studied species can be classified as alpine at above 2500 m, sub-alpine at 1200–2500 m and mountain slopes-to-lowlands at below 1200 m. Species richness severely decreases at above 2000 m. The family distribution is denoted by elevation from 0–4000 m: *Convolvulus* L. at 0–4000 m (Fig. 4), *Cressa* L. to 1500 m, and *Calystegia* R.Br. at 0–2000 m. *Convolvulus persicus*, and *C. commutatus* show the longest and shortest range of distribution, respectively (Fig. 6). *Calystegia* R.Br. is restricted to the northern slopes of the Alborz, but other genera show wider ranges of distribution. *Convolvulaceae* is distributed over 135 of 201 grid cells as follows: *Convolvulus* L. (105), *Calystegia* R.Br. (7), *Ipomoea* L. (3), and *Cressa* L. (47). *Convolvulus* L. and *Calystegia* R.Br. show the highest

and lowest abundance in the taxa studied. Sedimentary rock (32.6%), quaternary deposits (29%), igneous rock (19.5%), metamorphic rock (9.5%), ophiolitic rock (5%), and sedimentary-volcanic rock (4.4%) comprise the habitats of this family in the geological context of Iran. Of these, 48.9% occur in zonobiome III, 37.8% in zonobiome VII (rII), and 13.4% in both.

- Phytogeography and conservation

Phytochorions belong mainly to the Sudano-Zambezian (28 taxa; 62.2%), Irano-Turanian (18 taxa; 40%), and Hyrcanian (Euro-Siberian; 17 taxa; 37.7%) areas. Hemicryptophytes (17 taxa; 37.7%), chamaephytes (11 taxa; 24.4%) and therophytes (18 taxa; 40%) comprise the life form spectra of *Convolvulaceae* in Iran.

All endemic taxa of this family (10 species) belong to *Convolvulus* L. situated at 25° to 38° N latitude from 0–3600 m. They are mainly centered in the southern Zagros and include the Irano-Turanian as well as Sudano-Zambezian regions (Fig. 4). *Convolvulus ammocharis* Boiss. & Hausskn., and *C. koieanus* Bornm. ex Koeie. are found in sedimentary rock., *C. cephalopodus* Boiss., *C. stapfii* Rech.f., and *C. gonocladus* Boiss. are found in Quaternary deposits and sedimentary rock. *Convolvulus eremophilus* Boiss. & Buhse, and *C. schirazianus* Boiss.

are found in Quaternary deposits, sedimentary and igneous rock. *Convolvulus oxysepalus* Boiss., and *C. urosepalus* Pau. are found in Quaternary deposits, sedimentary, igneous and metamorphic rock. At the end of the endemic taxa is *C. turrillianus* Parsa found in Quaternary deposits, sedimentary and ophiolitic rock.

Because of the high sensitivity to demographic and environmental events, rare species are more threatened with extinction than common species in the same environmental conditions (Johnson 1998, Matthies *et al.* 2004). Rarity is calculated by RI that ranged from 1 (*C. cephaloporus* Boiss., *C. koieanus* Bornm. ex Koeie., *C. fatmensis* Kunze, and *C. ammocharis*) to 0.031 (*C. pilosellifolius* Desr.). Besides, rare taxa including very rare (4) only present in one grid cell as well as rare (17) which is present up to 10 grid cells including about 48.8% of Iranian *Convolvulaceae* as followings: *Convolvulus* L. (17), *Ipomoea* L. (1), and *Calystegia* R.Br. (2). The CV ranged from 1.60 (*C. cephaloporus*) to 2.871 (*C. arvensis*) in Iran (Fig. 5, Table 2). The family shows highest species richness in the central Alborz and southern Zagros mountains in Iran. Besides, on the basis of Extent of Occurrence (EOO); Area of Occupancy (AOO) 8(18.6) and 5(11.6), it is classified in threatened categories (CR, EN, VU), respectively (Table 2).

Table 1. Comparison of *Convolvulaceae* among some countries of the region

| Taxonomic rank | Iran | Iraq | Afghanistan | Pakistan | USSR | Turkey | Saudi Arabia |
|----------------|------|------|-------------|----------|------|--------|--------------|
| Genus | 4 | 2 | 4 | 5 | 4 | 4 | 2 |
| Species | 43 | 13 | 13 | 20 | 39 | 39 | 8 |




Fig. 1. Abundance pattern of the genera of *Convolvulaceae* in scale of Iran: a. *Calystegia*, b. *Convolvulus*, c. *Cressa*, d. *Ipomoea*, e. family *Convolvulaceae*.




Fig. 2. a. Richness of genera, b. Richness of species inside occupied grid cells.




Fig. 3. Richness of endemic taxa in *Convolvulaceae*.




Fig. 4. Richness of *Convolvulus* in altitudinal profile.

Table 2. Conservation status and rarity of studied taxa

| Taxon | RI | SDI | CV | EOO | CS | AOO | CS |
|---|-------|-------|-------|---------------|----|-------------|----|
| <i>Calystegia sepium</i> (L.) R.Br. | 0.166 | 0.970 | 1.966 | 83,153.388 | LC | 15,000.000 | LC |
| <i>C. silvatica</i> (Kit.) Griseb. | 0.5 | 0.990 | 1.87 | 2,108.413 | EN | 646.566 | VU |
| <i>Convolvulus acanthocladus</i> Boiss. | 0.038 | 0.870 | 1.958 | 506,135.063 | LC | 77,500.000 | LC |
| <i>C. ammocharis</i> Boiss. & Hausskn. | 1 | 0.995 | 2.265 | 0.000 | CR | 0.000 | CR |
| <i>C. arvensis</i> L. | 0.015 | 0.686 | 2.471 | 1,671,558.095 | LC | 217,500.000 | LC |
| <i>C. betonicifolius</i> Mill. | 0.333 | 0.985 | 1.928 | 19,356.778 | VU | 3,529.961 | NT |
| <i>C. buschiricus</i> Bornm. | 0.25 | 0.980 | 1.89 | 85,316.515 | LC | 15,000.000 | LC |
| <i>C. calverti</i> Boiss. | 0.2 | 0.975 | 1.725 | 23,449.273 | NT | 20,000.000 | LC |
| <i>C. cantabricus</i> L. | 0.052 | 0.905 | 2.227 | 784,353.565 | LC | 67,500.000 | LC |
| <i>C. cephalophorus</i> Boiss. | 1 | 0.995 | 2.435 | 0.000 | CR | 0.000 | CR |
| <i>C. cephalopodus</i> Boiss. | 0.1 | 0.950 | 1.6 | 317,266.876 | LC | 42,500.000 | LC |
| <i>C. chondrilloides</i> Boiss. | 0.055 | 0.910 | 2.125 | 225,054.195 | LC | 75,000.000 | LC |
| <i>C. commutatus</i> Boiss. | 0.031 | 0.840 | 2.311 | 809,148.126 | LC | 120,000.000 | LC |
| <i>C. dorycnium</i> L. | 0.076 | 0.935 | 1.951 | 324,156.082 | LC | 47,500.000 | LC |
| <i>C. elymaiticus</i> Mozaff. | 1 | 0.995 | 2.255 | 0.000 | CR | 0.000 | CR |
| <i>C. eremophilus</i> Boiss. & Buhse | 0.066 | 0.925 | 1.821 | 633,853.144 | LC | 45,000.000 | LC |
| <i>C. erinaceus</i> Ledeb. | 0.066 | 0.925 | 1.871 | 228,647.779 | LC | 55,000.000 | LC |
| <i>C. fatmensis</i> Kunze | 1 | 0.995 | 2.265 | 0.000 | CR | 0.000 | CR |
| <i>C. fruticosus</i> Pallas | 0.045 | 0.890 | 2.035 | 457,342.894 | LC | 75,000.000 | LC |
| <i>C. glomeratus</i> Hochst ex Choisy | 0.25 | 0.980 | 1.67 | 36,964.983 | NT | 15,000.000 | LC |
| <i>C. gonocladus</i> Boiss. | 0.062 | 0.920 | 1.752 | 213,042.980 | LC | 60,000.000 | LC |
| <i>C. koeieanus</i> Bornm. ex Koeie. | 1 | 0.995 | 2.265 | 0.000 | CR | 0.000 | CR |
| <i>C. kotschyanus</i> Boiss. | 0.2 | 0.975 | 1.835 | 104,592.306 | LC | 20,000.000 | LC |
| <i>C. leiocalycinus</i> Boiss. | 0.033 | 0.850 | 1.983 | 643,944.747 | LC | 122,500.000 | LC |
| <i>C. leptocladus</i> Boiss. | 0.071 | 0.930 | 1.831 | 545,561.149 | LC | 50,000.000 | LC |
| <i>C. lineatus</i> L. | 0.045 | 0.890 | 2.485 | 777,192.446 | LC | 72,500.000 | LC |
| <i>C. oxyphyllus</i> Boiss. | 0.066 | 0.925 | 1.761 | 425,971.192 | LC | 70,000.000 | LC |
| <i>C. oxysepalus</i> Boiss. | 0.071 | 0.930 | 1.991 | 204,431.560 | LC | 40,000.000 | LC |
| <i>C. pentapetaloides</i> L. | 0.5 | 0.990 | 1.93 | 14,087.951 | VU | 6,413.481 | LC |
| <i>C. persicus</i> L. | 0.2 | 0.975 | 2.055 | 333,987.713 | LC | 22,500.000 | LC |
| <i>C. pilosellifolius</i> Desr. | 0.031 | 0.840 | 1.921 | 1,371,612.443 | LC | 107,500.000 | LC |
| <i>C. prostrates</i> Forssk. | 0.166 | 0.970 | 1.686 | 211,644.850 | LC | 47,500.000 | LC |
| <i>C. pseudocantabrica</i> Schrenk | 0.090 | 0.945 | 1.975 | 187,080.193 | LC | 32,500.000 | LC |
| <i>C. reticulatus</i> Choisy | 0.090 | 0.945 | 1.975 | 211,644.850 | LC | 47,500.000 | LC |
| <i>C. schirazianus</i> Boiss. | 0.2 | 0.975 | 2.115 | 56,140.251 | LC | 15,000.000 | LC |
| <i>C. siculus</i> L. | 0.333 | 0.985 | 1.758 | 39,449.943 | NT | 4,654.443 | LC |
| <i>C. spinosus</i> Burm. f. | 0.071 | 0.930 | 1.831 | 449,006.399 | LC | 40,000.000 | LC |
| <i>C. stachydifolius</i> Choisy | 0.052 | 0.905 | 2.117 | 906,342.241 | LC | 75,000.000 | LC |
| <i>C. stapfii</i> Rech.f. | 0.2 | 0.975 | 1.835 | 19,158.059 | VU | 5,948.096 | LC |
| <i>C. urosepalus</i> Pau | 0.1 | 0.950 | 2.04 | 565,612.419 | LC | 30,000.000 | LC |
| <i>C. virgatus</i> Boiss. | 0.062 | 0.920 | 1.922 | 262,232.822 | LC | 55,000.000 | LC |
| <i>C. turrillianus</i> Parsa | 0.142 | 0.965 | 1.827 | 181,690.759 | LC | 22,500.000 | LC |
| <i>Cressa cretica</i> Linn. | 0.019 | 0.741 | 2.31 | 1,441,915.359 | LC | 160,000.000 | LC |
| <i>Ipomoea crassicaulis</i> (Benth.) B.L. Rob. | 0.333 | 0.985 | 1.758 | 5,525.909 | LC | 2,742.676 | LC |

RI (Rarity Index), SDI (Species Distribution Index), CV (Conservation Value), EOO (Extend of Occurrence), CS (Conservation Status), AOO (Area of Occupancy), and CS (Conservation Status).

Discussion

This recent study represents the first phytogeographical conservation assessment of *Convolvulaceae* based on the geographical information system at Iranian scale. The results revealed a higher rate of diversity and endemism than other countries in SW Asia. Several rare taxa in the family (Austin 1992) have been confirmed by the efforts of the current study. The endemic taxa are mainly distributed across a range of geology and this diversity has been reported as an essential factor of plant endemism in Iranian habitats (Hedge & Wendelbo 1978). Moreover, the centers of endemism of this family correspond to Iranian centers of endemism (Hedge & Wendelbo 1978, Davis *et al.* 1994). Diversity gently decreases toward NW Asia and sharply decreases towards the western zones of Asia. Iran, Turkey, and USSR are the most important centers of diversity in western and central Asia (Table 1). The family represents a greater rate of diversification than other flowering plants and colonization in a temperate bioclimate across three southern continents four million years ago (Ushimaru & Kikuzawa 1999).

The life forms showing adaptation to ecological factors, therefore, their analysis can help identify ecological properties of the habitats (Diaz & Cabido 1997). The highest diversity of life forms ranges between 1000–1500 m and abruptly falls over 2000 m. Moreover, it shows a peak between 26° to 31° latitude and with an irregularity decrease at the higher and lower latitudes (Fig. 6).

Central Asia especially Iran, shows the highest diversity of *Convolvulus* in Irano-Turanian territory (Wood *et al.* 2015). Several species (e.g. *C. oxyphyllus*, and *C. pilosellifolius*) have been reported as importance and grazing feedstock under desert conditions. *Convolvulus* species are known as important structures to shaping the vegetation especially shrubby species in Middle East (Zohary 1973).

Convolvulecta iranica subtropica comprises several communities in the in border zones of the Persian Gulf to the central lands of Iran and include

C. acanthocladus Boiss. and *C. leocalycinus* Boiss., *C. acanthoclada* Boiss., *C. turrillianus* Parsa, *C. cephalopodus* Boiss., *C. spinosus* Burm., and *C. virgatus* Boiss. (*l.c.*). They are prominent elements of the Sudano-Zambian region and are mainly distributed along coastlines and zones adjacent to the Persian Gulf (*l.c.*). Moreover, these elements (spiny habit and tomentose leaves) compose a different group of Eurasian clade of *Convolvulus* (Wood *et al.* 2015).

Besides, the endemics of *Convolvulaceae* cover Mediterranean (e.g. xeric-continental, desertic-continental) to tropical desertic climes that affected by arid and tropical bio-climatic conditions, respectively. Our results confirmed the Austin (1998) who believes that, endemism zones in *Convolvulaceae* centered mainly in tropical zones. *Convolvulus betonicifolius* Mil., *C. staechadifolium* Choisy., and *C. arvensis* L. are known as the segetal plant communities and synanthropic flora in the Middle East, including Iran (Zohary 1973). *Convolvulus arvensis* by a widely ecological distribution is the most spared taxa in Iran. This species is known as a biological disaster in pastures of Iran as well as throughout the world (Ditomaso 2000), so management actions seems to be necessary to its ecological control in Iran.

Calystegia R.Br. comprises about 26 species, two of which are distributed in Iran. *Calystegia sepium* (L.) R.Br. distributed throughout Europe, Turkey, Caucasia, the northern slopes of the Alborz mountains (N. Iran), and in confined zones of the SW and central Zagros mountain in Iran. Besides, Iran is the far eastern border of the distribution of the mentioned taxa worldwide. *Calystegia silvatica* (Kit) Griseb, similar to other species of genus (Ushimaru & Kikuzawa 1999) is distributed in the Hyrcanian forests. However, other drier Mediterranean bioclimes are the largest barriers to penetration of these taxa into the Iranian plateau.




Fig. 5. a. Species profile along latitude, b. Genera profile along latitude.




Fig. 6. Comparative life form diversity in: Altitudinal profile (a), Latitudinal profile (b). (Th=Therophyte, Ch=Chamaephyte, Hem=Hemicryptophyte).

Ipomoea crassicaulis (Benth.) Robinson is distributed in coastal areas of the Persian Gulf, Saudi Arabia, and Central America. Zohary (1973) classified them as the Mesogeion relict elements of heterogeneous origin of Mediterranean flora during Arcto-Tertiary migrations and Neogene geological events.

The current results show that, *Convolvulaceae* undergoes minor changes at 2000 m but sharply decreases above 2000 m (Fig. 5), so can be classified as mainly lowland to mountainous elements in Iranian habitats as a prominent zone to conservation plans.

A prominent percentage of Iranian *Convolvulaceae* appear to have very rare to rare distributions, out of which them *Convolvulus* mainly cover the threaded categories on the basis of IUCN criteria distributed in the Irano-Turanian phytochorion as a priority region for conservation planning. Moreover, this family represents the highest richness in the Alborz,

and central Zagros mountains. Moreover, Conservation value as a cumulative index appear a more comprehensive perspective to determine the priorities for conservation, so the *C. cephalopodus* and *C. arvensis* show the highest and lowest degree in context of priority for conservation. The presence of spine habit as well as leaves in some taxa (e.g. *C. spinosus*, and *C. acanthocladus*), it can reflect the ecological feasters of habitats as well as protecting habit against herbivory attacks (Hanley *et al.* 2007).

A meaningful proportion of priority taxa and habitats (Figs 2–3 & 5) are located in protected areas of Iran. Regardless, some of them experiment several threatening factors (e.g. land use change, deforestation, and overgrazing) outside protected areas. Emergent conservation actions (*in situ* and *ex situ* methods) appear necessary to protect these priorities for conservation of taxa. Habitat loss is the most important threatened factor

to rare and endangered species (Foin *et al.* 1998). In addition, availability of suitable habitats are considered as a prominent limited factor to rare plants (Aitken *et al.* 2007), mainly those taxa by narrowly geographic distribution. The ecological modelling methods provide a proper strategy to predict the habitat suitability as well as

alternative habitats to *ex situ* conservation of these endangered taxa. However, *in situ* conservation of near threatened taxa and priority habitats (e.g. endemism and diversity zones, threatened habitats), can be an effective strategy to ensure their survival.

References

- Afshar, A. 1979. Stratigraphy, tectonic and petroleum geology of Kopet-Dagh region, northeast Iran: Doctoral College of Science and Technology. University of London, England, 316 pp.
- Aghanabati, A. 1998. Jurassic stratigraphy of Iran. Vol. 1. Geological Survey of Iran, Tehran.
- Agard, P., Omrani, J., Jolivet, L. & Mouthereau, F. 2005. Convergence history across Zagros, Iran; constraints from coalitional and earlier deformation: *International Journal of Earth Sciences* 94: 401–419.
- Aitken, M., Roberst, D.W. & Schultz, L.M. 2007. Modeling distributions of rare plants in the Great Basin, Western North America: *Western North American Naturalist* 67: 26–38.
- Akhani, H. 1998. Plant Biodiversity of Golestan National Park, Iran. *Stapfia* 53: 1–411.
- Akhani, H. 2006. Flora Iranica: facts and figures and a list of publications by K.H. Rechinger on Iran and adjacent areas. *Rostaniha* 7(Suppl. 2): 19–61.
- Alavi, N.M. 1994. Tectonic of the Zagros, orogenic belt of Iran, new data and interpretation. *Tectonophysics* 299: 211–238.
- Alejandra Jaramillo, M. & Paul, S. 2001. Phylogeny and patterns of floral diversity in the genus *Piper* (*Piperaceae*): *American Journal of Botany* 88(4): 706–716.
- Anderson, S. 1994. Area and endemism. *Quarterly Review of Biology* 69: 451–471.
- Andrew, R., Marshall Philip, J., Platts, R., Gereau, E., Kindeketa, W., Kang'ethe, S. & Marchant, R. 2012. The genus *Acacia* (*Fabaceae*) in East Africa: Distribution, diversity and the protected area network. *Plant Ecology and Evolution* 145(3): 289–301.
- Arponen, A., Heikkinen, R., Thomas, C. & Moilanen, A. 2005. The value of biodiversity in reserve selection: representation, species weighting, and benefit functions. *Conservation Biology* 19(6): 2009–2014.
- Austin, D.F. 1992. Rare *Convolvulaceae* in the Southwestern United States. *Annals of the Missouri Botanical Garden*. 78(1):8–16.
- Austin, D.F. & Bianchini, R.S. 1998. Additions and corrections in American *Ipomoea* (*Convolvulaceae*). *Taxon* 47: 833.
- Austin, D.F. 1998. Parallel and convergent evolution in the *Convolvulaceae*. Pp. 201–234. *In*: Mathews, P. & Sivadasan, M. (eds). *Biodiversity and Taxonomy of Tropical Flowering Plants*, Calicut: Mentor Books.
- Austin, D.F. 2006. Noteworthy distributions and additions in Southwestern *Convolvulaceae*. *Canotia* 2(3): 79–106. Linnaeus, C. 1753. *Species plantarum* (ed. 1). Salvius, Stockholm, 1–1200.
- Barthlott, W., Lauer, W. & Placke, A. 1996. Global distribution of species diversity in vascular plants: Towards a world map of phytodiversity. *Erdkunde* 50: 317–328.
- Barthlott, W., Hostert, A., Kier, G., Kuper, W., Kreft, H., Mutke, J., Rafiqpoor, M.D. & Sommer, J.H. 2007. Geographic patterns of vascular plant diversity at continental to global scales. *Erdkunde* 61: 305–316.
- Berberian, M. & King, G.C.P. 1981. Towards a paleogeography and tectonic evolution of Iran.

- Canadian Journal of Earth Sciences 18(2): 210–265.
- Bibby, C.J., Collar, N.J., Crosby, M.J., Heath, M.F., Imboden, C., Johnson, T.H., Long, A.J., Stattersfield, A.J. & Thirgood, S.L. 1992. Putting Biodiversity on the Map: Priority Areas for a Global Conservation. ICBP, Cambridge.
- Bottrill, M.C., Joseph, L.N., Carwardine, J., Bode, M., Cook, C., Walker, S., Wilson, K.A. & Possingham, H.P. 2008. Is conservation triage just smart decision making?. Trends in Ecology & Evolution 23(12): 649–654. doi:10.1016/j.tree.2008.07.007.
- Boissier, E. 1879. Flora Orientalis sive enumeratio plantarum in Oriente a Graecia et Aegypto and Indiae fines hucusque observatarum. IV: *Coroliflorae* et *Monochlamydeae*. Geneva: H. Gorge.
- Bovell-Benjamin, A.C. 2007. Sweet Potato: A review of its past, present, and future role in human nutrition. Advances in Food and Nutrition Research 52: 1–59.
- Braun, G., Mutke, J., Reder, A. & Barthlott, W. 2002. Biotope patterns, phytodiversity and forestline in the Andes, Based on GIS and Remote Sensing Data. Pp. 75–89. In: Körner, C. & Spehn, E.M. (eds). Mountain Biodiversity: a Global Assessment. Parthenon Publishing, London.
- Breckle, S.W. 2002. Walter's vegetation of the earth. The ecological systems of the geobiosphere. Springer, Heidelberg, p. 527.
- Brown J.M., Brummitt R.K., Spencer, M. & Carine, M.A. 2009. Disentangling the bindweeds: hybridization and taxonomic diversity in *Calystegia* R.Br. Botanical Journal of the Linnean Society 160: 388–401. doi: 10.1111/j.1095-8339.2009.00987.
- Brummitt, R.K. 1972. *Calystegia*. Pp. 78–79. In: Tutin, T.G., Heywood, V.H., Burgess, N.A., Moore, D.M., Valentine, D.H., Walters, S.M. & Webb, D.A. (eds). Flora Europaea Vol. 3: *Diapensiaceae* to *Myoporaceae*. Cambridge University Press, Cambridge.
- Brummitt, N. & Lughadha, E.N. 2003. Biodiversity where is hot and where is not hot. Conservation Biology 17: 1442–1448.
- Buchman, A.S., Leurgans, S.E., Boyle, P.A., Schneide, R.J.A., Arnold, S.E. & Bennett, D.A. 2011. Combinations of Motor Measures More Strongly Predict Adverse Health Outcomes in Old Age: The Rush Memory and Aging Project, a Community-Based Cohort Study, BMC Medicine, 9: 42.
- Cao, J.J., S.C. Lee, S.C., Zhang, X.Y., Chow, J.C., An, Z.S., Ho, K.F., Watson, J.G., Fung, K., Wang, Y.Q. & Shen, Z.X. 2005. Characterization of airborne carbonate over a site near Asian dust source regions during spring 2002 and its climatic and environmental significance. Journal of Geophysical Research 110. D03203, doi:10.1029/2004JD005244.
- Clements, F.E. 1905. Research methods in ecology. University Publishing Company, Lincoln, NE.
- Cronquist, A. 1988. The Evolution and Classification of Flowering Plants (2nd ed.). Bronx, New York, USA: The New York Botanical Garden.
- Davis, S.D., Heywood, V.H. & Hamilton, A.C. 1994. Centres of Plant Diversity. A guide and strategy for their conservation. Vol. 1. Europe, Africa, South West Asia and the Middle East.
- Davis, S., Heywood, V.H. & Hamilton, A.C. 1994–1997. Worldwide Fund for Nature and International Union for Conservation of Nature and Natural Resources, Centers of Plant Diversity (Gland, Switzerland).
- Davis, P.H. 1978. Flora of Turkey 6. Edinburgh: Edinburgh University Press.
- Delisle, F., Lavoie, C., Jean, M. & Lachance, D. 2003. Reconstructing the spread of invasive plants: taking into account biases associated with

- herbarium specimens. *Journal of Biogeography* 30: 1033–1042.
- Dieckmann, M. 2003. Species indicator values as an important tool in applied plant ecology a review. *Basic and Applied Ecology* 4: 493–506.
- Diaz, S. & Cabido, M. 1997. Plant functional types and ecosystem function in relation to global change. *Journal of Vegetation Sciences* 8(4): 463–474.
- DiTomaso, J. 2000. Invasive weeds in rangelands: Species, impacts, and management. *Weed Science*. Published and copyrighted by: Weed Science Society of America 48(2): 255–265.
- Djamali, M., Akhiani, H., Khoshravesh, R., Anderieu-Ponel, P. & Brewer, S. 2011. Application of the global bioclimatic classification to Iran: implications for understanding the modern vegetation and biogeography. *Ecologia Mediterranea* 37(1): 91–114.
- Dobson, A.P., Rodriguez, J.P., Roberts, W.M. & Wilcove, D.S. 1998. Geographic distribution of endangered species in the United States. Reid, W.V. Biodiversity hotspots. *Trends in Ecology & Evolution* 13: 275–280.
- ESRI. 2000. ArcView Gis Ver. 3.2a. Environmental Systems Research Institute Inc., California.
- Faith, D.P. & Walker, P.A. 1996. Environmental diversity: on the best-possible use of surrogate data for assessing the relative biodiversity of sets of areas. *Biodiversity and Conservation* 5: 399–415.
- Fisher, W.B. 1968. Physical Geography. *In: Fisher, W.B. (ed.) The Cambridge History of Iran*, Cambridge University Press 1: 3–110.
- Ferrer-Castán, D. & Vetaas, O.R. 2003. Floristic variation, chorological types and diversity: do they correspond at broad and local scales? *Diversity and Distributions* 9: 221–235.
- Fuentes, N., Sañchez, P., Esquivel, J. & Marticorena, A. 2012. A new comprehensive database of alien plant species in Chile based on herbarium records. *Biological Invasions* 15(4): 847–858.
- Foin, T.C., Riley, S.P.D., Pawley, A.L., Ayres, D.R., Carlsen, T.M., Hodum, P.J. & Switzer, P.V. 1998. Improving recovery planning for threatened and endangered species: comparative analysis of recovery plans can contribute to more effective recovery planning. *BioScience* 48 (3): 177–184.
- Ghorbani, M. 2013. The economic geology of Iran. Springer. 581 pp.
- Groom, M.J., Meffe, G.K. & Carroll, C.R. 2006. Principles of Conservation Biology. Third edition. Sinauer Associated, Sunderland.
- Grigoriev, C. 1953. *Convolvulaceae*. Pp. 1–37. *In: Shishkin, B.K. et al. (eds). Flora USSR, Vol. 19: Tubiflorae*. USSR Scientific Academy, Moscow and Leningrad.
- Hansen, A.J. & DiCasteri, F. 1992. Landscape boundaries: consequences for biotic diversity and ecological flows. *Ecological Studies* 92. Springer-Verlag, New York.
- Hanley, G.P., Heal, N.A., Tiger, J.H. & Ingvarsson, E.T. 2007. Evaluation of a classwide teaching program for developing preschool life skills. *Journal of Applied Behavior Analysis* 40(2): 277–300. doi:10.1901/jaba.2007.57-06.
- Hawkins, B.A., Field, R. & Cornell, H.V. 2003. Energy, water, and broad scale geographic patterns of species richness. *Ecology* 84: 3105–3117.
- Hedge, C. & Wendelbo, P. 1978. Patterns of distributions and endemism in Iran. *Notes from the Royal Botanical Garden* 36: 441–464.
- Hernández, H.M. & Navarro, M. 2007. A new method to estimate areas of occupancy using herbarium data. *Biodiversity and Conservation* 16: 2457–2470.
- Heiser, C.B. 1969. Nightshades, the paradoxical plants. San Francisco: Freeman & Company. 200 pp.
- Hill, M.P. & Hulley, P.E. 2000. Aspects of the phenology and ecology of the South American weed, *Solanum sisymbriifolium*, in the Eastern Cape province of South Africa. *African Plant Protection* 6(2): 53–59.

- Hilty, J.A., Lidicker, W.A. & Merenlender, M. 2006. Corridor Ecology: The Science and Practice of Linking Landscapes for Biodiversity Conservation. Island Press, Washington.
- Hernandez-Stefanoni, J.L. 2006. The role of landscape patterns of habitat types on plant species diversity of a tropical forest in Mexico. *Biodiversity and Conservation* 15: 1441–1457.
- Heywood, V.H. & Dulloo, M.E. 2005. *In situ* conservation of wild plant species: a critical global review of best practices. IPGRI Technical Bulletin 11. IPGRI, Rome, Italy.
- Homke, S. 2007. Timing of Shortening and Uplift of the Pusht-E Kuh arc in the Zagros Fold-and-Thrust belt (IRAN). A Combined Magnetostratigraphy and Apatite Thermochronology Analysis, Universidad de Barcelona Facultad de Geología, Departamento de Geodinámica y Geofísica.
- IUCN. 2011. Guidelines for appropriate uses of IUCN Red List Data. Incorporating the Guidelines for Reporting on Proportion Threatened and the Guidelines on Scientific Collecting of Threatened Species. Version 2. Adopted by the IUCN Red List Committee and IUCN. SSC Steering Committee. 78 pp.
- IUCN Standards and Petitions Subcommittee. 2011. Guidelines for Using the IUCN Red List Categories and Criteria. Version 9.0. Prepared by the Standards and Petitions Subcommittee.
- Jalili, A. & Jamzad, Z. 1999. The Red data book of Iran. Research Institute of Forests and Rangelands, 748, Tehran.
- Jedicke, E. 2001. Biodiversität, Geodiversität, Ökiodiversität. Kriterien zur Analyse der Landschaftsstruktur-ein konzeptioneller Diskussionsbeitrag. *Naturschutz und Landschaftsplanung* 33: 59–68.
- Jetz, W., Rahbek, C. & Colwell, R.K. 2004. The coincidence of rarity and richness and the potential signature of history in centres of endemism. *Ecology Letters* 7: 1180–1191.
- Johnson, K.T.M. 1998. Experimental determination of partition coefficients for rare earth and high-field-strength elements between clinopyroxene, garnet, and basaltic melt at high pressures. *Contributions to Mineralogy and Petrology* 133(1–2): 60–68.
- Kala, C.P. 2005. Indigenous uses, population density, and conservation of threatened medicinal plants in protected areas of the Indian Himalayas: *Conservation Biology* 19: 368–378.
- Kandari, L.S., Rao, K.S., Maikhuri, K., Kharkwal, G., Chauhan, K. & Kala, C.P. 2011. Distribution pattern and conservation of threatened medicinal and aromatic plants of Central Himalaya, India: *Journal of Forestry Research* 22(3): 403.
- Kapos, V., Rhind, J., Edwards, M., Price, M.F. & Ravilious, C. 2000. Developing a map of the world's mountain forests. *In: Price, M.F. & Butt, N. (eds). Forests in sustainable mountain development: A state-of-knowledge report for 2000.* CAB International, Wallingford: 4–9.
- Kreft, H. & Jetz, J. 2007. Global patterns and determinants of vascular plant diversity. *PNAS* 104(14): 5925–5930.
- Kurbanov, D. 1994. Flora of Kopetdagh. Pp. 105–128. *In: Fet, V. & Atamuradov, K.I (eds). Biogeography and Ecology of Turkmenistan.* Kluwer Academic Publishers, Dordrecht.
- Lavoie, C. 2013. Biological collections in an ever changing world: Herbaria as tools for biogeographical and environmental studies. *Perspectives in Plant Ecology, Evolution and Systematics* 15: 68–76.
- Leslie Dowe, J. 2010. *Biogeography, Ecology and Systematic* Csiro. Scientific and Industrial Research Organisation.
- Léonard, J. 1981–1989. à l'étude de la flore et de la végétation des déserts d'Iran: étude des aires de distribution les phytochories, les chorotypes.

- Fascicules 1–9. Meise, Jardin Botanique National de Belgique.
- Linder, H.P. 2001. Plant diversity and endemism in sub-Saharan tropical Africa. *Journal of Biogeography* 28: 169–182.
- Lovett, J.R., Rudd, S., Taplin, J. & Frimodt-Moller, C. 2000. Patterns of plant diversity in Africa south of the Sahara and their implications for conservation management. *Biodiversity and Conservation* 9: 37–46.
- Mabberley, D.J. 1987. *The Plant-book*. Cambridge. Cambridge University Press.
- MacArthur, R.H. 1972. *Geographical Ecology*. New York: Harper & Row.
- Mack, R.N., Simberloff, D., Lonsdale, W.M., Evans, H., Clout, M. & Bazzaz, F.A. 2000. Biotic invasions: causes, epidemiology, global consequences and control. *Ecological Society of America* 10: 689–710.
- Malalavidhane, T.S., Wickramasinghe, S.M. & Jansz, E.R. 2000. Oral hypoglycaemic activity of *Ipomoea aquatica*. *Journal of Ethnopharmacology* 72: 293–298.
- Matthies, A., Rajagopalan, K.V., Mendel, R.R. & Leimkuhler, S. 2004. Evidence for the physiological role of a rhodanese-like protein for the biosynthesis of the molybdenum cofactor in humans. *Proceedings of National Academy of Science* 101: 5946–5951.
- Mandaville, J.P. 1990. *Flora of eastern Saudi Arabia*. Kegan Paul International Ltd., London and New York, with National Commission for Wildlife Conservation and Development, Riyadh, 1–482.
- McNaughton, S.J. 1994. Conservation goals and the configuration of biodiversity. *In*: Forey, P.L., Humphries, C.J. & Vane-Wright, R.I. (eds). *Systematics and conservation evaluation, the systematic association (special issue)*, Vol. 50. Clarendon Press, Oxford.
- Meusel, H., Jäger, E., Rauschert, S. & Weinert, E. 1978. *Vergleichend Chronologie der zentrleuropalschen Flora-Kareten*, Band II. Gustav Fischer Verlag. Jena.
- Mehrabian, A.R., Sheidai, M., Noormohammad, Z., Mozaffarian, V. & Asri, Y. 2012. Interpopulations diversity in *Onosma microcarpa* (*Boraginaceae*): Morphological and molecular (ISSR) approach. *Science Medical* 3(3): 187–198.
- Michael, J. & Hopkins, G. 2007. Modeling the known and unknown plant: biodiversity of the Amazon Basin, *Journal of Biogeography* 34: 1400–1411.
- Miller Mittermeier, R.A., Myers, N., Thomsen, J.B. & da Fonseca, G.A.B. 1998. Biodiversity hotspots and major tropical wilderness areas: approaches to setting conservation priorities. *Conservation Biology* 12: 516–520.
- Mittermeier, R.A., Myers, N., Gil, P.R. & Mittermeier, C.G. 1999. Hotspots: Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions (CEMEX, Conservation International and Agrupacion Sierra Madre, Monterrey, Mexico).
- Mittermeier, R.A., Gil, P.R., Hoffman, M., Pilgrim, J., Brooks, T., Mittermeier, C.G., Lamoreux, J. & da Fonseca, G.A.B. 2005. Hotspots revisited: Earth's Biologically Richest and Most Endangered Ecoregions. Conservation International (CI). Washington DC.
- Mostafavi, H., Kiabi, B., Abdoli, A., Mehrabian, A.R., Mahini, A.R., Kami, H. & Naqinezhad, A.R. 2006. *Biodiversity of Mond Protected Area*. Shahid Beheshti University Publishing, 254 pp.
- Mozaffarian, V. 1378. *Flora Khuzestan: Markaz-e Tahghighat-e Manabe Tabiee va Dam-e Khuzestan*, 670 pp. (In Persian).
- Mozaffarian, V. 1379. *Flora of Yazd: Moasese-ye Entesharat-e Yazd*, 636 pp. (In Persian).
- Mozaffarian, V. 2010. Three new species and two species records from Iran, Ilam province. *Iranian Journal of Botany* 16: 204–212.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G. & da Fonseca, G.A.B. 2000. Biodiversity hotspots for conservation priorities, *Nature* 403.

- Nabavi, M. 1978. An Introduction to Geology of Iran. Geological Survey of Iran, Tehran.
- Negi, H.R. & Gadgil, M. 2002. Cross-taxon surrogacy of biodiversity in the Indian Garhwal Himalaya. *Biological Conservation* 105: 143–155.
- Niels, R., Marco, C., Roos, J.W., Ferry, S.E. & Loon, E. 2009. Botanical richness and endemic patterns of Borneo derived from species distribution models. *Ecography* 32: 180–192.
- Noroozi, J., Akhiani, H. & Breckles, S.W. 2008. Biodiversity & phytogeography of alpine flora of Iran. *Biodiversity and Conservation* 17(2): 493–521.
- Norouzi, M. 2001. *Convolvulaceae* Juss. In: Assadi *et al.* (eds). Flora of Iran. Vol. 40. Research Institute of Forests and Rangelands, Tehran, Iran (In Persian).
- Olmstead, R.G. 2013. Phylogeny and biogeography in *Solanaceae*, *Verbenaceae*, and *Bignoniaceae*: a comparison of continental and intercontinental diversification patterns. *Botanical Journal of Linnean Society* 171: 80–102.
- Prasad, K.N., Divakar, S., Shivamurthy, G.R. & Aradhya, S.M. 2005. Isolation of a free radical-scavenging antioxidant from water spinach (*Ipomoea aquatica* Forsk). *Journal of Food and Agriculture organization* 85: 1461–1468.
- Pedrotti, F. 2013. Plant and vegetation mapping. Springer, Heidelberg.
- Petrus, C. & Cunninghamia, H. 1998. Some New South Wales coastal plant distributions: a comparison of herbarium records with transect survey data. Vol. 5(3): 645–664.
- Prendergast, J.R., Quinn, R.M., Lawton, J.H., Eversham, B.C. & Gibbons, D.W. 1993. Rare species, the coincidence of diversity hotspots and conservation strategies. *Nature* 365: 335–337.
- Priscilla, H.C. & Bruce, W. 2009. Can herbarium records be used to map alien species invasion and native species expansion over the past 100 years? *Journal of Biogeography* 36: 651–661.
- Rabinowitz, D. 1981. Biological Aspects of Rare Plant Conservation. Wiley, New York, pp. 205–217.
- Raunkaier, C. 1905. Types biologiques pour la géographie botanique. *Oversigt over Det Forhandlinger*: 347–438.
- Raunkaier, C. 1908. Livsformernes Statistik som Grundlag for biologisk Plantegeografi. *Botanisk Tidsskrift* 29: 42–43.
- Raunkaier, C. 1934. The life forms of plants and statistical plant geography. Clarendon Press, Oxford, 632 pp.
- Raven, P.H. & Axelrod, D.I. 1974. Angiosperm biogeography and past continental movements. *Annals of Missouri Botany Garden* 61: 39–637.
- Rechinger, K.H. 1963. Flora Iranica, Akademische Druck-U. Verlagsanstalt. Graz. Vol. 1.
- Rechinger, K.H. 1989. Fifty years of botanical research in the Flora Iranica Area (1937–1987). In: Tan, K. (ed.). Plant taxonomy, phytogeography and related subjects. The Davis & Hedge Festschrift, Edinburgh, pp. 301–349.
- Regens, N. 1997. Floristic biodiversity and history of African arid regions. *Biodiversity and Conservation* 6: 495–514.
- Reveal, J.L. 1981. The concept of rarity and population threats in plant communities. *Rare plant conservation*, ed. Morse, L.E. & Henefin, M.S. Bronx: New York Botanical Garden, pp. 41–46.
- Riedl, H. 1968. *Boraginaceae*. In: Rechinger, K.H. (ed.). Flora Iranica 48. Akademische Druck-u. Verlagsanstalt.- Graz: Austria.
- Risser, P.G. 1995. The status of the science of examining ecotones. *Bioscience* 45(5): 318–325.
- Rivas-Martínez, S., Sánchez-Mata, D. & Costa, M. 1997. Syntaxonomical synopsis of the potential natural plant communities of North America I. *Itinera Geobotanica* 10: 5–148.
- Sa'ad, F. 1967. The *Convolvulus* species of the Canary Islands, the Mediterranean region and the near and Middle East. *Medelingen van het Botanisch*

- Museum en Herbarium van de Rijks universities Utrecht, 281: 1–288.
- Sarasan, V., Cripps, R., Ramsay, M.M., Atherton, C., McMichen, M., Prendergast, G. & Rowntree, J.K. 2006. Conservation in vitro of threatened plants-progress in the past decade. *In Vitro Cellular Developmental Biology* 42(3): 206–214.
- Santi, E., Maccherini, S., Rocchini, D., Bonini, I., Brunialti, G., Favilli, L., Perini, C., Pezzo, F., Piazzini, S., Rota, E., Salerni, E., Chiarucci, A. & Sarfatti, G. 2010. Simple to sample: Vascular plants as surrogate group in a nature reserve: *Journal for Nature Conservation* 18: 2–11.
- Sankaran, M. 2009. Diversity patterns in Savanna grassland communities: implications for conservation strategies in a biodiversity hotspot Mahesh Biodiversity Conservation 18: 1099–1115.
- Selvi, F. 1997. Rare plants on mount Aiata, Italy: vulnerability to extinction on an ecological “Island”. *Biology Conservation* 81: 257–266.
- Schery, R.W. 1951. *Plants for Man*, Prentice Hall, Adapted from Vavilov Englewood Cliffs, NJ, 1972.
- Schmidt, M., Kreft, H., Thiombiano, A. & Zizka, G. 2005. Herbarium collections and field data-based plant diversity maps for Burkina Faso. *Diversity and Distributions* 11: 509–516.
- Schonbeck-Temesy, E. 1972. *Solanaceae*. In: Rechinger, K.H. (ed.). *Flora Iranica*, Vol. 100. Akademische Druck-u. Verlagsanstalt, Graz, Austria.
- Schulman, L., Toivonen, T. & Ruokolainen, K. 2007. Analyzing botanical collecting effort in Amazonia and correcting for it in species range estimation. *Journal of Biogeography* 34(8): 1388–1399.
- Shakur, A., Roshan, Gh., Najafe, R. & Kanei, A. 2010. Evaluating climatic potential for palm cultivation in Iran with emphasize on degree-day index, *African Journal of Agricultural Research* 13: 99–118.
- Simpson, M.G. 2007. *Plant systematics*, Elsevier Academic Press. 603 pp.
- Singh, B. 1981. Establishment of First Gene Sanctuary in India for *Citrus* in Garo Hills. Concept Publishing, New Delhi, India.
- Solymos, P. & Feher, Z. 2005. Conservation prioritization based on distribution of land snails in Hungary. *Conservation Biology* 19(4): 1084–1094.
- Spooner, D.M. & Hijmans, R.J. 2001. Potato systematics and germplasm collecting. *American Journal of Potato Research* 78: 237–268.
- Stattersfield, A.J., Crosby, M.J., Long, A.J. & Wege, D.C. 1998. *Endemic bird areas of the world*. Birdlife International, Cambridge.
- Stocklin, J. 1974. Northern Iran: Alborz Mountains. In: Spencer, A.M. (ed.). *Mesozoic-Cenozoic belts data for orogenic studies; Alpine-Himalayan Orogens*: Geological Society of London (special publication) 4: 212–234.
- Svhmidt, M., Kreft, H., Thiombiano, A. & Zizka, G. 2005. Herbarium collections and field data-based plant diversity maps for Burkina Faso. *Diversity and Distribution* 11: 509–516.
- Takin, M. 1972. Iranian geology and continental drift in the Middle East: *Nature* 235: 147–150.
- Thiers, B.M. 2008. Continuously updated: Index Herbariorum.
- Tilman, D. 1999. The ecological consequences of changes in biodiversity: a search for general principles. *Ecology* 80: 1455–1474.
- Toledo, J.A., Kaminski, J., Santanna, M.A. & Santos, D.R., 2012. Tampão Santa Maria (TSM) como alternativa ao tampão SMP para medição da acidez potencial de solos ácidos. *Revista Brasileira de Ciência do Solo* 36: 427–435.
- Ushimaru, A. & Kikuzawa, K. 1999. Variation of breeding dystem, floral rewards, and reproductive success in clonal *Calystegia* species (*Convolvulaceae*). *American Journal of Botany* 86(3): 436–446.

- Ushimaru, A. & Kikuzawa, K. 1999. Variation in breeding system, floral rewards, and reproductive success in clonal *Calystegia* species (*Convolvulaceae*). *American Journal of Botany* 86: 436–446.
- Vega, I.L., Ayala, O.N.N. & Conteras-Medina, R.L. 2004. Patterns of diversity, endemism and conservation: an example with Mexican species of *Ternstroemiaceae* Mirb. ex DC. (*Tricolpates: Ericales*). *Biodiversity and Conservation* 13: 2723–2739.
- Wagnetiz, G. 1986. *Centaurea* in South-West Asia: Patterns of distribution and diversity. *Proceedings of the Royal Society of Edinburgh. Sect. B. Biological Sciences*, 89B: 11–21.
- Wood, S., Nolte, S., Burrige, M., Rudloff, D. & Green, W. 2015. 'Dimensions and location of high-involvement management: fresh evidence from the UK Commission's 2011. Employer Skills Survey'. *Human Resource Management Journal* 25: 2: 166–183.
- Zohary, M. 1973. *Geobotanical foundations of the Middle East*. 2 Vols. Stuttgart. 739 pp.