

GIS

*

Moghaddam@tabrizu.ac.ir

Fijani@tabrizu.ac.ir

Nadiri@tabrizu.ac.ir

// :

// :

Arc View

.(Thapinta, and Hudak, 2003)

.(Vrba & Zaporozec, 1994)

(Worrall, and Besien, 2004; Babiker, et al., 2005)

()

(Antonakos, and Lambrakis,

)

.2007)

(

.(Vrba, and Zoporozec, 1994)

.(Kim, and Hamm, 1999)

(Samey, and

.(Rahman, 2007)

.(Zabet, 2002)

()


Gang, 2008)

)

.(Asghari Moghaddam & Fijani .2008

GIS

()



()

()

() ۶

()

:

()

(Alavi Naini, Bolurchi, 1973)

(Yilmaz, et al., 1998)

()

:

()

(δ)

)

(

()

()

()

(Evans, and Mayers, 1990; Al-Adamat, et al., 2003; Babiker, et al., 2005)

δ) (Asghari moghaddam & Fijani , (.2009)

)
(

() (LeGrand, 1964)
(Panagopoulos, et al., 2006) ²PCSM

(Panagopoulos, et al., 2006)

PCSM

() (Aller, et al., 1987)
()

:()

GIS	GIS		D R A S T I C

با (Merchant, 1994)

GIS

...

GIS

(Soper, 2006)

)

6

GIS

ArcView-GIS

GIS

(

()

Spatial


Geoprocessing ,

Analyst

Model Builder

GIS

GIS



GIS

() :

ArcView-GIS

Model Builder

()

GIS


GIS

ArcView


(

.()


.()



:()



:()




:()

.(Singhal, & Gupta, 1999)

6

.()

.()



:()

(Aller, et

)

6

6


.al., 1987)

...

()

ArcView

()




()

GIS

ArcView

()



()

Model Builder

)


(

د

()

)

() ()



()


ArcView-GIS

۵

GIS

۵


های



:()

۵

۵



:()

- 1- Drastic
- 2- Point Count System Models
- 3-Depth of groundwater
- 4-Recharge
- 5-Aquifer media
- 6-Soil media
- 7-Topography
- 8-Impact of the vadose zone media
- 9-Conductivity
- 10-Rate
- 11-Weight
- 12- Extension
- 13- Clip
- 14- Merge
- 15- Overlay
- 16- Digital Elevation Model (DEM)
- 17- Kriging
- 18- Overlay
- 19- Arithmetic
- 20- Weighted

()

GIS

GIS

.ArcView-GIS

GIS

6

Al-Adamat,R.A.N., I.D.L.,Foster, S.M.J.,Baban . 2003. Groundwater vulnerability and risk mapping for the Basaltic aquifer of the Azraq basin of Jordan using GIS, remote sensing and DRASTIC. *Applied Geography* 23, 303–324.

Alavi Naini,M., M.H.,Bolurchi .1973. Explanatory text Maku Quadrangle Map 1:250000. Geological Survey of Iran.

Aller,L., et al. 1987. DRASTIC: A Standardized System for Evaluating Ground Water Pollution Potential Using Hydrogeologic Settings, EPA 600/2-87-035. Ada, Oklahoma: U.S. Environmental Protection Agency.

Antonakos,A.K., N.J.,Lambrakis .2007. Development and testing of three hybrid methods for the assessment of aquifer vulnerability to nitrates, based on the drastic model, an example from NE Korinthia, Greece. *Journal of Hydrology* 333, 288– 304.

Asghari Moghaddam, A., E.,Fijani .2008. Distribution of fluoride in groundwater of Maku area, northwest of Iran. *Environment Geology*, 56, 281-287.

Asghari Moghaddam, A., E.,Fijani .2009. Hydrogeologic framework of the Maku area basalts, north western Iran. *Hydrogeology Journal*, 17, 949-959.

Babiker,I.S., et al. 2005. A GIS-based DRASTIC model for assessing aquifer vulnerability in Kakamigahara, Heights, Gifu Prefecture, central Japan. *Science of the Total Environment* 345, 127–140.

Evans,B.M., W.L.,Mayers .1990. A GIS-based approach to evaluating regional groundwater pollution potential with DRASTIC. *Journal of Soil and Water Conservation* 45, 242–245.

-
- LeGrand,H.E. 1964. System for evaluating the contamination potential of some waste sites. *Journal of American Water Works Association* 56, 959–974.
- Kim,Y.J., S.Y.,Hamm .1999. Assessment of the potential for ground water contamination using the DRASTIC/EGIS technique, Cheongju area, South Korea: *Hydrogeol. J.*, 7 (2): 227-235.
- Merchant,J. 1994. GIS-based groundwater pollution hazard assessment: a critical review of the DRASTIC model. *Photogrammetric Engineering and Remote Sensing* 60 (9), 1117–1127.
- Panagopoulos,G., A.,Antonakos, N.,Lambrakis .2006. Optimization of DRASTIC model for groundwater vulnerability assessment, by the use of simple statistical methods and GIS. *Hydrogeology Journal* (published online).
- Rahman,A. 2007. A GIS based model for assessing groundwater vulnerability in shallow aquifer in Algarh, India. *Applied Geogr.*, 28 (1): 32-53.
- Samey,A.A., C.,Gang .2008. A GIS based DRASTIC model for the assessment of groundwater vulnerability to pollution in west Mitidja: Blida city, Algeria. *Research Journal of Applied Sciences* 3 (7): 500-507.
- Singhal,B.B.S., R.P.,Gupta .1999. *Applied Hydrogeology of Fractured Rocks*. Kluwer Academic Publisher. 400 p.
- Soper,R.C. 2006. Groundwater vulnerability to agrochemicals: A GIS-based DRASTIC model analysis of Carrol, Chariton, and Saline Counties, Missouri, USA. Master science thesis, University of Missouri-Columbia.
- Thapinta,A., P.,Hudak .2003. Use of geographic information systems for assessing groundwater pollution potential by pesticides in Central Thailand. *Environmental International* 29, 87–93.
- Vrba,J., A.,Zoporozec .1994. *Guidebook on mapping groundwater vulnerability*. International Contributions to Hydrogeology. Verlag Heinz Heise GmbH and Co. KG.
- Worrall,F., T.,Besien .2004. The vulnerability of groundwater to pesticide contamination estimated directly from observations of presence or absence in wells. *Journal of Hydrology* 303 (1-4), 92–107.
- Yilmaz,Y., Y.,Guner, F.Saroglu .1998. Geology of the quaternary volcanic centers of the east Anatolia. *Journal of Volcanology and Geothermal Research* 85, 173-210.
- Zabet,T.A. 2002. Evaluation of aquifer vulnerability to contaminant potential using DRASTIC method. *Environ. Geol.*, 43(1-2): 203-208.