

()

()

//

(W-36)

x

/

(P< /)

/

(P< /)

()

Archive of SID

()

()

()

()

()

()

1. Relative Bioavailability Value (RBV)

()

:

(±)

/

/

/

/

/

()

(AOAC)

(.)

/

()

Archive of SID

W-36)

/

(

x

/	/	/	/	/	/	/	/	/	
/	/	/	/	/	/	/	/	/	
/	/	/	/	/	/	/	/	/	
/	/	/	/	/	/	/	/	/	
/	/	/	/	/	/	/	/	/	
/	/	/	/	/	/	/	/	/	
/	/	/	/	/	/	/	/	/	
/	/	/	/	/	/	/	/	/	-
/	/	/	/	/	/	/	/	/	-
									()
/	/	/	/	/	/	/	/	/	"
/	/	/	/	/	/	/	/	/	"
/	/	/	/	/	/	/	/	/	"
/	/	/	/	/	/	/	/	/	"
									meq/kg Na + K - Cl
/	/	/	/	/	/	/	/	/	"
/	/	/	/	/	/	/	/	/	"

A :

/ B₁₂ K₃ E

/ () (H₂O) :

() () () (H₂O) ()

()

:

()

()

()

SAS

()

(RBV)

()

SAS

(P< /)

(P< /)

() /

()

(P< /)

(P< /)

(P< /)

()

-
- 1. Futura
 - 2. Slope Ratio

/ a	/ ab	/ ab	/ c	/ ab	/ ab	/ b	/ c	/ d	(%)
± /	± /	± /	± /	± /	± /	± /	± /	± /	()
/ b	/ ab	/ ab	/ a	/ b	/ b	/ b	/ ab	/ b	()
± /	± /	± /	± /	± /	± /	± /	± /	± /	()
/ ab	/ ab	/ bc	/ bc	/ a	/ bcd	/ cde	/ e	/ de	()
±	± /	±	± /	±	± /	±	±	± /	()
/ a	/ ab	/ cd	/ ef	/ ab	/ bc	/ de	/ f	/ f	()
± /	± /	± /	± /	± /	± /	±	±	± /	()
/ ab	/ ab	/ ab	/ ab	/ ab	/ b	/ ab	/ a	/ b	()
± /	± /	± /	± /	± /	± /	± /	± /	± /	()
/ b	/ b	/ ab	/ ab	/ b	/ b	/ b	/ a	/ ab	()
± /	± /	± /	± /	± /	± /	± /	± /	± /	()
/ a	/ a	/ ab	/ bc	/ a	/ a	/ bc	/ cd	/ d	()
± /	± /	± /	± /	± /	± /	± /	± /	± /	()
/	/	/	/	/	/	/	/	/	()
± /	± /	± /	± /	± /	± /	± /	± /	± /	()

(P < /)

Archive of SID



/ b	/ a	/ a	(%)
± /	± /	± /	()
/ b	/ ab	/ a	()
± /	± /	± /	()
/ b	/ ab	/ a	()
± /	± /	± /	()
/ b	/ a	/ a	()
± /	± /	± /	()

(P < /)

()

:

/	/	/	/	/	/	/	/
/	/	/	/	/	/	/	/
/	/	/	/	/	/	/	/
/	/	/	/	/	/	/	/
/	/	/	/	/	/	/	/



REFERENCES

1. AOAC, 1990. Official methods of analysis of the Association of Official Analytical Chemists. P. O. Box. 540 Benjamin, Franklin station Washington D.C.
2. Ammerman, C. B., D. H. Baker & A. J. Lewis, 1995. Bioavailability of nutrients for animals amino acids, minerals and vitamins. Academic press, INC.
3. Burnell, T. W., G. L. Cromwell & T. S. Stahly, 1990. Effects of particle size on the biological availability of calcium and phosphorus in difluorinated phosphate for chicks. Poultry Science. 69:1110.
4. Forbes, R. M. & J. W. Erdman, 1983. Bioavailability of trace mineral elements. Annu. Rev. Nutrition, 3:213.
5. Guinotte, F. Nys, Y. de Monredon F., 1991. The effects of particle size and origin of calcium carbonate on performance and ossification characteristics in broiler chicks. Poultry Science,70:1908.
6. Henry, M. H. & G. M. Pesti, 2002. An investigation of calcium citrate malate as a calcium source for young broiler chicks. Poultry Scienc, 81: 1149.
7. Hunter, J.E., R.A. ditcher & H.C. Kandel, 1933. Relative utilization of calcium from calcium carbonate and calcium gluconate by chickens. Proc. Soc. Exp. Biol, 31:70.
8. McNaughton, J. L., B. C. Dilworth & E. J. Day, 1974. Effect of particle size on the utilization of calcium supplements by chick. Poultry Science, 53:1024.
9. Poneros-Schneier, A. G. & J. W. Erdman, 1989. Bioavailability of calcium from sesame seeds, almond powder, whole wheat bread, spinach and non fat dry milk in rats. Journal of Food Science.54:150.
10. Roland, D. A., Sr., 1986. Egg shell quality: oyster shell versus limestone and the importance of particle size or solubility of calcium source. World Poultry Science. 42:166.
11. Roland, D. A., Sr. & H. R. Harms, 1973. Calcium metabolism in the laying hen. 5. Effect of various sources and sizes of calcium carbonate on shell quality. Poultry Science. 52:369.
12. SAS Institute, Inc. 1998. SAS/STAT® user's guide. Version6. 4th ed. SAS Institute Inc, Cary, NC.
13. Saunders-blades, J. L. & D. M. Anderson, 2003. Effect of calcium source and particle size on production performance and bone quality of the laying hen. Fact sheet # 18. Nova Scotia Agricultural College. Truro, N.S.
14. Littell, R. C., P. R. Henry, A. J. Lewis & C. B. Ammerman, 1997. Estimation of relative bioavailability of nutrients using SAS procedures. J. Animal Science, 75:2676-2683.
15. Yoshida, M. & H. Hoshii, 1982. Relationship between ash content of the toe and hardness of the tibia bone of meat type chick. Jpn. Poultry Science. 19:126.
16. Zhang, B. & C.N. Coon, 1997. Improved *in vitro* methods for determining limestone and oyster shell solubility. J. Appl. Poultry Res. 6:94-99.