

()

//

Archive of SID

(/ /)

(FAO)

۳/

(iv)

()

(/)

()

(.)

(/)

)

(

()

()

()

()

(۱۷)

E-mail: khodarahmi_m@yahoo.com

()

()

()

()

()

(Juanillo 92)

K O

K O

()

-
- 1.CIMMYT
 - 2. α -lattice

()

Puma-super

//

()

()

()

()

(/)

()

()

()

Archive of SID

()

()

$$= \frac{\dots}{\dots} + \frac{\dots}{\dots} \text{BY+} / \text{HI} \quad (\text{BY}) \quad (\text{HI})$$

**	ns	ns	ns	ns	**	**	ns	ns	ns	ns
ns	**	ns	ns	ns	ns	ns	ns	ns	**	
ns	**	ns	ns	ns	ns	ns	ns	**		
*	ns	*	ns	ns	ns	**	ns			
**	ns	ns	ns	**	ns	*				
**	**	*	ns	ns	ns					
**	ns	ns	ns	*						
**	ns	ns	**							
ns	ns	ns								
ns	ns									
ns										

% % : ns ** *


/ (R²)

()

()

()

$$\begin{aligned}
 & \text{(HI)} \quad \text{(NS)} \quad \text{((NSP)} \\
 & \quad \quad \quad \text{(LSP)} \quad \text{(TKW)} \\
 & \text{()} \quad \text{NSP} + / \quad \text{NS} + / \quad \text{HI} + / \quad \text{TKW} + / \quad \text{LSP} \\
 & \quad \quad \quad = / + / \\
 & \quad \quad \quad / \quad (\text{R}^2) \\
 & \text{()} \quad \text{()}
 \end{aligned}$$



(/ /)

(/)

()

()

)

(/ /)

()

(/ / /)

()

(1913)

:

/ **

/ **

/ **

/ **

_____ : -

/ **

/

/

/

/ **

/

/

/

/

/ **

/

/

/

/

/ **

/

/

/

/ *

/

()

(/)

)

(

REFERENCES

6. Aly, R. M., & A. Y. A. El-Bana. 1994. Grain yield analysis for nine wheat cultivars grown in newly cultivated sandy soil under different N fertilization levels .Zagazing .J .Agric.Res.21:67-77.
7. Bhatt, G. M. 1973. Significance of path coefficient analysis in determining the nature of character association.Euphytica 22:338-343.
8. Bittle, D. C., & J. P. Gustafson. 1991. High molecular weight glutenin from wheat triticale flour improvement .Proc.2nd.Int.Triticale Symp.Mexico. D.F. CIMMYT. P.550-553.
9. Briggs, K. G. 2001 The growth potential of triticale in western Canada . A report that outlines the characteristics and potential of triticale as a in w. Canada and identifies the barriers to reaching this potential .
10. Dewey,D. R. & K. H. Lu. 1959. A correlation and path-coefficient analysis of component of crested wheat grass seed production Agron.J.51:515-518.

- :
11. Gebeyhoue, G. D., R. Knott, & R. J. Baker. 1982. Relationship among duration of vegetative and grain filling phases ,yield components and grain yield in durum wheat cultivars.Crop Sci.22:287-290
 12. Gill, K. S., M. L. Verma, & G. S. Sandha. 1976. Studies on variability and character association in triticale .Crop Improv. 3:64-69
 13. Karpenstein-Machan, M. & J. Heyne. 1992. Yield and yield structure of winter cereals triticale and wheat in the middle mountain areas of northern Hessen. Agribiol. Res.45:88-96
 14. Reddy. V. R. K. 2001. Character association in hexaploid triticale . Crop Res. 22(1) : 94-98.
 15. Singh, K. N. & R. S. Rana. 1983.Genetic variability and path analysis in triticale grown in alkali soil.Indian J.Agric.Sci.53:1-4.
 16. Singh, H. B. & G. S. Sethi. 1974. Path and regression analysis in triticale . Plant Sci. 6:11-16
 17. Smith,R. L., M. E. Schweder, & R. D. Barnett. 1994. Indentification of glutenin alles in wheat and triticale using PCR generated DNA markers.Crop Sci .34:1373-1378.
 18. Yagbasanlar, T., & H. Ozkan. 1995. Correlation and path coefficient analysis for ear characters in triticale under Mediterranean climatic condition.Crop Sci. 174:297-300.

Archive of SID