





NAG

(.) (Pyelonephritic Scaring)

)  
 ) ( )  
 - -  
 ((CRP) (ESR)  
 ) DMSA  
 ( )  
 ESR  
 DMSA  
 NAG DMSA VCUG  
 NAG (N-Acetyl glucose aminidase)  
 NAG  
 NAG )  
 RA 1000 ( )  
 Jaffee  
 ) NAG  
 ( ) ELISA ( NAG )  
 DIAZYME ( )





نمودار ۱- ROC Curve مساسیت و اختصاصیت NAG ادرار در تشفیص پیلونفریت ماد

SPSS

DMSA

NAG

VCUG

regression analysis t-test

% ±

NAG

%

NAG/Cr NAG

) NAG

CRP ESR CBC)

جدول ۱ - نتایج آزمایش های مورد بررسی در کودکان مبتلا به پیلونفریت ماد در بیمارستان مفید سال ۱۳۸۵

%

( (VCUG DMSA

%

/ ± /	/ ± /	( )
/ ± /	/ ± /	( ) NAG
/ ± /	±	( ) NAG/Cr

DMSA

Candice 1990

NAG

Metro health

NAG/Cr

NAG

NAG ( P < / )

NAG

%

%

NAG

( )

CBC

NAG

NAG

CRP ESR

( )

VCUG DMSA

	Jantusch	NAG
	DMSA	
Cuartero Jantusch	( DMSA )	
Candice	Belli %	NAG %
	( )	
	Cuartero	
	NAG	NAG
	( )	
		Belli
		( )
		NAG
		NAG
		( )

## REFERENCES

1. Ellis D Avner "Pediatric Nephrology" 5<sup>th</sup> Edt. Lippincott Williams & Willkins 2004, New York. p: 850.
2. Behrman, Nelson textbook of pediatrics, 2004, Saunders. p: 1785-89.
3. Johnson CE. Urinary NAG and selection of children for radiologic evaluation after UTI. Pediatrics 1990;86:211-5.
4. Jantusch BA. Urinary NAG and Beta 2 microglobulin in the diagnosis of UTI. Pediatr Infec Dis J 1994;13:294-9.
5. Roilides E. Increased urine interleukin-6 concentrations correlate with pyelonephritic changes on Tc-Dimercaptosuccinic acid scans in neonates with urinary tract infections. J Infec Dis 1999; 180: 904-7.

- 
6. Haraoka M. Elevated interleukin-8 levels in the urine of children with renal scarring and/or vesicouretral reflux. *J Urol* 1996; 155(2): 678-80.
  7. Fretzayas A. Polymorphonuclear elastase as a diagnostic marker of acute pyelonephritis in children. *Pediatrics* 2000; 105: E28.
  8. Cuartero RA. Urinary NAG in patients with UTI. *Eur J Urol* 1998; 33: 348-50.
  9. Belli A . Evaluation of NAG In upper and lower UTI in childhood. *Minerva Pediatr* 1996; 48: 503-7.

Archive of SID