

//
//

()

*

STEC

EPEC EAEC

(Shiga Toxin Escherichia coli)

ETEC

()
STEC

STEC

(.)

f_jaafari580@yahoo.com :

Salmonella-Shigella Agar

(Promadisa, manufacture by Hispanlab, S.A
Madrid)

(.)

%

°C

()

Campy Blood Agar

(Himedia M 9960 Imdia)

°C

Yersinia (% CO2 % O2 % N2)

(Merck KGaA Darmstadt, – Selective Agar

°C

Germany)

(.)

(.)

(Merck

GaA- Germany)

DNA

°C

/ ml

DNA PCR .

st, lt stx2, stx1, eae

()

pCVD432

stx⁺

(Mast company) O157:H7

μl

μl

PCR

/ ,DNA

/ ()

/

/

Tag

(ice-pack)

(Eppendorf AG , 2233, Hamburg, Germany)

DNA

°C

(XLD) Xylose-Lysine-Desoxycholate-agar

UV °C °C
 kb °C
 PCR (*LT*⁺, *ST*⁺) *E. coli* ATCC 35401 :
 O157:H7 (*stx1*⁺, *stx2*⁺, *eae*⁺) *E. Coli* ATCC 43894
stx (Agarose % / PCR
 O157:H7 (Mast Company) LE Roche applied science, Germany)

| | (5'-3') | (bp) |
|----------------------|--|------|
| STEC (<i>stx1</i>) | CAGTTAATGTGGTGGCGAAG CTGCTAATAGTTCTGCGCATC | |
| STEC (<i>stx2</i>) | CTTCGGTATCCTATTCCCGG GGATGCATCTCTGGTCATTG | |
| EPEC (<i>eae</i>) | TGCGGCACAACAGGCGGCGA CGGTCGCCGCACCAGGATTC | |
| EAEC (pCVD432) | CTGGCGAAAGACTGTATCAT CAATGTATAGAAATCCGCTGTT | |
| ETEC (<i>lt</i>) | TCTCTATATGCACACGGAGC CCATACTGATTGCCGCAAT | |
| ETEC (<i>st</i>) | TCTGTATTGTCTTTTTACC TTAATAGCACCCGGTACAAGC | |

(% /)
 (% /)
 %
 . () / :
 EPEC ,EAEC (% /) () % /
) STEC () %
 LT ETEC . ()
 (% /) ST (% /) ()
 (% /) % /
 STEC . () % /
eae⁻, *stx2*⁺, *stx1*⁺
stx2, *stx1*, *eae*

| | | <i>E.coli</i> O157-H7 <i>stx1, stx2</i> | |
|-------------------------------------|------------------|---|--------|
| (n=) | : | | |
| a (%) | | (% /) | |
| (% /) | | | |
| | STEC | (% /) | |
| (% /) | <i>stx1</i> | | |
| (% /) | <i>stx2</i> | | |
| (% /) | <i>stx1/stx2</i> | (n =) | |
| (% /) | EAEC | | |
| (% /) | EPEC | () | |
| (% /) | | () | |
| | ETEC | (/) | |
| (% /) | <i>LT</i> | (/) | |
| (% /) | <i>ST</i> | (/) | |
| (% /) | <i>LT/ST</i> | | () |
| | | (/) | |
| (%) a | | (/) | |
| | | (/) | |
| | | (/) | |
| | | (/) | |
| | | (/) | |
| | | (/) | |
| | | (/) | |
| | (%) | : / | (/) |
| <i>Shigella flexneri</i> /EAEC | (/) | / ± / | (± SD) |
| <i>Salmonella paratyphi A</i> /ETEC | (/) | / ± / | (± SD) |
| <i>Shigella boydii</i> /EAEC | (/) | / ± / | |
| <i>Salmonella typhi</i> /EPEC | (/) | () | |
| <i>Salmonella paratyphi C</i> /EPEC | (/) | (/) | |
| <i>Shigella sonnei</i> /EAEC | (/) | (/) | |
| <i>Salmonella group D</i> /EAEC | (/) | (/) | |
| <i>Shigella flexneri</i> /STEC | (/) | (/) | |
| <i>Campylobacter</i> /STEC | (/) | (/) | |
| <i>Campylobacter</i> /EAEC | (/) | (/) | |
| | (/) | (/) | |

()

(% /)

(%)

:

| (%) | (%) | (%) | (%) | (%) | () | |
|-----|-----|-----|-----|-----|-------|-------------|
| / | / | / | / | / | ± / | |
| / | / | / | / | / | / ± / | STEC |
| / | / | / | / | / | / ± / | EAEC |
| | / | / | / | / | / ± / | EPEC |
| / | / | | / | / | ± / | |
| / | / | / | / | / | / ± / | |
| / | | / | / | / | ± / | ETEC |

(% /)

(% /)

()

Archive of SID

()

()

(%)

()

()

(% /)

EPEC

(% /)

(% /)

()

()

(HUS)

()

Diarrhegenic E.coli

ETEC EAEC

ETEC (%)

(%) EAEC

()

(% /) STEC (*Diarrhegenic E.coli*)

PCR

() (%) / % /)

STEC

()

STEC (HUS)

STEC EPEC

HUS

REFERENCES

1. Bern C, Martinez J, de Zoysa I, Glass RI. The magnitude of the global problem of diarrhoeal disease: a ten-year update. Bull World Health Organ. 1992; 70(6):705-14.
2. Yamashiro T, Nakasone N, Higa N, Iwanaga M, Insisiengmay S, Phounane T et al. Etiological study of diarrhoeal patients in Vientiane, Lao People's Democratic Republic. J Clin Microbiol 1998; 36:2195-9.
3. Wheeler JG, Sethi D, Cowden JM. Study of infectious intestinal disease in England: rates in the community, presenting to general practice, and reported to national surveillance. BMJ 1999; 318:1046-50.
4. Svenungsson B, Lagergren A, Ekwall E, Evengard B, Hedlund KO, Karnell A et al. Enteropathogens in adult patients with diarrhea and healthy control subjects: a 1-year prospective study in a Swedish clinic for infectious diseases. Clin Infect Dis 2000; 30(5):770-8.
5. Kelly MT, Brenner DJ, Farmer J. Enterobacteriaceae. In: Clennette EH, Ballows A, Tausler Jr WJ and Shadomy HJ. Manual of clinical microbiology. 4th ed. Washington, DC: ASM press.1985; P: 263-277.

6. Olsvik O, Strockbine NA. PCR detection of heat-stable, heat-labile, and shiga-like toxin genes in *Escherichia coli*. In: Persing DH, Smith TH, Tenover F, White TJ, editors. Diagnostic molecular microbiology-Principles and applications. Washington DC: ASM. 1993; P: 271-6.
7. Heuvelink AE, Van de Kar NC, Meis JF, Monnens LA, Melchers WJ. Characterization of verocytotoxin-producing *Escherichia coli* O157 isolates from patients with haemolytic uremic syndrome in Western Europe. *Epidemiol Infect* 1995; 115:1-14.
8. Schmidt H, Knop C, Franke S, Aleksic S, Heeseman J, Karch H. Development of PCR for screening of enteroaggregative *Escherichia coli*. *J Clin Microbiol* 1999; 33: 701-5.
9. Salmanzadeh-ahrabi S, Habibi E, Jafari F, Zali MR. Molecular epidemiology of *Escherichia coli* diarrhea in children in Tehran. *Ann Trop Ped Liverpool, UK*. 2005; 25: 35-9.
10. Katouli M, Jaafari A, Farhoudi-Moghaddam AA, Ketabi GR. Etiological studies of diarrhoeal diseases in infants and young children in Iran. *J Trop Med Hyg* 1990; 93: 22-7.
11. MoezArdalan K, Zali MR, Dallal MM, Hemami MR, Salmanzadeh-Ahrabi S. Prevalence and pattern of antimicrobial resistance of Shigella species among patients with acute diarrhoea in Karaj, Tehran, Iran. *J Health Popul Nutr* 2003; 21(2): 96-102.
12. Jamal WY, Rotimi VO, Chugh TD, Pal T. Prevalence and susceptibility of Shigella species to 11 antibiotics in a Kuwait teaching hospital. *J Chemother* 1998; 10(4): 285-90.
13. Bellido JB, Galiano JV, Tirado MD, Gonzalez-Cano JM, Safont L. Incidence of sporadic cases of the most frequent intestinal infections in Castellon. *Rev Esp Salud Publica* 2003; 77(5): 629-38.
14. Katouli M, Ketabi GR, Ketabi GR. The role of diarrhoeagenic *Escherichia coli* in acute diarrhoeal diseases in Bandar-Abbas, Iran. *J Med Microbiol* 1988; 27(1):71-4.
15. Soltan-Dallal M.M, Moezardalan K. Frequency of Yersinia species infection in paediatric acute diarrhea in Tehran. *Eastern Mediterranean Health Journal*. 2004; 10(1,2):152-8.
16. Aslani MM, Badami N, Mahmmoodi M, Bouzari S. Verotoxin-producing *Escherichia coli* (VTEC) infection in randomly selected population of Ilam Province (Iran). *Scand J Infect Dis* 1998; 30 (5) :473-6.
17. Seung-Hak Cho, Jong-Hyun Kim, Jong-Chul Kim, Hyun-Ho Shin, Yeon-Ho Kang, Bok-Kwon Lee. Surveillance of Bacterial Pathogens Associated with Acute Diarrheal Disease in the Republic of Korea During One Year, 2003. *J Microbiol Soc Korea* 2006; 44(3): 327-35.
18. Rezayi M, Salmanzadeh-ahrabi S, Edalatkhah H, Jafari F, MoezArdalan S, Zolfagarian K, et al. Epidemiology of bacterial-induced acute diarrhea in Varamin. *Pejouhandeh Bimonthly Research Journal* 2004; 8(7): 467-474.
19. Aslani MM, Bouzari S. An epidemiological study on Verotoxin-producing *Escherichia coli* (VTEC) infection among population of northern region of Iran (Mazandaran and Golestan provinces). *Eur J Epidemiol* 2003; 18(4): 345-9.
20. Keskimaki M, Eklund M, Pesonen H, Heiskanen T, Siitonen A; Study Group. EPEC, EAEC and STEC in stool specimens: prevalence and molecular epidemiology of isolates. *Diagn Microbiol Infect Dis* 2001; 40(4): 151-6.
21. Black RE, Morris SS, Bryce J. Where and why are 10 million children dying every year. *Lancet* 2003; 361: 2226-34.