

*

Yale -Pas

Bieri facial pain score

% / %
% %
% . (P < /)
% /
% . (P < /)
%

()

.()

.()

()

(recovery)

.()

.()

Lloyd .()

%

management

.()

()

.()

(IV line)

.()

/ mg/kg

(%)

/ mg/kg

(% /)

mg/kg

(P< /)

(sedation)

(%)

(%)

(% /)

(%)

(%)

Bieri Facial Pain

(P< /)

Scale

(.)

Yale-PAS

(.)

Likert

(P< /) %

% /

(.)

(P< /)

% /

(P< /) % /

(%)

(% /)

(%)

(P< /)

	(/)	(/)	/ ± /	(/)	(/)
	(/)	(/)	/ ± /	(/)	(/)
P value	()	()	()	()	()
	«	»	«	»	«
P < /	(/)	(/)	(/)	(/)	(/)
P < /	(/)	(/)	(/)	(/)	(/)
P < /	(/)	(/)	(/)	(/)	(/)
P < /	(/)	(/)	(/)	(/)	(/)

)	())	()
((((
/ ± /	± /	±	± /
± /	± /	± /	± /

Crock

±

.()

Ljungman

Bieri Facial Pain

.()

(%)

(% /)

(%)

Lejus

Archive of SID

)

(

.()

Ljungman

Lam

%

.()

Lloyd

.()

%

/ ± /

Yale

/ ± /

.()

()

Malinovsky

.()

pH

REFERENCES

1. Pizzo PA, Poplack DG. Principles and Practice of Pediatric Oncology. 4th ed. Philadelphia: Lippincott, Williams & Wilkins; 2001; PP: 1340-5.
2. Ljungman G, Kreuger A, Andréasson S, Gordh T, Sörensen S. Midazolam nasal spray reduces procedural anxiety in children. *Pediatrics* 2000; 105: 73-8.
3. Bhatt-Mehta V, Rosen DA. Sedation in children: current concepts. *Pharmacotherapy* 1998; 18: 790-804.
4. Crock C, Olsson C, Phillips R, Chalkiadis G, Sawyer S, Ashley D, et al. General anaesthesia or conscious sedation for painful procedures in childhood cancer: the family's perspective. *Arch Dis Child* 2003; 88: 253-7.
5. Lloyd CJ, Alredy T, Lowry JC. Intranasal midazolam as an alternative to general anaesthesia in the management of children with oral and maxillofacial trauma. *Br J Oral Maxillofac Surg* 2000; 38: 593-5.
6. Everitt IJ, Barnett P. Comparison of two benzodiazepines used for sedation of children undergoing suturing of a laceration in an emergency department. *Pediatr Emerg Care* 2002; 18: 72-4.
7. Vivarelli R, Zanotti F, Battaglia D, Caggese G, Stella G, Gilli G, et al. Premedication with intranasal midazolam in children of various ages. *Minerva Anesthesiol* 1998; 64: 499-540.

-
8. Davis PJ, Tome JA, McGowan FX, Cohen IT, Latta K, Felder H. Preanesthetic medication with intranasal midazolam for brief pediatric surgical procedures: effect on recovery and hospital discharge times. *Anesthesiology* 1995; 82: 2-5.
 9. Kain ZN, Mayes LC, Cicchetti DV, Bagnall AL, Finley JD, Hofstadter MB. The Yale Preoperative Anxiety Scale: how does it compare with a "Gold Standard"? *Anesth Analg* 1997; 85: 783-8.
 10. Lejus C, Renaudin M, Testa S, Malinovsky JM, Vigier T, Souron R. Midazolam for premedication in children: nasal vs. rectal administration. *Eur J Anaesthesiol* 1997; 14: 244-9.
 11. Lam C, Udin RD, Malamed SF, Good DL, Forrest JL. Midazolam premedication in children: a pilot study comparing intramuscular and intranasal administration. *Anesth Prog* 2005; 52: 56-61.
 12. Malinovsky JM, Lejus C, Servin F, Lepage JY, Le Normand Y, Testa S, et al. Plasma concentrations of midazolam after i.v., nasal or rectal administration in children; *Br J Anaesth* 1993; 70: 617-20.

Archive of SID