

//
//

()

PCR

*

(*Human Papilloma virus*) HPV

HPV

(E9 E6)

HPV

PCR

PCR

HPV

HPV

HPV

HPV16

HPV %

HPV

HPV

Premalignant

PCR

HPV:

Rigoni-Stezn

HPV

()

()

Kessler

HPV₁₈
E₇ E₆

HPV SCC HPV₁₈ HPV₁₆ HPV

HPV Hella (.)

E₇ E₆ mRNA HPV HPV

())

Premalignant (HPV

()

% (Early stage) HPV

(.) HPV

HPV (.)

) :

(HIV)

()

(%) ()

()

HPV HPV CIN

HPV () HPV₁₆ HPV

/

HPV

DNP DNA

(DN8115C)

DNA

PCR

HPV

koilocyte

Sacace) HPV Screening HPV

(V-26-50R

HPV

PCR-mix-1

HPV A6 , A7 , A9

PCR

()

DNA

: /

2.5X PCR-mix-1 HPV

Hot Start Polymerase / buffer-red

Reaction Mix

DNA

°C

°C

°C

°C

°C

UV PCR

/

Squamous Cell Carcinoma (SCC)

) HPV Typing HPV High risk

(V-25-50R Sacace

HPV

PCR-mix-1

DNA

HPV (%)
 () ()
 .() ()

HPV
 /
 PCR-mix-1 :
 / 2.5X buffer-red
 Hot Start Polymerase
 Mix Reaction
 DNA
 °C : °C °C
 °C °C
 UV PCR

.()

HPV

() () () ()
 ()
 () () ()
 () ()) /
 .()

HPV SCC

HPV ()
 .()

HPV

HPV
 ()

PCR

.()
 HPV

HPV16

/ /
 /
 / / <

HPV16 Masumato, Silins, Ma CL
 HPV
 ()

HPV
 HPV
 PCR
 DNA probs Hybrid Capture2 (HC2)
 HPV ()
 ()

HPV
 ()

HPV

Premalignant

REFERENCES

2. Murray PR, Rosenthal KS, Kobayashi GS. Medical Microbiology. 4th ed. USA: Mosby Inc. 2001; PP: 459-62.
3. Kufe DW, Pollock RE, Weichselbaum RR, Bast RC, Gansler TS, Holland JF, et al. Cancer Medicine. Canada: BC Decker Inc. 2003.
4. Kenneth J. Ryan. Kistner's Gynecology & Women's Health, 7th ed. 1999; PP: 100-9.
5. Winn WC, Koneman EW, Allen SD, Janda WM, Schreckenberger PC, Procop GW, et al. Koneman's Color Atlas and Textbook of Diagnostic Microbiology. 6th ed. USA: Lippincott Williams & Wilkins. 2006; PP: 1362-4.
6. Evidence reports and summaries, Numbers 1-60/Evaluation of Cervical Cytology. Available at: www.nlm/ahrq

7. Kraus I, Molden T, Holm R, Lie AK, Karlsen F, Kristensen GB, et al. Presence of E6 and E7 mRNA from human papillomavirus types 16, 18, 31, 33, and 45 in the majority of cervical carcinomas. *J Clin Microbiol* 2006; 44(4): 1310-7.
8. Andreoli TE, Carpenter CJ, Griggs RC, Loscalzo J. *Cecil Essentials of Medicine*, 6th ed. USA: WB Saunders Company. 2004; P: 660.
9. Berrington de Gonzalez A, Green J. Comparison of risk factors for invasive squamous cell carcinoma and adenocarcinoma of the cervix: collaborative reanalysis of individual data on 8,097 women with squamous cell carcinoma and 1,374 women with adenocarcinoma from 12 epidemiological studies. *Int J Cancer* 2007; 120(4): 885-91.
10. Eleuterio Jr, Giraldo PC, Goncalves AK, Cavalcante DI, de Almeida Ferreira FV, Mesquita SM, et al. Prognostic markers of high-grade squamous intraepithelial lesions: the role of p16INK4a and high-risk human papillomavirus. *Acta Obstet Gynecol Scand* 2007; 86(1): 94-8.
11. de Silva R, Karunaratne K, Mendis LN, Ramesh R, Chow VT. PCR detection and typing of human papilloma virus DNA in squamous carcinoma of the cervix in a cohort of Sri Lankan women. *Ceylon Med J* 2006; 51(3): 114-7.
12. Tornesello ML, Duraturo ML, Botti G, Greggi S, Piccoli R, De Palo G, et al. Italian HPV Working Group. Prevalence of alpha-papillomavirus genotypes in cervical squamous intraepithelial lesions and invasive cervical carcinoma in the Italian population. *J Med Virol* 2006; 78(12): 1663-72.
13. Hadzisejdic I, Simat M, Bosak A, Krasevic M, Grahovac B. Prevalence of human papillomavirus genotypes in cervical cancer and precursor lesions. *Coll Antropol* 2006; 30(4): 879-83.
14. Peedicayil A, Abraham P, Sathish N, John S, Shah K, Sridharan G, et al. Human papillomavirus genotypes associated with cervical neoplasia in India. *Int J Gynecol Cancer* 2006; 16(4): 1591-5.
15. Rajaram S, Gupta G, Agarwal S, Goel N, Singh KC. High-risk human papillomavirus, tumor suppressor protein p53 and mitomycin-C in invasive squamous cell carcinoma cervix. *Indian J Cancer* 2006; 43(4): 156-62.
16. Gudleviciene Z, Didziapetriene J, Suziedelis K, Lapkauskaitė L. Investigation of human papillomavirus, its types and variants. *Medicina (Kaunas)* 2005; 41(11): 910-5.
17. Wall SR, Scherf CF, Morison L, Hart KW, West B, Ekpo G, et al. Cervical human papillomavirus infection and squamous intraepithelial lesions in rural Gambia, West Africa: viral sequence analysis and epidemiology. *Br J Cancer* 2005; 93(9): 1068-76.
18. Asato T, Maehoma T, Nagai Y, Kanazawa K, VAzato H, Kariya K. A large case-control study of cervical cancer risk associated with human papillomavirus infection in Japan, by nucleotide sequencing-based genotyping. *J Infect Dis* 2004; 189:1829-32.
19. Konidaris S, Kouskouni EE, Panoskaltsis T, Kreatsas G, Patsouris ES, Sarivalassis A, et al. Human papillomavirus infection in malignant and benign gynaecological conditions: a study in Greek women. *Health Care Women Int* 2007; 28(2): 182-91.
20. An HJ, Kim KR, Kim IS, Kim DW, Park MH, Park IA, et al. Prevalence of human papillomavirus DNA in various histological subtypes of cervical adenocarcinoma: a population-based study. *Mod Pathol* 2005; 18(4): 528-34.

-
21. Qiu AD, Wu EQ, Yu XH, Jiang CL, Jin YH, Wu YG, et al. HPV prevalence, E6 sequence variation and physical state of HPV16 isolates from patients with cervical cancer in Sichuan, China. *Gynecol Oncol* 2007; 104(1): 77-85.
 22. You K, Liang X, Qin F, Guo Y, Geng L. High-risk human papillomavirus DNA testing and high-grade cervical intraepithelial lesions. *Aust N Z J Obstet Gynaecol* 2007; 47(2): 141-4.
 23. Ozçelik B, Serin IS, Gökahmetoğlu S, Başbuğ M, Erez R. Human papillomavirus frequency of women at low risk of developing cervical cancer: a preliminary study from a Turkish university hospital. *Eur J Gynaecol Oncol* 2003; 24(2): 157-9.
 24. Ozsaran AA, Dikmen Y, Akercan F, Zekioglu O, Terek MC, Mgoyi L, et al. The triage of squamous cell abnormalities of cervical cytology by human papilloma virus screening. *Eur J Gynaecol Oncol* 2003; 24(6): 535-8.
 25. Silins I, Wang X, Tadesse A, Jansen KU, Schiller JT, Avall-Lundqvist E, et al. A population-based study of cervical carcinoma and HPV infection in Latvia. *Gynecol Oncol* 2004; 93(2): 484-92.
 26. Ma CL, Li YJ, Zhang FC, Wang GQ, Zheng YJ, Kai LM, et al. Detection of HPV16 E6 gene in cervical tissues by quantitative polymerase chain reaction. *Zhonghua Yi Xue Za Zhi* 2004; 84(6): 469-73.
 27. Masumoto N, Fujii T, Ishikawa M, Mukai M, Ono A, Iwata T, et al. Dominant human papillomavirus 16 infection in cervical neoplasia in young Japanese women; study of 881 outpatients. *Gynecol Oncol* 2004; 94(2): 509-14.
 28. Liu J, Rose B, Huang X, Liao G, Carter J, Wu X, et al. Comparative analysis of characteristics of women with cervical cancer in high- versus low-incidence regions. *Gynecol Oncol* 2004; 94, 803-10.