

(AO)

Malgaigne

"

mm
(exact reduction)

Lambotte .()
()

()

%

()

Archive of SID

experimental

Odd's Ratio

()

% /

()

(P< /) . ()

III

(P< /) (Pin tract infection)

(P< /)

() .

	(n=)	(n=)
--	-------	-------

P< /	(/)	II I
	(/)	IIIB IIIA

mm)

P< /	(/)	(%)	mm
	(/)		

%

(P< /)

mm

P< /

> mm

(Odd's Ratio=)

III

(B A)

4. Melendez EM, Colon C. Treatment of open tibial fractures with the Orthofix fixator. *Clin Orthop* 1989; 241: 224-30.
5. Lawyer RB, Lubbers LM. Use of the Hoffmann apparatus in the treatment of unstable tibial fractures. *J Bone Joint Surg* 1980; 62A: 1264-73.
6. Schmidt A, Rorabeck CH. Fractures of the tibia treated by flexible external fixation. *Clin Orthop* 1983; 178: 162-72.
7. Egger EL, Gottsauner –Wolf F, Palmar J, Aro HT, Chao EYS. Effects of axial dynamization on bone healing. *J Orthop Trauma* 1993; 34: 185-92.
8. Helland P, Boe A, Molster AO. Open tibial fractures treated with the Ex-fi-re external fixation system. *Clin Orthop* 1996; 326: 209-20.
9. Henley MB, Chapman JR, Agel J, Harvey EJ, Whorton AM, Swiontkowski MF. Treatment of the type II, IIIa and IIIb open fractures of the tibial shaft: a prospective comparison of unreamed interlocking intramedullary nails and half-pin external fixators. *J Orthop Trauma* 1998; 12: 1-7.
10. Kenwright J, Richardson JB, Cunningham JL, White SH, Goodship AE, Adams MA, et al. Axial movement in tibial fractures. A controlled randomised trial of treatment. *J Bone Joint Surg Br* 1995; 77B: 645-8.

Archive of SID