

1 2 2 *1

86/9/24 :

1
2
*

E-mail:nadermirfakharee@yahoo.com

12

()

1382

12

()

Path analysis of Biomass and Seed Yield Under Three Irrigation Regimes in

Vicia dasycarpa

N Mirfakhraee^{1*}, M Moghaddam², S Aharizad² and A Razban Haghighi¹

¹East Azarbaijan Agricultural Research Center, Tabriz, Iran

²Department of Agronomy and Plant Breeding, Faculty of Agriculture, University of Tabriz, Tabriz, Iran

*Corresponding author:E-mail:nadermirfakharee@yahoo.com

Abstract

This study was conducted to evaluate the effects of different irrigation conditions on 12 genotypes of vetch (*Vicia dasycarpa*) at the Agriculture and Natural Resources Center of East Azarbaijan province,

Iran, during 2003. Field trial was conducted as a split plot design based on randomized complete blocks with three replications. Three irrigation regimes were considered as regular (irrigation at 5-day intervals), water deficit (irrigation at 10-day intervals) and rainfed (no irrigation). Several characters including biomass and grain yield were measured. Path analysis of biomass and grain yield under the above mentioned conditions, using stepwise ridge regression, indicated that some yield components had significant direct effect on these characters. Under regular condition the direct effect of number of grains per pod and number of lateral branches were positive and significant and that of stem length was negative and significant in relation to the biomass. Also the direct effect of number of grains per pod was positive and significant and that of grain density was negative and significant in relation to grain yield. Under water deficit condition the direct effects of number of grains per pod and grain density on biomass and 1000 grain weight and stem length on grain yield were positive and significant. In the rainfed condition because of severe stress, grains were not matured and path analysis was carried out only on biomass. Results showed that in this condition, only number of grains per plant had important effect on the biomass.

Key Word: Biomass, Grain yield, Hairy vetch, Stepwise ridge regression

(*Vicia dasycarpa*)

3

2

1

14

(2004)

pH

8/2

4/9

¹3- isoxazolinonyl

²Allelopathic effects

)
(1371)
(2000)
(1989)
(1999)
12 %50
(1993)
14 24 62
4/95 (1375)
8/4
81 82
246/4 (1998)
7/57 81 82
18/53
1837/2
12 ()
()
4 1382

/1 18 / ... 136
1387

24 3×1/5 1382

8
480

240

116/9

81 82
)

(
893/2

116/9

1384

Archive of SID

10

0/5

(1)

()

(1992)

4/5

80

4/5

(1381)

48

83/3

(1993)

16/1

75

(1995)

%50

(2)

(1984)

1

با عملکرد همبستگی بیولوژیک

اثر مستقیم

صفت

0/473**

0/112

0/280

1/027 **

0/095*

0/011

0/505

0/57*

0/328*

0/011

0/21

0/55*

= 0/740

=0/01

%1 %5

: ** *

2

همبستگی با عملکرد دانه

اثر مستقیم

صفت

0/407

0/076

0/611*

0/492

0/078

0/60*

=0/02

%5

:*

=0/554

%60

(3)

			3		
صفت	اثر مستقیم	همبستگی با بیوماس هوایی			
	0/78	0/055			
	0/77	0/271	0/056		
	%5	=0/01			:*
		=0/617			

%76

(4)

(2000)

Archive of SID

()

()

%40

(5)

4

صفت	اثر مستقیم	همبستگی با عملکرد دانه
	0/69**	0/11
	0/35*	0/570
	%5	=0/01
:** *		=0/76

()

5

0/693	0/685**
=0/01	%1
=0/475	:**

.1381

.54

.1375

.1371

Abdali QN, 1992. Variation in some agronomic characteristics in three populations of chickpea. J. of Amman 8:231-235.

Anonymous, 2004. *Vicia villosa*. Roth. <http://w.w.F.A.O.database/pdf>.

Arraudeau MA, 1989. Breeding strategies for drought resistance. Pp. 107-116. In: Baker FWG (eds). Drought resistance in cereals. CAB International. London, UK.

- Chang CH, 1984. Effects of growth environments on yield and its components in kidney bean (*Phaseolus vulgaris*). Annual Report. 1983-84. Institute of Botany, Academic Sinica Beijing, China.
- Lazcano IF and Lovatt CJ, 1999. Relationship between relative water content, nitrogen pools and growth of *Phaseolus vulgaris* L. and *P. acutifolius* during water deficit. Crop Sci 39: 467-475.
- Ney D and Turk O, 1993. A heat unit based on description of the reproductive development on pea. Crop Sci 33: 510-514.
- Nielsen DC and Nelson NO, 1998. Black bean sensitivity to water stress at various growth stages. Crop Sci 38: 422-424.
- Seppetoglu AH, 1993. A study to determine components affecting seed yield in cowpea (*Vigna unguiculata*). Dogatuk Tarim ve Ormancilik Dergisi 17: 775-784.
- Singh IS, Hassain M and Gupta AK, 1995. Correlation studies among yield and yield contributing traits in F2 and F3 chickpea populations. Inter Chickpea and Pigeonpea Newsletter 2: 11-13.
- Yilmaz VE, 2000. Path analysis of seed yield and biological yield in *Vicia dasycarpa*. w.w.mku.edu.tr/ziraat dergi/2000/12-5. Yilmaz E. pdf.

Archive of SID