(Eucalyptus

camaldulensis) (PAGE)

11:

		(EC , ,	,)	
()			
()	Coleus (Colmant model6/2)			Hepler
		()	:	

¹ Differentiation

		() PAGE	
			()(
	•				
/11		()	
(II)	(ASTM-			
III				. D1	106-84)
111					
	·	c. \supset			
				()
	()	3 3 · · ·)		
			,		
			•		
		()		
			`		
		()		
			()	
			() ()

			%			
					()	
Ŷ/Ŷ¥	-	YV/49	-	٣/١	-	
٣٥/-	-	۲۵/۸۳	-	٣/٧	-	
۴۸/۳	-	Y 0/49	-	٣/٣	-	
۲۳/۳	-	1	-	٣/٢	-	
1	1	1	1	7	1	I
/	1	1	1		1	II
1	1	1	1		1	III
	1	1	1	1	1	1
/	1	1		1	1	
/	1	1	1	1	1	
	1				1	
	1		V		1	
	1				1	

- 1-Ebermann, R., Korori, S.A.A. and Lickl, E. 1991. Temperature dependent alteration of peroxidase and amylase isoenzymes in Quercus robur. Phyton 31:121-128.
- 2-Gaspar, Th., Penel, C.C., Castukki F.J. and Greppin, H. 1985. A Two step control of basic and acidic peroxidase and its sighificance for growth and development. Physiol. Plant. 64:418-423.
- 3-Gaspar, Th., Penel C., Thorpe A, and grepping h.1982. A surbey of their biochemical and physiological rolesof higher plauts peroxidase 1970-1980. Univ. Geneva, centre do botanique. Geneva 324pp.
- 4-Hepler P.K., Fosket. D.E. and newcomb, E.M. 1970. Lignifiation during secondary wall formation in cokeus: An electron microscope study A.m. J.Bot. 57, 85-96.
- 5-Higuchi. T. 1982. Biosynthesis of lignin in Biosynthesis and biodegradation of wood comporeuts ed. T. Higuch: Academic press.
- 6-Saka S. and Goring D.A. I 1985. Localization of lignins in wood cell walls in Biosynthesis and biodegradation of wood comporents (ed. T. Higuch) Academic press. Orlando. PP.51-62.

Enzymatical Studies related to wood production in Eucolyptus

N. Toghraie¹

D. Parsapajouh²

A.Hosseinzadeh³

S.A.A Korori⁴

Abstact

To make a primary investigation on lignin content variation related to content of peroxidase enzyme in different seasons and sites, We sampled from four regions in two season, The eucalyptus (Eucalyptus camaldulensis) trees.

Qualitative and quantitative survey was done PAGE and spectrophotometry methods.

The Content of Klason lignin was determined using extractive free wood sawdust.

The results show that peroxidase isoenzymes patterns change due to region and season. This can be understood as a part of tree physiological preparing period against coldness. It was also observed that there are differences between ecological regions and between trees in the same region. Genetically the lignin content in Khuzistan region decreased in winter and so did enzyme activity. The content of lignin was related directly to branch diameters and enzyme activity.

Keywords: Lignin, Peroxidase, PAGE, Lignification, Eucalyptus Camaldulensis

¹ -Research inst. Of Forests and Rangelands ² -Culty of Natural , University of Tehran, Karaj ³ -