
)

() (

()

p< /

()

p< /

p< /

()

Archive of SID

(*Sorbus torminalis*)

()

,Bassi Piagnani)

,Lanier ,

Demersure

, Tsitsa-txardi) (

(

Sorbus

.(,Demersure)

.(Anonymus)

.(Zare ,)

.()

() Meyer

() Asthalter () Ivenko

Razumova () Bassi Piagnani

()

()

() Palashev Lyapova .

() Yagihasi .

Cecil)

.(Fare

()

)

-) (

(

.(Meyer)

t

t-test

)

p<

p<
p<

()

Strouts)

(,McDonald
(,Pepper)

Polycyclism

...

()

					/	
				/	** /	
			* /	/	* /	
		** /	** /	/	** /	
	** /	** /	** /	/	** /	

*

**

()

$p < ,$

Kolmogorov- Smirnov

$F = ,$)

t-test

($p < ,$

()

Kolmogorov- Smirnov

()	()	()	()	()	
/	/	/	/	/	K - S Sign
ns /	ns /	ns /	ns /	ns /	
/	/	/	/	/	

ns

ns / /	ns / /	ns / /	* / /	/	F Sign

ns

*

T

()

p<0.05

()

(p<0.01)

()

()

()

p<0.01

(T)

Sign	t	()			
/	* /	/	/	()	
/	* /	/	/	()	
/	/ **	/	/	()	
/	** /	/	/	()	
/	** /	/	/	()	

/ () /

	()		()		()		
	/	/	/	/	/	()	
	/	/	/	/	/	()	
	/	/	/	/			

() p<0.05

() p<0.05

() Sanches Casal .

() p<0.01

() ABA

()

() Fare Cecil .

Schmidtlin . ()
()
()

Barnola) / /
Sagheb-Talebi Roloff
(

() Hussendoerfer

() Tacon ()
()

()

- ()

()

()
(*Pistacia vera* L)

Archive of SID

-
- 8-Anonymus., 1967. Propagation from seed. Pt. 8: Fall planting of fruit and hardwood seeds. Am. Nurseryman 126(4): 12-13, 85-90.
- 9-Asthalter, K., 1980. Causes and site-related occurrence of drought and possible influences for tree species. Allgemeine-Forstzeitschrift 19: 510-512.
- 10-Barnola, P., Alatou, D., Parmentier, C., Vallon, C. 1993. Approche du déterminisme du rythme de croissance endogène des jeunes chênes pédonculés par modulation de l'intensité lumineuse. Ann. Sci. For. 50: 257-272.
- 11-Casal, j., Sanches, R.A. 1998. Phytochromes and seed germination. Seed Science Research, 8: 317 – 329.
- 12-Cecil, P., Fare, D. 2002. Effect of seed source on first year Growth of *Quercus phellos* and *Q. shumardii*. SNA Research conference proceedings vol.47:295- 299.
- 13-Demesure, B., Guerroue, BL., Lucchi, G., Part, D., Petit, RJ. 2000. Genetic variability of a scattered temperate forest tree: *Sorbus torminalis* L. Annual Forest. Science 57: 63-71.
- 14-Hussendoerfer, E., 1992. Untersuchungen zur Verzweigungsform und anderen phänotypischen Merkmalen der Rotbuche (*Fagus sylvatica* L.) unter besonderer Berücksichtigung über zusammenhaenge mit ausgewählten Merkmalen. Dipl. Arbeit d. Forstwiss. Fak. Der Albert-Ludwig-Universitaet. Freiburg i. Br. 82 p.
- 15- Ivenko, Sl., 1952. *Sorbus torminalis* – A valuable species for planting in the steps. Lesn-Hoz 5: 7-35.
- 16- Lanier, N., 1993. Le boom sur le marche de l'Alisier torminalis Rev.for.fr.XIV, P:319.
- 17- Le Tacon, F., 1983. La plantation en plein découvert: une des causes de la mauvaise forme du hêtre dans le nord-est de la France. Rev. Forest. Française.XXXV. 6: 452-459.
- 18-Lyapova, I., Palashev, I. 1988. Growth of seedling of *Platanus orientalis*, *Sorbus torminalis* & *Corylus avellana* to light, Gorsko-Stopavstvo, 44, 1: 24-25.
- 19- McDonald, S. E., 1984. Irrigation in Forest –Tree Nursery: Monitoring and Effects on Seedling Growth. In: Duryea, M.L and Landis, T.D. Forest Nursery Manual U.S.D.A Forest Service 384P. 107-122.
- 20-Meyer, U., 1980. Growing *Sorbus torminalis* in Forest District Grohnde Landes Forstverwaltung 33: 184-193.
- 14- Pepper, H.W., 1992. Forest fencing, Forestry Commission Bulletin 102, HMSO, London.
- 21- Piagnani, C., Bassi, D. 2000. In vivo and In vitro propagation of *Sorbus* ssp.from juvenile material, Italus Hortus 7: 3-7.
- 22- Razumova, Mv., 1987. Biology of seed germination in species of the genus of *Sorbus*. Botanicheskii Zhurnal 72: 77-83.
- 23- Roloff, A., 1985., Morphologie der kronenentwicklung von *Fagus sylvatica* L. (Rotbuche) unter besonderer Berücksichtigung moeglicherweise neuartiger Veraenderungen. Diss. Georg-August-Univ. Goetingen. 161p.
- 24- Sagheb- Talebi, K., 1995. Study of some characteristics of young beeches in the regeneration gaps of irregular shelterwood system (Femelschlag). In: Madsen, S. F. (Herg.) Genetics and silviculture of beech. Denmark. Forskningsserien Nr. 11: 105-116.
- 25- Schmidting, Rc., 1994. Using provenance test to predict response to climatic chang: Lobolly pine and Norway spruce . Tree physiology 14 : 805-817.

- 26- Strouts, R. G., Gregory, S. C., Heritage, S. G., 1994. Protection against climatic damage, fungal diseases, insects and animal pests. In: Aldhous, J.R and Mason. W.L. Forest nursery practice, Forest Commission Bulltin 111. HMSO, London 268p. 181- 194.
- 27- Tsitsa–Tzardi, E., Loukis, A., Philianos, S. 1991. Constituent of *Sorbus torminalis* fruits. *Fitoterapia* 62: 282-283.
- 28-Tsitsa–Tzardi E., Loukis, A., Philianos, S. 1992. Constituents of *Sorbus torminalis* leaves. *Fitoterapia* 63: 189-190.
- 29-Yagihashi, T., Hayashida, M., Myamoto, T. 1998. Effect of birds ingestion on seed germination of *Sorbus comixica*. *Oecologia* 114: 209-212.
- 30-Zare, H., Tabari, M., Espahbodi, K. 2002. Ecological characteristics on quantitative condition of *Sorbus torminalis* (a case study in the northern forests of Iran) International Congress of Ecology. Seoul, Korea August 11-18 .

Archive of SID

Effect of Seed Source Altitude in Wild Service Tree, on Seed Germination

K. Espahbodi¹

H. Mirzaie – Nodoushan²

M. Tabari³

M. Akbarinia³

Y. Dehghan Shooraki²

Abstract

In order to determine the effects of *Sorbus torminalis* seed origin altitude on the germination of seed, 200 seeds of 40 individual wild service trees from two origins, namely Ashak (2100-2200m a.s.l) and Sangdeh (1700-1800m a.s.l), were planted in plastic flower-vases, in a nursery at 1500 m a.s.l. Germination of seed, survival and height growth of seedlings were recorded during the first year. Analyse is of data showed that, the effect of seed source altitude on seed germination was significant ($p<0.05$). The effect of seed source altitude on seedling survival ($p<0.05$) and the height growth of seedlings ($p<0.01$) was significant too. Germination rate of seeds collected from higher altitudes alttitude(Ashak) was higher than the germination of seed from lower altitude (Sangdeh). Despite that, the survival and height growth in seedlings of seeds from lower altitude was more than those in seeds collected from higher altitudes. It is therefore suggested to collect seeds from the nearest origin to the nursery.

Keywords: *Sorbus torminalis*, Germination, Site altitude, Survival, Height growth, Nursery

1-Scientific Staff Member, Natural Resources Research Center, Mazandaran, Ph.D. Student at Forestry Dpt of Tarbiat Modarres Universit E-mail: espahbodi2002@yahoo.com

2- Scientific Staff Member, Forests and Rangelands Research Institute

3- Scientific Staff Member, Natural Resources Faculty, Tarbiat Modarres University