
Archive of SID

(/ / : / / :)

pH

(PCA)

(PCA)

:

...

(Gypsids)

/ (Salids)

.()

() .()

()

Halocnemum strobilaceum

()

Seidlitzia

Aeluropus littoralis rosmarinus

Atriplex veruciferum

()

()

()

()

()

()

.()

()

pH EC

()

Alphen and Romero

Smith and Robertson

()

)

(

Weak Aridic

()

()

pH

Fabaceae

Chenopodiaceae

Poaceae

Asteraceae

Archive of SID

)

(

()

(

)

West and Ibrahim
et al. Abd El-Ghani

(eigenvalue)

()

()

(EC)

()

pH PW 9527

EC

EYELA 2000

pH

Walkley & Black

()

EDTA

()

(USDA 2003)

Fine loamy gypsic thermic Typic

(PCA)

Clayey gypsic thermic Calcigypsid

PC-ORD

GypsicAquisalids

PCA (ver. 4.7)

PCA

(eigen vector)

(eigen value)

sp.

Sasp.)Halocnemum strobilaceum-Salsola

(eigen vectors)

Halocnemum strobilaceum (Hast-

() *Salsola sp.*

Seidlitzia rosmarinus:

Principal Component Analysis

	<i>Hordeum</i>	<i>Zigophyllum fabago</i>
<i>Tamarix sp.-</i>		<i>morinum</i>
<i>Nitraria schoberi (Tasp.-Nisc)</i>		
<i>Aeloropus littoralis:</i>	<i>Seidlitzia</i>	
<i>Halocnemum Artemisia sieberi</i>	<i>Haplophyllum glaberrimum</i>	<i>rosmarinus</i>
<i>Hordeum Salsola crassa strabilaceum</i>	<i>Haplophyllum</i>	(Hagl-Sero)
<i>.Chenopodium album morinum</i>		<i>glaberrimum</i>
		<i>Seidlitzia rosmarinus</i>
	<i>Lycium</i>	:
		<i>Artemisia sieberi ruthenicum</i>
	<i>Ducrosia sp.</i>	<i>Zigophyllum fabago</i>
	<i>Hyosiamus niger</i>	<i>Eryngium bungei</i>
		<i>Scariola orientalis,</i>
	<i>Astragalus spp.</i>	<i>Eremurus sp.</i>
	<i>Alhagi</i>	<i>Annual Grasses</i>
		<i>Salsola sp. camelorum</i>
		<i>Anchusa sp. Halocnemum strabilaceum</i>
	<i>Salsola rigida-</i>	
		(Sari-Peau) <i>Petropyrum aucheri</i>
		<i>Salsola rigida</i>
		() <i>petropyrum aucheri</i>
		:
		<i>Ziziphora tenuis, Artemisia sieberi,</i>
		<i>Acanthophyllum sp., Peganum harmala,</i>
		<i>Stipa barbata, Echinops robustus ,</i>
		<i>Centaura depressa, Astragalus glacanthus,</i>
		<i>Scabiosa sp, Cousinia sp., Allysum</i>
		<i>strigosum, Hordeum morinum, Noea</i>
		<i>mucronatha Lotus corniculatus,</i>
		<i>Eremopyrum sp. , Bromus tectorum,</i>
		<i>Onopordon acanthium, Tavatenia sp.,</i>
		<i>Scariola orientalis,...</i>

()

/	/	/	
/	/	/	%
/	/	/	

Hagl-Sero HaSt-Sasp.

Halocnemum

strobilaceum

Sari-Ptau

Tasp.-Nisc

()

)

(

)

(

Aeloropus sp.

) *Halocnemum sp.*

(

() () ()

()

Halocnemum strobilacei

()

:()

(Ph.D)

:()

:()

:()

16- Abd El-Ghani, M.M., Wafaa M. Amer.2003. soil – vegetation relationships in coastal desert plain of southern Sinai,Egypt. Journal of Arid environment 55(607-628)

17- Alphen Van, G.J and De los Rios Romero F.1971. Gypsiferous soils, note on characteristics and management. International institute for reclamation and improvement. - Bulletin 12/ Wageningen/ The Nethelands.

-
- 18- FAO, 1990, Management of gypsiferous soils, Bulletin 62.FAO, ROME FAO,1990.
- 19 - Hesse, P. R (1976) Particle size distribution in gypsic soils. Plant and Soil.44,241-247
- 20-Jafari.M, Zare chahouki M.A,Tavili.A, Azarnivand.H, Zahedi Amiri.GH(2004) Effective Environment Factors In The Distribution Of Vegetation Types In Poshtkouh Rengelds Of Yazd Province (Iran), J. Arid Environments 56 (627-641)
- 21-Jafari.M, Zare chahouki M.A,Tavili.A, Azarnivand.H,Tavili A, 2003. soil – vegetation relationships in Hoz-e-soiltan region of Qom Province, Iran. Pakistan Journal of nutrition, 2(6) 329-334
- 22- Jongman,R.H.G. and O.F.R., Van Tongeren, 1995. Data analysis in community and landscape ecology. Center fire agricultural publishing and documentation.
- 23-Smith, r & Robertson V.S. 1962. Soil irrigation classification of shallow soil overlying gypsum beds northern Iraq. Journal of Soil Science 13: 106-115
- 24-Soil Survey Staff. 1993. Soil Survey Manual. U.S..Dep. agr. Handbook 18.U.S. Gouv Printing office. Washington.D.C
- 25- USDA, 2003. keys of soil taxonomy,soil survey staff, 9th edition.
- 26- USDA, 2004. Soil Survey Labrotoary Methods And Precedures For Collecting Soil Samples.
- 27-West, N.E. and K.I. Ibrahim, 1967. Soil –vegetation relationship in the shad scale zone of southern Utah ECology 49: 45-456

Archive of SID

Soil-vegetation relationships in saliferous and gypsiferous soils in winter rangelands (Eshtehard)

A. Ahmadi^{*1}, Gh. Zahedi Amiri², Sh. Mahmoodi³ and E. Moghiseh⁴

¹ member of scientific board of Islamic Azad university of Arak, I. R. Iran

² Associate prof, Faculty of Natural Resources, University of Tehran, I. R. Iran

³ Professor, Department of Soil Science, University of Tehran, I. R. Iran

⁴ Ph.D. Student of Soil Science, University of Tehran, I. R. Iran

(Received 2006 Nov 5, Accepted 2008 Janu 7)

Abstract

Saliferous and gypsiferous aridisols are among the vast extension soils of rangelands in arid and semi-arid regions of Iran. Vegetation cover with low density is representative of special physicochemical characteristics of these soils. The aim of the present study was to identify gypsophilous and halophilic plants and also to investigate plants distribution related to physicochemical characteristics of saliferous and gypsiferous in Eshtehard winter rangelands. Vegetation sampling was conducted in the key area based on randomized-systematic pattern, and data including canopy cover and density along transects in each plot were determined. After determination of plant types, the soil profiles were excavated and soil samples collected. Based on the standard methods, physicochemical characteristics, including soil texture, electrical conductivity in saturated extract, soil reaction, organic carbon content, gypsum and equivalent calcium carbonate percentage, soluble anions and cat ions were determined. Multivariate method of principal component analysis (PCA) was used to analyze the collected data. The results showed that the vegetation distribution pattern was mainly related to soil characteristics such as salinity, texture, chloride, soil reaction, gypsum and gravel percentage.

Key words: saliferous and gypsiferous soils, principal component analysis, cover, physicochemical characteristics, Eshtehard

* Corresponding author: Tel: 0912-4522394 , Fax: 0261-2249313 E-mail: bsahmadi@yahoo.com