

(Nigella sativa)

\*\*\*

\*\*

\*


\*

\*\*

\*\*\*

// :

// :



.()

.()

.()

LDL

mg/dl . ( ) HDL

( )

( )

)

( )

(S TZ: Streptozotocin)

( )

STZ

mg/dl (Nigella Sativa)

( )

(n=44) Wistar

( )

( ) Spectronic 20 °c

Mean±S.E.M.

repeated measure ANOVA /

student's paired t-test

One-way

Tukey's Post-hoc test ANOVA

P<0.05

)

( )

( )

( )

(P<0.01)

/ / /

...(Nigella sativa)

(P<0.001)

.( ) (p<0.01)

p<0.05)

.(p<0.05)

(p<0.01

(p<0.05)

( )

(P<0.001)

.(p<0.05)

(p<0.01)

	Body weight (g)			Serum GLUCOSE (mg/dl)		
	Week 0	Week 4	Week 8	Week 0	Week 4	Week 8
	295.2 ± 5.1	309.4 ± 6.2	323.1 ± 6.9	131.5 ± 13.05	126.7 ± 11.08	123.4 ± 9.7
+	307.8 ± 7.6	314.5 ± 9.2	320.1 ± 6.7	159.5 ± 25.2	131.2 ± 9.5	137.6 ± 7.1
	287.2 ± 6.6	234.1 ± 10.3*	207.1 ± 9.8**	128.9 ± 6.8	409.4 ± 8.4***	401.3 ± 8.4***
+	300.7 ± 7.6	259.1 ± 10.5	238.1 ± 10.6*	124.8 ± 10.1	351.8 ± 14.1***	359.6 ± 20.7***

\* P<0.05, \*\* P<0.01, \*\*\* P<0.001 ( )

NS = Nigella sativa; week 0 =

	Total Cholesterol (mg/dl)			Triglyceride (mg/dl)		
	Week 0	Week 4	Week 8	Week 0	Week 4	Week 8
	59.1 ± 3.9	56.4 ± 5.7	55.4 ± 4.3	101.7 ± 6.8	112.3 ± 7.3	98.9 ± 6.4
+	47.1 ± 4.9	49.2 ± 4.1	45.8 ± 4.9	80.7 ± 6.5	75.4 ± 5.6	74.9 ± 6.3
	54.8 ± 6.1	75.6 ± 5.8**	74.9 ± 6.1**	99.2 ± 6.3	122.4 ± 6.4*	124.1 ± 5.7*
+	50.2 ± 5.8	62.5 ± 5.6*	53.8 ± 4.3*	100.2 ± 7.1	109.1 ± 8.3	102.7 ± 7.1

\* P<0.05, \*\* P<0.01 ( )

NS = Nigella sativa; week 0 =

Archive of SID

1. Tripathi BK, Srivastava AK. Diabetes Mellitus: Complications and Therapeutics. Med Sci Monit 2006;12:RA130-47.

2. Gleckman R, Mory J. Diabetes-Related Foot Infection. Journal of Contemporary Internal Medicine 1994; 6: 57-62.

3. Swanston- Flatt SK, C, Bailey CJ, Flatt PR. Evaluation of Traditional Plant Treatments for Diabetes: Studies in Streptozotocin Diabetic Mice.

4. Dahri AH, Chandiol AM, Rahoo AA, Memon RA. Effect of Nigella Sativa (Kalonji) on Serum Cholesterol of Albino Rats. J Ayub Med Coll Abbottabad 2005; 17:72-74.

5. Vahdati-Mashhadian N, Rakhshandeh H, Omidi A. An Investigation on LD50 and Subacute Hepatic Toxicity of Nigella Sativa Seed Extracts in Mice. *Pharmazie* 2005; 60:544-547.
6. Ilhan A, Gurel A, Armutcu F, Kamisli S, Iraz M. Antiepileptogenic and Antioxidant Effects of Nigella Sativa oil Against Pentylentetrazol-Induced Kindling in mice. *Neuropharmacology* 2005;49: 56-64.
7. Farah N, Benghuzzi H, Tucci M, Cason Z. The Effects of Isolated Antioxidants from Black Seed on the Cellular Metabolism of A549 Cells. *Biomed Sci Instrum* 2005;41:211-6.
8. Mansour M, Tornhamre S. Inhibition of 5-Lipoxygenase and Leukotriene C4 Synthase in human blood cells by thymoquinone. *J Enzyme Inhib Med Chem* 2004;19:431-436.
9. Rchid H, Chevassus H, Nmila R, Guiral C, Petit P, Chokairi M, Sauvaire Y. Nigella Sativa Seed Extracts Enhance Glucose-induced Insulin Release from Rat-Isolated Langerhans Islets. *Fundam Clin Pharmacol* 2004; 18:525-529.
10. Suboh SM, Bilto YY, Aburjai TA. Protective Effects of Selected Medicinal Plants Against Protein Degradation, Lipid Peroxidation and Deformability Loss of Oxidatively Stressed Human Erythrocytes. *Phytother Res* 2004;18:280-284.
11. Ramadan MF, Kroh LW, Morsel JT. Radical Scavenging Activity of Black Cumin (*Nigella Sativa* L.), Coriander (*Coriandrum sativum* L.), and Niger (*Guizotia Abyssinica* Cass.) Crude Seed Oils and Oil Fractions. *J Agric Food Chem* 2003; 51:6961-6969.
12. Ali BH, Blunden G. Pharmacological and Toxicological Properties of Nigella Sativa. *Phytother Res* 2003; 17: 299- 305.
13. Yildirim O, Buyukbingol Z. Effect of Cobalt on the Oxidative Status in Heart and Aorta of Streptozotocin-Induced Diabetic Rats. *Cell Biochem Funct* 2003; 21:27-33.
14. Stoenoiu MS, De Vriese AS, Brouet A, Moulin P, Feron O, Lameire N, Devuyst O. Experimental Diabetes Induces Functional and Structural Changes in the Peritoneum. *Kidney Int* 2002; 62:668-678.
15. Martin-Nizard F, Shapaz S, Furman C, Fruchart JC, Duriez P, Billeul F. Natural Phenylpropanoids Protect Endothelial Cells Against Oxidized LDL-Induced Cytotoxicity. *Planta Med* 2003;69:207-211.
16. VanderJagt TJ, Ghattas R, VanderJagt DJ, Crossey M, Glew RH. Comparison of the Total Antioxidant Content of 30 Widely Used Medicinal Plants of New Mexico. *Life Sci* 2002; 70:1035-1040.
17. Jelodar Gholamali A, Maleki M, Motadayen MH, Sirius S. Effect of Fenugreek, Onion and Garlic on Blood Glucose and Histopathology of Pancreas of Alloxan-Induced Diabetic Rats. *Indian J Med Sci* 2005; 59:64-69.
18. Karawya MS, Wahab SAA. Diphenylamine, an Antihyperglycemic Agent from Onion and Tea. *Journal of Natural Products* 1984; 47: 775-780.
19. Shimizu M, Ito T, Terashima S, Mayashi T, Arisawa M, Morita N, Kurokawa S, Ito K, Hasimato Y. Inhibition of Lens Aldose Reductase by Flavonoids. *Phytochemistry* 1984; 23: 1885-1888.

---

## Survey The Hypoglycemic and Hypolipidemic Effect of Chronic Oral Administration of Nigella Sativa in Diabetic Rat

Roghani M.(Ph. D.), Baluchnejadmojarad T.(Ph.D.), Roghani-Dehkordi F.(MD.)

### Abstract

**Introduction:** There are some reports about Anti-Diabetic effect of black seed in Islamic and traditional medicine. Therefore, hypoglycemic and hypolipidemic effect of this medicinal plant was investigated.

**Objective:** Survey the hypoglycemic and hypolipidemic effect of chronic oral administration of Nigella Sativa in diabetic rat.

**Materials and Methods:** Male Wistar rats (n=44) were randomly divided into 4 groups, including control, black seed-treated control, diabetic, and black seed-treated diabetic. For induction of diabetes, Streptozotocin (STZ; 60 mg/Kg; i.p.) was used at a single dose intraperitoneal. A serum glucose level higher than 250 mg/dl was considered as diabetic state. The treatment groups received oral administration of black seed-mixed pelleted food (6.25%) for two months.

**Results:** The results showed there were significant increase in serum glucose level in diabetic group 4 and 8 weeks after intervention as compared with the week before periment( $P<0.001$ ), Although there were no significant difference between same groups at the same time Nigella Sativum treatment of diabetic rats did not show any significant effect. In addition. Also, triglyceride level in diabetic group increased 8 weeks after intervention in comparison with related data one week before the study ( $P<0.05$ ) and there was a significant lower level of triglyceride in Nigella Sativum-treated diabetic rats ( $p<0.05$ ). Furthermore, a similar high significant reduction was obtained for black seed treated-diabetic group as compared to diabetic group regarding serum cholesterol level ( $p<0.05$ ).

**Conclusion;** The obtained results demonstrated that oral chronic administration of Nigella Sativum could significantly reduce Serum Triglyceride and Cholesterol Level in Diabetic rats and this may reduce some Diabetic complications.

**Key words:** Cholesterol/ Diabetes Mellitus/ Hypoglycemia/ Nigella sativa/ Rats/ Triglycerides