

:
/ / :
/ / :

± /
/ ± / / ± / , / ± / , /
()
() -
B A (/ ± /)
(/ ± / - / ± /)
(P < /) -

Email : sbolboli@yahoo.com



Archive of SID

(.)

()

(.)

()

(.)

(.)

1- Graded Exercise Test

Archive of SID

(B A)

1- Cardiovascular Fitness Ratio



()

()

(**Seca**)

(**ACSM**)

()

()

()

1- American College of Sport Medicine

C° %)) (

CFR = / - / * ()

(C)

CFR = / - / * ()

(B)

(A)

$$\mathbf{A: CFR} = \frac{\text{---}}{\text{---}} \times \text{---} \quad (\text{---})$$

$$\mathbf{B: CFR} = \frac{\text{---}}{\text{---}} \times \text{---} \quad (\text{---})$$

)

()

-

(r^2 r ,SEE

()

-


MedCalc(8.2.1.0)

(/)	()	()	()
/ ± /	/ ± /	/ ± /	/ ± /




A) , (,)

(**B**

- 1- Agreement
- 2- Bland – Altman (1986)




/	/	/	/ /	/	-C
/	/	/	/ /	/	-A
/	/	/	/ /	/	-B



() A ()
 (X) () (B)
 (y) () - ()

()
() (%)



Archiv


$$- = / - / * ()$$

A

$$- = / - / * ()$$

B

$$- = / - / * ()$$



()B

()A

()C

-

% , -
()

-

	%		
P< /	/ /	/ /	C
P< /	/ /	/ /	A
P< /	/ /	/ /	B

T , -
()

	T		
P< /	/ /	/ /	C
P< /	/ /	/ /	A
P< /	/ /	/ /	B

-

- (C)

()

- % .()

- (C)

.()

(C)

-

—
—————

()

—

(, , , ,)

A

()

A

A

—

—

—

()

—

Archive of SID

"()

:"(—)

-
3. American College of Sports Medicine. (2000). "ACSM'S guidelines for exercise testing and prescription (6th ed)". Philadelphia PA: Lippincott Williams & Wilkins :91.
4. American College of Sport Medicine. (1990). "ACSM'S guidelines for exercise testing and prescripting. 4th edition, lea and febiger, P:148.
5. Arrais GA, Wigle DT, and Mao Y. (1992). "Risk assessment of physical activity and physical fitness in the Canadian Health survey follow up study". *J Clin Epidemiol* ; 45: PP:419-28.
6. Blair S.N. Kohl H.W. (1989). "Physical fitness and all cause mortality : a prospective study of Health men and women". *J of the American Medical Association*; 262 : PP:2395-2401.
7. Blair , S.N. Kohl, H.W., Gordon, N.F., & Paffenbarger, R.S. (1992). "How much physical activity is good for health? Annual Review of public Health, 13, PP:99-126.
8. Blair S.N., Ming, W., and chong, D.L (1998). "Cardiorespiratory fitness determined by exercise heart rate as a predictor of mortality in the aerobic center longitudinal study". *J Sport sci*; 16, PP: S47-S55.
9. Bland, M.J. & D.G. Altman, (1986). "Statistical methods for assessing agreement between two methods of clinical measurement". *The Lancet*, PP:1307-310.
10. Bradshaw DI, George JD, Hyde A, Lamonte MJ, Vehrs PR, Hager RL, Yanowitz FG. (2005). "An accurate VO₂max nonexercise regression model for 18-65-year-old adults". *Res Q Exerc Sportt.*;76(4) : PP:426-32.
11. Byrne NM, Hills AP, Hunter GR, Weinsier RL, Schutz Y. (2002). "The metabolic equivalent : One size does not fit all". *J Appl Physiol*. 14; [Epub ahead of print].
12. Donald C. Dibloo and Jeffery K. Moffit (2003). "A comparison of bioelectrical impedance and NEAR-Infrared Interact ance to skinfold measures in determining minimum wrestling weight in collegiate wrestlers". *International Electronic Journal Vol. 6. No.2* : PP:26-36.
13. Ekelund L.G., Haskell WL. (1988). "Physical fitness as a predictor of cardiovascular mortality in asymptomatic North American men". *N Eng J Med*, 319 : PP:1379-84.

-
14. Franz X. Kleber . (2004). "The predictive value of cardiorespiratory fitness". *European Heart Journal* 25, PP: 1374-1375.
15. Gibbons L, Blair SN, Kohl HW, Cooper K (1989). "The safety of maximal exercise testing". *Circulation*; 80 : PP:846-852.
16. Hien H.O. (1992). "Physical fitness or physical activity as a predictor of ischemic heart disease". *J intern Med*, 232; PP:471-9.
17. James D.G., Pat R.V. & Garth J.B. (2000). "A modified submaximal cycle ergometer test designed to predict treadmill VO_{2max} , Vol, 4, PP:229-243.
18. Lie H. Mundal R. and Erikssen J. (1985). "Coronary risk factor and incidence of coronary death in relation to physical fitness". *Eur heart J*, 6 : PP:147-57.
19. Seefeldt V, RM Malina, and MA Clark. (2002). "Factors affecting levels of physical activity in adults". *Sports Med.* 32; PP:143-168.
20. Slattery, M.L. Jacobs, D.R.Jr. (1988). "Physical fitness and cardiovascular disease mortality"; the US railroad study *Am J Epidemiol*, 127;PP:601-10.
21. Spangler, R.D. (1970). "A submaximal exercise electrocardiographic test as method of determining occult ischemic heart disease". *Am Heart J* 80: PP:725-58.
22. Thomas, J.R.& J.K. Nelson . (1990). "Research methods in physical activity". 2th ed, Chmpaign : *Human Kinetics*. PP:129-178.
23. Wei M.JB Kamper T, CE Barlow, MZ Nichaman, LW Gibbons, RS Paffenbarger Jr, ans SN Blair. (1999). "Relationship between low cardiorespiratory fitness and mortality in normal-weight, overweight, and obese men". *JAM A.* 282; PP:1547-1553.
24. Young DZ, Lampert S, Graboys TB, Lown B. (1984). "The safety of maximal exercise testing in patients at high risk for ventricular arrhythmia; 70;PP:184-191.