

\*

/ / : // :

## Inhibitory effect of sodium selenite on ferric nitrilotriacetate-induced oxidative stress in mice kidney tissues

Rezazadeh H.\*<sup>1</sup>, Habibi Asl B.<sup>1</sup>, Hassannajad S.<sup>1</sup>, Somi M.H. <sup>2</sup>

<sup>1</sup>Department of Pharmacology and Toxicology School of Pharmacy, <sup>2</sup>Liver and Gastrointestinal diseases Research Center, Tabriz University of Medical Sciences

Received: 2005/19/15 , Accepted: 2007/1/16

**Objectives:** Intraperitoneal injection of ferric nitrilotriacetate (Fe-NTA) to laboratory animals leads to oxidative stress, lipid peroxidation (LPO) and tissue injuries. However, recent reports indicate that selenium can inhibit oxidative stress, and delays the benign tumor conversion into malignant carcinoma. In this study we showed that selenium can inhibit the harmful effects of Fe-NTA in mice kidneys. **Methods:** Female albino swiss mice were divided into 11 groups. Group-I received saline normal and served as control. Different doses of selenium (0.5, 1, 1.5 mg /kg) were injected before and after Fe-NTA treatment. Also the groups of animals received either selenium (0.5, 1, 1.5 mg/kg) or Fe-NTA (9 mg Fe/kg) alone. The selenium was injected daily for a period of 21 days. All the animals were killed and the kidneys were taken out. The biochemical estimation such as protein and LPO were carried out on the kidney tissue. **Results:** selenium at the dose of 0.5 mg/kg inhibits the LPO induction-induced by Fe-NTA. Other doses of selenium (1 and 1.5 mg/kg) were not effective on the inhibition of LPO. Our data indicate that selenium is more effective before Fe-NTA treatment. **Conclusion:** The results indicated that the inhibitory effect of selenium may act through different mechanisms and involvement of selenium in the diminishment of LPO may be due to inhibition and scavenging harmful free radicals.

**Key words:** Ferric nitrilotriacetate, Sodium selenite, Oxidative stress, Lipid peroxidation.

( / / mg/kg) ( / / mg/kg) ( mg Fe/kg) / mg/kg / mg/kg

\*Corresponding Author: Dr Hassan Rezazadeh, Assistant Professor, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tel: 0411- 3341315; Fax:0 411 - 3344798; E-mail: rezafar81@hotmail.com

( )

( )

( )

( )

(Sigma )

DNA

( )

)

( Merck

(Albino Swiss)

( )

±

( )

)

(

( )

( )

:(Fe-NTA)

( )

Awai

( )

( )

PH=7.40

cc (Mohandas 1984)

% cc

RPM

nmol (MDA)

Polytron RPM

$M^{-1}cm^{-1}$

Dunnets

P < 0.05 Variance

,In vivo

I


II

V IV ,III

VIII VII ,VI

(Saline)

XI ,X ,IX



Mean±Se

\* a P < 0.05 ; \*\*P < 0.05 ;

( )

wright et al.


/ cc

/ M

/ cc

/

(Saline)




/

/ mg/kg

mg/kg

/ mg/kg



Archive of SID

Dose mg/kg	Saline	Sodium selenite	Fe-NTA+	
			Sodium selenite	Fe-NTA
0	0	0	0	0
0.5	0	0	0	0
1	0	2	3	3
1.5	0	4	3	2

.( )

.( )

.( )

.( )

(Marker) 8-OH-dG DNA  
 DNA ( , )  
 OH  
 ( ) ( ) Eltayeb

S

( )

/

( )  
 (O )

DNA (Cross-Links)  
 (8-OH-dG)

## 6- References:

- 1- HU M.L., Louie S., Cross C.E., Motchnik P., Halliwell B. Antioxidant protection against hypochlorous acid in human plasma. *J Lab Clin Med.*, 1993, 121 (2): 257-262.
- 2- Rosa A., Deiana M., Corona G., Atzeri A., Incani A., Dessi MA. Cholesterol as target Fe-NTA-induced lipid peroxidation in rat tissues. *Toxicol Lett.*, 2005, 157(1): 1-8.
- 3- Khan N., Sultana S. Chemomodulatory effect of *Ficus racemosa* extract against chemically induced renal carcinogenesis and oxidative response in wistar rats. *Life Sci.*, 2005, 77 (11): 1194-1210.
- 4- Lipinski B. Rationale for the treatment of cancer with sodium selenite. *Med Hypotheses*, 2005, 64 (4): 806-810.
- 5- Igbal M., Rezazaseh H., Ansar S., Athar M. Alpha -Tocopherol (Vitamin-E) ameliorates ferric nitrilotriacetate (Fe-NTA)-dependent renal proliferative response and toxicity: diminution of oxidative stress. *Hum Exp Toxicol.*, 1998, 17 (3): 163-171.
- 6- Adamowicz A., Trafikowska U., Trafikowska A., Zachara B., Manitius J. Effect of erythropoietin therapy and selenium supplementation on selected antioxidant parameters in blood of uremic patients on long-term hemodialysis. *Med Sci Monit.*, 2002, 8 (3): 202-205.
- 7- Kennedy A.L., Lyons T.J. Glycation, oxidation and lipoxidation in the development of diabetic complications. *Metabolism*, 1997, 46 (1): 14-21.
- 8- Becker D.J., Reul B., Ozcelikay A.T., Buchet J.P., Henquine J.C., Brichard S.M. Oral selenite improves glucose homeostasis and partly reverses abnormal expression of liver glycolytic and gluconeogenic enzymes in diabetic rats. *Diabetologia*, 1996, 39: 3-11.
- 9- Battell M.L., Delgatty H.L., McNeill J.H. Sodium selenate corrects glucose tolerance and heart function in STZ diabetic rats. *Mol Cell Biol Chem.*, 1998, 179: 27-43.
- 10- Dasilva C., Poupon J., Guillausseau P.J., Chappuis P. In: *Medicine and Biology, Serum selenium, micro and macrovascular complications in diabetic patients, in therapeutic uses of trace elements.* New York: Plenum press, 1996, 167-170.
- 11- Peskin A.V., Koen Y.M., Zbarsky I.B., Konstantinov M. Superoxide dismutase and glutathione peroxidase in tumors. *FEBS Lett.*, 1997, 78: 41-45.
- 12- Schrauzer G.W. Selenium-mechanistic aspects of anticarcinogenic action. *Biol Trace Elem Res.*, 1993, 33: 51-62.
- 13- Awai M., Narasaki M., Yamanoi Y., Seno S. Induction of diabetes in animals by the parenteral administration of ferric nitrilotriacetate. A model of experimental hemochromatosis. *Am. J. Pathol.*, 1979, 95: 663-672.
- 14- Mohandas J., Marshall J.J., Duggin G.G., Horvath J.S., Tiller D.J. Differentiation distribution of

- 
- glutathione and glutathione-related enzymes in rabbit kidney. Possible implication in analgesic nephropathy. *Biochem pharmacol.*, 1984, 33 (11): 1801-1807.
- 15- Wright J.R., Colby H.D., Miles P.R. Cytosolic factors which affect microsomal lipid peroxidation in lung and liver. *Arch Biochem. Biophys.*, 1981, 206 (2): 296-304.
- 16- Zimmermann T., Leonhardt H., Kersting S., Albrecht S., Range U., Eckelt U. Reduction of postoperative lymphedema after oral tumor surgery with sodium selenite. *Biol Trace Elem Res.*, 2005, 106 (3): 193-204.
- 17- Neve J., In: Trace elements in medicine and biology: Biological function of selenium. Walter de Gruyter, Berlin, 1989, 97-111.
- 18- Takahashi M., Sato T., Shinohara F., Echigo S., Rikiishi H. Possible role of glutathione in mitochondrial apoptosis of human oral squamous cell carcinoma caused by inorganic selenium compounds. *Int J Oncol.*, 2005, 27 (2): 489-495.
- 19- Clark L., Combs G., Turnbull B., Slate E., Chalker D., Chow J. Effect of selenium supplementation for cancer prevention in patients with carcinoma of the skin. *JAMA.*, 1996, 276: 1957-1963.
- 20- Eltaveb A.A., Liu Q., Gan L., Liu H., Xu H. Antagonistic effect of scutellarin on the toxicity of selenium in rat livers. *Bio Trace Elem Res.*, 2004, 98: 253-264.

Archive of SID