

()

*
Archive of SID

(NRS)

% /
(VAS)

NRS VAS

()

()

Pre-

()

(emptive analgesia

:

(

(

(

(

()

()

()

()

()

Archive of SID

(CNS

Plasticity)

()

()

SPSS 9.01

/

% mg/kg

% /

/ mg/kg

/

/

% /

(% /)

()

()

()

VAS

VAS

(p= /)

(p= /)

(p= /)

(P< /)

VAS

) NRS () ()

NRS

(

(p= /))

(P< /)


.(

(P< /)

.(P< /)


(P < /) (/ ± /)

(/ ± /)



()

VAS



()

NRS



()

()

()

()

()

Archive of SID

()

()

()

()

(%) (Power)

%

()

Archive of SID

References:

1. Johan J, Boica Post operative pain; the management of pain. Philadelphia: Lea and Febiger 1990, 461-80.
2. Tuffin J, Cun liffe D, Shaw S. Do local analgesics injected at the time of third molar removal under general anesthesia reduce significantly post operative analgesic requirements? A double-blind controlled trial. *Br J Oral Maxillofac Surg* 1989; 27: 27-32.
3. Tversky M, Cozacov C, Ayache M, et al. Postoperative pain after inguinal herniorrhaphy with different types of anesthesia. *Anesth Analg* 1990; 70: 29-35.
4. McQuay H, Carroll D, Moore R. Postoperative orthopedic pain—the effect of opiate premedication and local anesthetic blocks. *Pain* 1988; 33: 291-95.
5. Ke RW, Portera SG, Bagous W, et al. A randomized, double-blind trial of preemptive analgesia in laparoscopy. *Obstet Gynecol* 1998; 92: 972-5.
6. Woolf CJ. Evidence of a central component of post injury pain-hypersensitivity. *Nature* 1983; 306: 686-8.
7. Woolf CJ, Wall PD. Relative effectiveness of primary afferent fibers of different origin in evoking a prolonged facilitation of the flexor reflex in the rat. *J Neurosci* 1986; 6: 1433-42.
8. Willard PT, Blair NP. Is wound infiltration with anesthetic effective a pre-emptive analgesia? A clinical trial in appendectomy patients. *CJS* 1997; 40: 213-7.
9. Ko CY, Thompson JE Jr, Alcantara A, et al. Preemptive analgesia in patients undergoing appendectomy. *Arch Surg* 1997; 132: 874-8.
10. Conno FD, Carcaceni A, Gamba A, et al. Pain measurement in cancer patients: A comparison of six methods. *Pain* 1994; 57: 161-6.
11. Price DD, Bush FM, Long S, et al. Comparison of pain measurement characteristics of mechanical visual analogue and simple numerical rating scale. *Pain* 1994; 56: 217-26.
12. Chapman CR, syjala K. Measurement of pain. In: *The management of pain*. Philadelphia: Lea and Febiger, 1990, 580-94.
13. Jebels J, Reilly JS, Gutierrez J, et al. The effect of pre-incisional infiltration of tonsils with bupivacaine on the pain following tonsillectomy under general anesthesia. *Pain* 1991; 47: 305-8.
14. Buggedo G, Carcamo C, Mertens R, et al. Precutaneous ilioinguinal and iliohypogastric nerve block with 0.5% bupivacaine for post-herniorrhaphy pain management in adults. *Reg Anesth* 1990; 15: 130-3.
15. Woolf C, Chong MS. Preemptive analgesia-treating postoperative pain by preventing the establishment of central sensitization. *Anesth Analg* 1993; 77: 362-79.
16. Heard S, Edwards W, Ferari D. Analgesic effects of intraarticular bupivacaine or morphine after arthroscopic knee surgery: a randomized, prospective, double blind study. *Anesth Analg* 1992; 74: 822-6.
17. McQuay H, Carroll D, Moore R. Postoperative orthopedic pain – the effect of opiate premedication and local anesthetic blocks. *Pain* 1988; 33: 291-5.
18. Woolf C, Wall P. Morphine-sensitive and morphine-insensitive actions of C-fiber input on the rat spinal cord. *Neurosci lett* 1986; 64: 221-5.
19. Ejlersen E, Andersen H, Eliassen K, et al. A comparison between preincisional and postoncisional lidocaine infiltration and postoperative pain. *Anesth Analg* 1992; 74: 495-8.