

()

*

Archive of SID

()

(% /)

(% CI; /)% /

BMI

BMI

// : : - // :

()

(Low back Pain)

%

(.)

%

() %

(.)

()

(.)

(.)

% /

%

%

(.)

%

()

(

)

(.)

)

(

() Nordic

(.)

)

BMI

(...

()

EPI- Info SPSS

χ

(P < /)

()

(P < /)

(/)

(BMI)

(% /)

(% /)

/ (± /) / (± /)

BMI

(/ CI: /) /

(P < /)

/ (± /)

(P < /) / (± /)

<

(SE= /)

(Flexion)

(P= /)

(SE= /)

(Extension) <

/

BMI

(Twisting) <

(Lateral Bending) <

(P= /)

(% /)

(% /)

%

% / % /

%

%

(

P-value	CI	OR	
/	/ /	/	BMI
/ *	/	/	BMI<18
/ *	/ /	/	BMI>27
/ *	/ /	/	>
/	/ /	/	
/ *	/ /	/	kg
/	/ /	/	kg
		<	
/	/ /	/	%
/	/ /	/	%
			()
/ *	/ /	/	
/ *	/ /	/	<
			$\alpha = /$ *

ARC

% / % / % /

(P> /)

...

kg

()%

()%

()% /

OR= /

OR= /

Multi factorial

/

()

/

.()

()

.()

/

.()

(% /)

(% /)

()

.()

.()

/

()

OR= /

.()

/

.()

.()

BMI

()

% /

%

.()

.()

OR= /

()

.()

OR= /

<

. ()

()

(%)

(% /)

()

. ()

)

. () (/ /

()

. ()

. ()

. ()

ICU

% /

% /

. ()

. ()

. ()

. ()

Archive of SID

References:

1. Edlich R, Woodard CR, Haines, MJ. Disabling back injuries in nursing personnel. *J Emerg Nurs* 2001;27:150-5.
2. Knibbe JJ, Friele RD. Prevalence of back pain and characteristics of the physical workload of community nurses. *Ergonomics* 1996;39:186-98.
3. Smedley J, Egger P, Cooper C, *et al.* Manual handling activities and risks of low back pain in nurses. *Occup Environ Med* 1995;52:160-63.
4. diagnosed disorders and perception of pain. *Applied Ergonomics* 1999;30: 325-30.
5. Ando S, Yuichirono B, Shimaoka M, *et al.* Associations of psychosocial and individual factors with three different categories of back disorder among nursing staff. *J Occup Health* 2004;46:100-8.
6. Burdorf A, Scorock G. Positive and negative evidence of risk factors for back disorders. *Scand J Work Environ Health* 1997;234:243-56.
7. Levy BS, Wegman DH. Occupational Health: recognizing and preventing work related disease and injury, selected groups of workers, 4th ed. Lippincott Williams & Wilkins, 2000,770-771.
8. Bernard B. Musculoskeletal Disorders and Workplace Factors: A Critical Review of Epidemiologic Evidence for Work-Related Musculoskeletal Disorders of the Neck, Upper Extremity, and Low Back. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, Cincinnati, OH. 1997.
9. Kuorinka I, Jonsson B, Kilbom A, *et al.* Standardised Nordic questionnaires for the analysis of musculoskeletal symptoms. *Applied Ergonomics* 1987; 18: 233-7.
10. Kaergaard A, Andersen JH, Rasmussen K, Mikkelsen S. Identification of neck-shoulder disorders in a 1 year follow-up study. Validation of a questionnaire-based method. *Pain* 2000; 86: 305-10.
11. Bjrksten MG, Boquist B, Talbkck M, The validity of reported musculoskeletal problems. A study of questionnaire answers in relation to
12. Prevalence of low back pain among staff in a rural hospital in Nigeria. *Occupational Med* 2000; 50:107-10.
13. Rossi A, Marino G, Barbieri L, *et al.* Backache from exertion in health personnel of the Istituti Ortopedici Rizzoli in Bologna. a case-control study of the injury phenomenon in the 10-year period of 1987-1996. *Epidemiol Prev* 1999; 23:98-104.
14. leighton DJ, Reilly T. Epidemiological aspects of back pain the incidence and prevalence of back pain in nurses compared to the general population. *Occup Med* 1995; 45:263-7.
15. De Zwaart BCH, Frings-Dresen MHW, Van Dijk Fjh. Physical workload and the ageing workert a review of the literature. *Int Arch occup Environ Health* 1995; 68:1-12.
16. Venning PJ, Walter SD, Stitt LW. Personal and job-related factors as determinants of incidence of **back** injuries among nursing personnel. *J Occup Med* 1987; 29:820-5.
17. Lagerstr M, Wenemark M, Hagberg M, *et al.* The Moses Study Group. Occupational and individual factors related to musculoskeletal symptoms in five body regions among Swedish nursing personnel. *Inter Arch Occupat Environmen Health* 1995; 68: 27-35.
18. Stubbs DA, Buckle PW, Hudson MP, *et al.* Back pain in the nursing profession. epidemiology and pilot methodology. *Ergonomics* 1983;26:755-65.
19. Eriksen W. The prevalence of musculoskeletal pain in Norwegian nurses' aides. *Int Arch Occup Environ Health* 2003; 76:625-30.
20. Leboeuf-Yde C, Kyvik KO, Bruun NH. Low back pain and lifestyle. Part 1: Smoking. Information from a population-based sample of 29,424 twins. *Spine* 1998; 23:2207-13.
21. An HS, Silveri CP, Simpson JM, *et al.* Comparison of smoking habits between patients with surgically confirmed herniated lumbar and

cervical disc disease and controls. *J Spinal Disord* 1994; 7:369-73.

23. American Association of Neurological Surgeons. Sprain and strain. 17 April, 2002.

24. Mandel JH, Lohman W. Low back pain in nurses the relative importance of medical history, work factors, exercise and demographics. *Res Nurs Health* 1987; 10:165-70.

25. Estyn-behar M, Kaminski M, Peigne E, et al. Strenuous working conditions and musculo-skeletal disorders among female hospital workers. *Int Arch Occup Environ Health* 1990; 62:47-57.

26. Byrns G, Reeder G, Jin G, Pachis K, Risk factors for work-related low back pain in registered nurses, and potential obstacles in using

mechanical lifting devices. *J Occup Environ Hyg* 2004; 1:11-21.

27. Retsas, A, Pinikahana J. Manual handling practices and injuries among ICU nurses, *Aust J Adv Nurs* 1999; 17: 37-42.

28. Smith DR, Ohmura K, Yamagata Z. Musculoskeletal disorders among female nurses in a rural Japanese hospital. *Minai J Nurs Health Sci* 2003; 5:185-8.

29. Guo H-R. Working hours spent on repeated activities and prevalence of back pain. *Occup Environ Med* 2002; 59:680-8.

30. Wickstrom GJ, Pentti J. Occupational factors affecting sick leave attributed to low-back pain. *Scand Work Environ Health* 1998; 24:145-8.

Archive of SID