



() —

()

%

%

%

()

()

)

(


I

()

II


()

()



%

% % % %






(%)

% % %

()

() %





()

(%)

(%)

(%)

%

()

%

()

References:

- 1- Brooks D.M.: Psoas Compartment block. CRNA, 2000 May; 11(2): 62-5.
- 2- Jankowski C.I., Hebl J.R., Co. workers: A Comparison of psoas compartment block and spinal and general anesthesia for outpatient knee arthroscopy. Anesth Analg, 2004 May; 98(5): 1501.
- 3- Turker G., Uckunkaya N., Co. workers: Comparison of the Catheter-technique Psoas Compartment block and the epidural block for analgesia in partial hip replacement surgery. Acta Anesthesia Scand. 2003 Jan; 47(1): 30-6.
- 4- Capdevila X., Co. workers: Continuous Psoas Compartment block for postoperative analgesia after total hip arthroplasty. Anesth Analg. 2002 Jun; 94(6): 1606-13.
- 5- Horlocker T.T., Hebl J.R., Co. workers: Opioid-free analgesia following total knee arthroplasty-a multi modal approach using continuous lumbar plexus (Psoas Compartment) block, acetaminophen and ketorolac. Reg Anesth Pain Med. 2002 Jan-Feb; 27(1): 105-8.
- 6- Miller R.D.: Anesthesia, USA, Churchill Livingstone. Fifth edition, volume1, 2000; 1530-33.
- 7- Hogan Q. Abrams; Neural blockade for diagnosis and prognosis, A review, Anesthesiology, 1997; 86-216.
- 8- Sansone V., Depoint A., Fanelli G., Agostini M.: Combined sciatic and femoral nerve block for knee arthroscopy: 4 years experience, Arch Orthop Trauma Surg, 1999; 119(3-4): 163-167.
- 9- Goranson B.D., Lang S., Cassidy J.D., Dust W.N., Mckerrell J.: A comparison of three regional anesthesia techniques for out patient knee arthroscopy. Can J Anesth, 1997, 44(4): 371-376.
- 10- Bonicalzi V., Gallino M.: Comparison of two regional anesthetic techniques for Knee arthroscopy. Arthroscopy, 1995: 11(2): 207-212.