



() CAPD
Mass
(P> /)
CAPD
(P< /)
CAPD
CAPD

CAPD
()
% / CAPD
(P> /)
(P< /)
CAPD

:

CAPD: Continuous ambulatory peritoneal dialysis

碑

(.)

()

()

(%) (%)

(%)

/

%

% / CAPD

% / CAPD

% / CAPD

% / CAPD

CAPD

() (p> /)

SGA

%

%

Subjective Global Assessment
Fat pad
pinching

--

%	%	%	CAPD
% /	%	% /	

< %	% -	> %	
%	%	%	CAPD
% /	% /	% /	

P> /

% CAPD

%
% /

(p> /)

% CAPD

%
% /

pvalue

(p> /)

CAPD

%	%	%	CAPD
% /	% /	% P> /	

%	%	%	CAPD
% /	% /	%	

P> /

% CAPD

()
()
) % /

% CAPD

%
% /
%
) (p> /).
Pvalue
()

(p> /) CAPD
) P> /
(p> /) (

CAPD

()

(p < /)

()

() IGF-1

() CAPD

(RRF)

() CAPD CAPD ()

/ / CAPD (PC < /)

()

()

CAPD ()

% / %

CAPD ()

(P > /)

(P > /) BIA ()

CAPD () DEXA

(P < /) CAPD ()

CAPD () %

() ()

References:

- 1- Ikizler T. Alp. Hakim R.: Nutrition in end stage renal disease. *Kidney int*, 1996; 50: N: 2, 343-353.
- 2- Rocco Micheal V., B Lumen krantz M.J.: Nutrition, Hand Book of dialysis. 3th ed: Lippincote william and wilkins, philadelphia, 2001; 420-445.
- 3- Baker J.P., Detzky A.S., Wesson O.E., et al: Nutrition assessment, A comparision of clinical judjment and objective measurement. *N Engl J medicin*, 1982; 306: 969-992.
- 4- Enia G., Sicsysco C., Alati G., eT al: Subjective global assessment of nutrition in dialysis patients. *Nephrology, Dialysis, trnasplant (NDT)*, 1993; N: 8 1094-1098.
- 5- Lowrie E.G., Lew N.L.: Death risks in HD paitents. The predictive value commonly measured variable and an evaluation of death rate differences between facilities. *Am j kidney disease*, 1990; 15: 458-482.
- 6- Young G.A., Kopple J.D., Lindholm B., et al: Nutritional assessment of CAPD patients An international study. *Am j Kidney disease*, 1991; 17: 462-471.
- 7- Kopple J.D., Effect of nutrition on morbidity and mortality in maintenance dialysis patients *Am j kidney disease*, 1994; 24: 1002-1009.
- 8- Herrmann F.R., Safranc, Levkoff S.E., et al: Serum albumin level on admission as a predictor of death, length of stay and readmission, *Arch Inter med*, 1992; 152: 125-130.
- 9- Lowrie E.G., Huang W.H., Lew N.L.: Death risk predictors among P.D. and H.D. patients, a preliminary comparison, *Am j Kidney disease*, 1995; 26: 220-228.
- 10- Uram M.M., Mitmann N., bonomini Zet., al: Marker for survival in dialysis. A 7 years prospective study (Analysis of United State Renal Data Sestem = USRDS). *Am j Kidney disease*, 1995; 26: 209-219.
- 11- Kople J.D., Berg R., Houser H., et al: Nutrition status of patients with different level of CRF. *Kidney int*, 1989; 36: 184-194.
- 12- Walser M.: Does prolonged protein restriction preceeding dialysis lead to protein malnutrition as the onset of dialysis? *Kidney Int*, 1993; 44: 1139-1144.
- 13- Krieg. J.r, Santos F.: Chan JCM, Growth Hormon, and Insulin like growth factor and kidney. *Kidney Int*, 1995; 48: 321-336.
- 14- Schulman G., Wingard R.L., Hutchinson R.L., et al: Effect of recombinant human GH in intradialytic parenteral nutrition in malnourished hemodialysis patients, *Am j kidney disease*, 1993; 21: 527-534.
- 15- Parker T.F., Wingard R.L., Husni L., et al: Effect of the membrane biocompatibility on nutrition parameters in chronic hemodialysis patients. *Kidney int*, 1996; 49: 551.
- 16- Hakim R.M., Levin N.: Malnutrition in HD patients. *Am j kidney disease*, 1993; 21: 125-137.
- 17- Jones M.R.: Etiology of sever malnutrition: results of an international cross-sectional study in continous peritoneal dialysis patients. *Am j kidney disease*, 1994; 23: 412-420.
- 18- Teehan B.P., Schlefier C.R., Brownj.: Adequacy of continous peritoneal dialysis morbidity and

mortality in chronic peritoneal dialysis .
Am j kidney disease, 1994; 25: 990-1001.

19- Kopple, J.D., Hirschberg R.: Nutrition and peritoneal dialysis, Nutrition and kidney edited by mitch WE, Klahr S, little Brown company, 1993; 290-299.

20- Lidnholm B., Bergstormj.: Nutrition management of patients undergoing peritoneal dialysis, edited by Nolh KD, Dordrecht, Kleuwer, Academic publisher, 1989; 230.

Archive of SID