




() -



%

%

%

()

()

()

(%)

C_C

()	()	
/	/	

(%)

(%)

(%)

(%)

(% /)

(%)

(%)

C - C

(%)

(%)

(% /)

(%)

%

(%)

(%)

(%)

)

(%)

:(

(%)

C - C

(%)

(% /)

C - C

(%)

(% /)

C - C

(% /)

(%)

C - C

(%)

(%)

%

(%)

)

(%)

(%)

:(

(%)

(%)

(%)

(% /)

(%)

BPH¹

(% /)

(% /)

(%)

(%)

(%)

(%)

(%)

C - C

(%)

(% -)


C - C

(%)

-C

(% /)

¹ Benign prostatic hypertrophy



% /

C

. ()

% /

C-C

(%)

. ()

/

/

. ()

% /

%

%

%

%

. ()

()		
		C-C
		C-C
/	/	C-C
		C-C

. ()

%

()

-)

(

(%)

:

(% /)

(%)

(% /)

(% /)

(% /)

%

(%)

(%)

%

% /) (% /)

(

(%)

C C

% / % /

² Sampath

³ Young

⁴ Lansford

/

) C C C C
(% - - /

.()

C -C C -C

.()(% /)

% C -C C -C C -C

.() C -C C -C

% /

% /

% /

. %

%

Archive of SID

⁵ Clark

References:

- 1- Victor Maurice., Ropper Allen H.: Principles of Neurology. Seventh Edition, Macgraw – Hill ompany, 2001; 1322.
- 2- Clark charles R.,: Cervical Spine Third Edition, Lippincott – Raven, 1998; 735-803.
- 3- Youmans Julian R.: Neurological surgery. Fourth Edition, Saunders, 1996; 2241-2271.
- 4- Rowland Lewis P-merrit,s: Neurology. Tenth Edition, Lipponcot Williams and Wilkins, 2000; 430-433.
- 5- Schmideck Henry H.: Operative Neurosurgical Techniques. Fourth Edition, Saunders, 2000; 1988-1990.
- 6- Young W.: Cervical Spondylotic Myelopathy, A Common Cause of Spinal Cord Dysfunction in Older Persons. Am Fam Physician, 2000 sep1; 62(5): 1064-70.
- 7- Mihara H., and et al: Cervical Myelopathy Csused by C3-C4 Spondylosis in Elderly Patients: A Radiographic Analysis of Pathogenesis. Spine, 2000 Apr 1; 25(1): 796-800.
- 8- YUA W.M. and et al: The Torg Parlov Ratio in Cervical Spondylotic Myelopathy. A comparative Study between Patients with Cervical Spondylotic Myelopathy and a non Spondylotic Myelopathy and A Non Spondilotic, Non Myelopathic Population, spine, 2001 Aug 15; 26(16):1760-64.
- 9- Misfud V., Pullicino P.: Spinal Cord MRI Hyperintensities in Cervical Spondylosis. An Ischemic Pathogenesis? J.Neuroimaging, 2000 Apr; 10(2): 96-100.
- 10- Matsumoto M., and et al: Relationships between Outcomes of Concervative Treatment and Magnetic MRI Findings in Patients with Mild cervical Myelopathy Caused by Soft Disc Herniation. Spine, 2001 Jul 15; 26(14): 1592-8.
- 11- Matsumoto M., et al: Increased Signal Intensity of the Spinal Cord on MRI in Cervical Compressive. Spine, 2002 Mar 15; 25(6): 677-82.
- 12- Morio et al: Correlation between Operative Outcomes of Cervical Compression Myelopathy and MRI of the Spinal Cord. Spine, 2001 jun 1; 26(11): 1238-45.
- 13- Yonanobu K.: Cervical Radiculopathy and Myelopathy when and what Can Surgery Contribute to Treatment. Eur spine J, 2000 Feb; 9(1): 1-7.
- 14- Sampath p., et al: Outcome of Patients Treated for Cervical myelopathy. A Prospective Multicenter Study with Independent Clinical Review. Spine, 2000 Mar 15; 25(6): 670-6.
- 15- Lunsford L.D., Bissonette D.J., Zorub D.S.: Anterior Surgery for Cervical Disc disease Part 2, Treatment of Cervical Spondylatic Myelopathy in 32 Cases. J. Neurosurgery, 1980; 53:12-19.
- 16- Shimizu T.: Scapulohumeral Reflex Its Clinical Significace and Testing Maneuvers Spine, 1993; 2182-2190.

Archive of SID