



.( )

( )

.( )

LDL/HDL APOB<sub>1</sub> APOA<sub>1</sub> LDL HDL  
FBs-APOA1/ APOB<sub>1</sub>

.( )

t

.( )

%

%

.(p= / )



-

$\frac{1}{2} + \frac{1}{2}$       %      ( )  
 $\frac{1}{2} + \frac{1}{2}$       %      ( )  
 = / )      .(p= / )  
 .( ) (p

HDL

DLC

FBS

FBS

.(p= / )

\*\*\*\*\*

**References:**

1- Brown N., and et al.: Quality of life four years after acute myocardial infraction: Short from 36 scores compared with a normal population.Heart, 1999; 31: 352-358.  
 2- Bennet S.J., And et al.:Comparison of Quality of life measures in heart failure. Nursing research, 2003; 52 (4): 207-216.

5- Betsy A., Finkelmeier.: Cardiothoracic surgical nursing, 2000 (2).  
 6- Iowa CHAMPS: Cardiac Rehabilitation Guide. Risk Factors for coronary artery disease.  
 7- American Family physician: Using Apotipoprotein B level to assess cardiac Risk.  
 8- Farmer J. A., Gotto AMJR.: Dyslipidemia and other risk factors for coronary artery diseases. In Braounwalde (Ed). Heart Disease: A textbook of cardiovascular medicine ed.5: Philadelphia, WB Saunders, 1997.