




|| : || :



AM

_____ ()

¹ FLACC

² Wong & Baker



()

)
(

Archive of SID

O₂ N₂O

mL/kg / /

% /

(ml/kg /)

ml/kg

(%)

FLACC

()

« »

mg/kg)

(

/ ± /

(/)

FLACC

()

()

()

rs= /

Archive of SID

References:

- 1- Patrick H, Mckenna, Fernard. A Fever, Prenatal and postnatal urologic emergencies. In: A Barry Belman, Lowell R King, Stephen A Kramer. Clinical pediatric urology. 4th ed; 2002: 185.
- 2- Merkel SI, Voepel-Lewis T, Shagevitz JR, et al. **Pediatric Nurs** 1997 May-Jun; 23(3):293-7
- 3- Bruce A. Bevn stein and Lee M. Pachter, cultural consideration in children pain. **IN: Neil.L SCHECHTER, chartes B. Berde, Myron yaster, pain in infants, children and Adolescents. 2nd ed; 2003:151.**
- 4- Dalens B, Vanneuville G, Dechelotte P. Penile block via the subpubic space in 100 children. **Anesth Analg** 1989; 69:41.
- 5- Navil F. Sethna, Charles B Berde. Pediatric regional anesthesia. **In: George A. Gregory, Pediatric Anesthesia.4th ed; 2002.p.302.**
- 6- Soliman MG Tremblay NA. Nerve block of the penis for postoperative pain relief in children. **Anesth Analg** 1978; 57:475-8.
- 7- Bacon AK.An alternative block for post-circumcision analgesia. **Anesth Intensive care** 1977; 5:63-4.
- 8- Bernard J.Dalens. Regional anesthesia in children. **In: Ronald Miller. Anesthesia. 5th ed; 2000:1573-74.**

Archive of SID