



E

// : // :



()E

E E

E TIBC

SPSS

(p< / . / ± / ± / ± / : / ± /)

E

E

E- Mail: ghanefsh@yahoo.com



E
C
E

: CRF

.()E .() /

ESRD

<
%
%

u/n

C E

CRF

() NAD

NADPH

.()

.()

E

()

³ End-stage Renal Disease

¹ Chronic Renal Failure
² Nicotinamid Adenin Dinucleotide

/ ± /

/ ± /

TIBC

(Hb)

TIBC

(p< /)

(HCT)

HCT Hb

-

()

±

/					
/	/		-		
/					
/					
/	/				

E

Archive of SID

p< /

(% /)

(% /)

(% /)

± /

± /

(% /)

⁴ Total Iron binding Capacity



B12
ROM

	(n=7)	(n=8)
/	/ ± /	/ ± /
/	/ ± /	/ ± /
/	/ ± /	/ ± /
/	/ ± /	/ ± /
/	/ ± /	/ ± /
/	/ ± /	/ ± /
/	/ ± /	/ ± /
/	/ ± /	/ ± /

()

(HCT₃-HCT₀)

/ ± / / ± /

()

(T= / p= /)

(Hb₃- Hb₀)

/ ± / / ± /

()

t- test

(T= / p= /)

()

RBC

(CRF)

() E

GFR

E

()

()

⁶ Reactive Oxygen Metabolites

⁵ Glumerolar Filtration Rate

()

(p< /)

E

%

()

E

()

E

E

()

E

()

E

E

E

)

() (

E

E

E

E

E

()

() (p< /)

E

⁷ Nemeth

⁸ Cristol

(

)

References:

- 1- Avner E, Harmon WE, Niaudet P. Conservative management of chronic renal Insufficiency. Fine RN, whyte DA, Boydston I. Pediatric Nephrology: Lippincott. Williams & Wilkins; 2004: 1291-1311.
- 2- Rigatto C, Singal PK. Oxidative stress in uremia: Impact on cardiac stress in dialysis patients: **seminars in dialysis** 1999; 12:91-93.
- 3- Giardini O, Gallacci M. Effect of alpha-Tocopherol administration on RBC membrane lipid peroxidation in hemodialysis patient. **Clin Nephrol** 1984; 21:174-180.
- 4- Carrel R, Winter Bourn C, Rachmile WF. Activated oxygen and hemodialysis. **Br.J.Horemnet** 1976; 20:259-265.
- 5- Suhayakin A, Yurtkaruon M. The effect of vitamin E therapy on plasma and erythrocyte lipid peroxidation in chronic hemodialysis patient. **Clinical chimica Acta** 1984; 185: 109-120.
- 6- Dinarello C.A. The biology of interleukin. I and its relevance to hemodialysis. **Blood purif** 1983; 11:197-204.
- 7- Canoud B, Cristol M. Imbalance of oxidative and antioxidants in hemodialysis patient. **Blood purification** 1997; 17:99-105.
- 8- Tatsuya N, Yoshiaki T, Kenji T. The effect of vitamin E-bonded dialyzer membrane on red blood cell survival in hemodialyzed patients. **Artificial organs** 2003; 27(3):214-217.
- 9- Usberti M, Bufano G, Lima G, Gazzotti RM, Tira P, Geradi G, Di Lorenzo D. Increased red blood cell survival reduces the need of erythropoetin in hemodialyzed patients treated with exogenous glutathione and vitamin E-modified membrane. **Contrib Nephrol** 1999; 127:208-214.
- 10- Jiao Y, Okumiya T, Saibara T, Tsubosaki E, Mstsumura H, Park K, Sugimoto K, Kageoka T, Sasaki M. An enzymatic assay for erythrocyte life time. **Clin Biochem** 1998; 31:59-65.
- 11- Luciak M. Antioxidants in the treatment of patients with renal Failure. **Rocz Akad Med Bialymst** 2004; 49:157-161.

- 12- Eiselt J, Racek J, Trefil L, opatrný K. Effect of a vitamin E-modified dialysis membrane and vitamin C infusion on oxidative stress in hemodialysis patients. **Artif organs** 2001; 25: 430- 436.
- 13- Turi S, Nemeth I, Varga I, Matkovics B. Erythropoietin and oxidative stress in hemodialysis: beneficial effects of vitamin E supplementation. **Nephrol Dial Transplant** 1999; 14: 252-253.
- 14- Smith KS, Lee CL, Ridlington JW, Leonard SW, Devaraj S, Traber MG. Vitamin E supplementation increases circulating vitamin E metabolites tenfold in end stage renal disease patients. **Lipids** 2003; 38:813-819.
- 15- Behrman R, Kliegman R, Jenson H. Toxic Nephropathies – Renal Failure. Beth A, Vogt and Ellis T, Avner. Nelson textbook of pediatrics: Saunders; 2004:1766-1782.
- 16- National kidney foundation, kidney disease outcomes quality initiative. Clinical practice guidelines for anemia of chronic kidney disease, 2000:21.
- 17- Eschbach JW. The anemia of chronic renal failure: pathophysiology and the effect of recombinant erythropoietin. **Kidney Int** 1989; 35:134-148.
- 18- Parker PA, Izard MW, Maher JF. Therapy of iron deficiency anemia in patients on maintenance dialysis. **Nephron** 1979; 23:181-186.
- 19- Hocken AG. Hemodialysis in chronic renal failure. **Nephron** 1982; 32:28-35.
- 20- Longnecker RE, Goffinet JA, Hendler ED. Blood loss during maintenance hemodialysis. **Trans Am Soc Artif Intern Organs** 1974; 20:135-138.
- 21- Tacob HJ, Eaton JW, Yawata NY. Shortened blood cell survival in uremic patients, Beneficial and deleterious effects of dialysis. **Kidney Int** 1975; 7:139-143.
- 22- Himmelfarb J, Lazarus JM, Hakim R. Reactive oxygen species production by monocyte and polymorphonuclear leukocytes during dialysis. **Am J Kidney Dis** 1991; 17:271-276.
- 23- Rosenkranz AR, Templ E, Traundl O, Hinzl H, Ziabinger GJ. Reactive oxygen product formation by human neutrophils as an early marker for biocompatibility of dialysis membranes. **Clin Exp Immunol** 1994; 98:300-305.
- 24- Usberit M, Lima G, Arisi M, Bufano G, Avanzo L, Gazzotti RM. Effect of exogenous reduced glutathione on the survival of red blood cells in hemodialyzed patients. **J Nephrol** 1997; 10: 261-265.
- 25- Galli F, Canestrari F, Buoncristiani U. Biological effect of oxidative stress in hemodialysis. **blood purif** 1999; 17; 79-94.
- 26- Galli F, Canestrari F, Bellomo G. Pathophysiology of the oxidative stress and its implication in uremia and dialysis. **Contrib Nephrol** 1999; 127:1-31.
- 27- Galli F, Rovidati S, Chiarantini L, Campus G, Canestrari F, Buoncristiani U. Bioreactivity and biocompatibility of a vitamin E-modified multilayer hemodialysis filter. **Kidney Int** 1998; 54:580-589.
- 28- Bnnefont RD, Lehmann E, Jaudon MD, Delatre J, Perrone B, Rechke JP. Blood oxidative stress and lipoprotein oxidizability in hemodialysis patients: effect of the use of a vitamin E-coated dialysis membrane. **Nephrol Dial Transplant** 2000; 15:2020-2028.
- 29- Usberti M, Bufano G, Lima G, Gazzotti RM, Tira P, Geradi GM, Di Lorenzo D. Increased red blood cell survival reduces the need of erythropoietin in hemodialyzed patients treated with exogenous glutathione and vitamin E modified membrane. **Contrib Nephrol** 1999; 127:208-214.

30- Usberti M, Geradi G, Micheli A, Tira P. Effects of a vitamin E-bonded membrane and of glutathione on anemia and erythropoietin requirements in hemodialysis patients. **JN** 2002; 15:558-564.

31- Nemeth I, Turi S, Haszon I, Berczki C. Vitamin E alleviates the oxidative stress of erythropoietin in uremic children on hemodialysis. **Pediatr Nephrol** 2000; 14:13-17.

32- Cristol JP, Bosc JY, Badiou-Leblanc M, Lorrho R, Descomps B, Canaud B. Erythropoietin and oxidative stress in hemodialysis: beneficial effects of vitamin E supplementation. **Nephrol Dial Transplant** 1997; 12:2312-2317.

Archive of SID