



MD

MD

MD

\* MD

// : // :

( )

( )  
(Bil> 8mg/dl)

Rh ABO

G6PD

/ gr/kg

IVIG

IVIG

IVIG

( )

\*

Email: Hasanboskabadi@yahoo.com



Rh

(.)

Rh

mg

(.)

O

B A

% /

(.)

( )

IVIG

(.)

IVIG

IVIG

(.)

G6PD

p

/

( )

NICU

Rh ABO

<sup>1</sup> Intravenous immunoglobulin



( )

P value			
	=	=	
/	/ ± /	/ ± /	/
/	/ ± /	±	( )
/	/ ± /	/ ± /	( )
/	/ ± /	/ ± /	( )
/	(% /)	(% /)	ABO
/	(% /)	(% /)	Rh

( )

Rh

( )

ABO

( )

P value			
	=	=	
/	/ ±	/ ± /	( )
/	/ ± /	±	( )
/	(% /)	(% /)	
/	/ ± /	/ ± /	
/		(% /)	



FC

IVIG  
ABO Rh

IVIG

IVIG  
FC

Rh

IVIG

( ) % % / ( )

ABO Rh

IVIG

IVIG

IVIG

( )  
(

Rh  
( )

IVIG Rh E

ABO  
IVIG

IVIG  
ABO,RH

IVIG ( )

( )

D FC  
FC

<sup>2</sup> Rubo  
<sup>3</sup> Arad,Ergas



## References:

- 1- Zipursky A, Bowman JM. Isoimmune hemolytic disease, In: Nathan OG, Oski FA (eds). **Hematology of infancy and childhood**, 4<sup>th</sup> ed. WB Saunders C, 1993; 1: 66-69.
- 2- Koenig JM. Evaluation and treatment of erythroblastosis fetalis in the neonate. In: Christensen RD. **Hematologic problems of the neonate**. 1<sup>th</sup> ed. Philadelphia; W.B Saunders 2000: 185-203.
- 3- Alpay F Sarici SU, Okutan V, Erdem G, Ozcan, Goky E: High dose intravenous immunoglobulin therapy in neonatal immune hemolytic jaundice. **Acta-Paediatr** 1999; 88: 216-9.
- 4- Jackson JC. Adverse events associated with exchange transfusion in healthy and ill newborns, *Pediatrics* 1997; 7 (99).
- 5- Dennery PA. Pharmacological interventions for the treatment of neonatal jaundice. **Semin-Neonatal** 2002; 7: 111-119.
- 6- Lakatos L, Csathy L, Nemes E. Bloodless treatment of a Jehovah's Witness infant with ABO Hemolytic Disease. **J-Perinatol**, 1999; 19(7): 530-32.
- 7- Gottstein RS, Cooke RW. Systemic review of intravenous immune globulin in hemolytic disease of the newborn. *Arch Dis Child, Fetal Neonatal Ed* 2003; 88: F8-F10.
- 8- Pishva N, Madani A, Homayoon k. Prophylactic intravenous immunoglobulin in neonatal immune hemolytic. **Iranian journal of Medical Sciences** 2000 Dec; 25, (3&4): 125-130.
- 9- Tanyer G, Siklaer Z, Dallar Y, et al: Multiple dose IVIG treatment in neonatal immune hemolytic jaundice. **J-Tropical pediatr** 2001; 47:50-53.
- 10- Porter TE, Silver RM, Jackson GM, Branch D, Scott Jr. Intravenous immune globulin in the management of severe RHD hemolytic disease. **Obstet-Gynecol-Surv** 1997; 52(3): 193-197.
- 11- Ergaz Z, Arad I: Intravenous immunoglobulin therapy in neonatal immune hemolytic jaundice. **J-Perinatol-Med** 1993; 21(3): 183-187.
- 12- Chitkara U, Bussel J, Alvarer M, et al: High dose IVIG: Dose it has a role in the treatment of severe erythroblastosis fetalis. **Obstet-Gynecol** 1990; 76: 703-708.
- 13- Hara T, Mizuno Y, Kawano M, et al: Treatment of immune hemolytic anemia with gammaglobulin. **J- pediatr** 1987; 110: 817-818.
- 14- Sato K, Hara t, kondo T et al: High dose intravenous gammaglobulin therapy for neonatal immune hemolytic jaundice due to blood group incompatibility. **Acta-Paediatr-Scand** 1991; 80(2): 163-166.
- 15- Kabo S, Ariga T, Tsuneta H: Can high dose immunoglobulin therapy be indicated in neonatal Rhesus hemolysis. A successful case of hemolytic disease due to Rhesus (C+E) incompatibility, **J- pediatr** 1991; 150: 507-8.
- 16- Rubo J, Albrecht K, Iasch P, et al: High dose intravenous immune globulin therapy for hyperbilirubinemia caused by Rh hemolytic disease. **J-pediatr** 1992; 121: 93-97.