



MD

MD

MD

\*

/ / :

/ / :

.( )

( )

.( )

( )

\*

ICU

BP

/

SGOT

ALP SGPT

mg/dl

mg/dl /

mg/dl /

/ /

( )

ICU

/

ICU

Archive of SID

( - ) %

( )

CPR

ET CO<sub>2</sub>

CPR

CPR

ET ( )

CPR CO<sub>2</sub>

<sup>1</sup> End -tidal Co<sub>2</sub>

( )

\*\*\*\*\*

### **References:**

- 1- Barret - Boyes K. **Cardiac Surgery.** 3<sup>rd</sup> ed. Churchill Livingstone. 2003; 2: 1799-1820.
- 2- Barret - Boyes K. **Cardiac surgery.** 3<sup>rd</sup> ed. Churchill Livingstone. 2003; 2: 1820-1821.
- 3- Edmunds LH. **Cardiac Surgery in the adult.** 2<sup>nd</sup> ed. Mc Graw-hill. 2003; 1: 439-469.
- 4- Vander Tj. Two techniques for the control of cardiac bleeding. **Ann.Thorac Surg** 1994; 57:762.
- 5- Fairman RM, Edmunds LH. Emergency thoacotomy in the surgical intensive care unit after open cardiac operation. **Ann Thorac Surg** 1981; 32:386.
- 6- Anthi A, Tzelepis GE, Alivizatos P, et al. Unexpected cardiac arrest after cardiac surgery:incidence, predisposing causes, and outcome of open chest cardiopulmonary resuscitation. **Chest** 1998; 113:15.
- 7- Feng WC, Bert AA, Browning RA, Sing AK: Open cardiac massage and periresusitative cardiopulmonary bypass for cardiac arrest following cardiac surgery. **J Cardiaovasc surg (torino)** 1995; 36:319.
- 8- Hill RC. Open – chest resuscitation and postcardiac surgery arrest . Chest 1998; 113:3.
- 9- Pottle A, Bullock I, Thomas J, Scott L. Survival to discharge following - open chest cardiac compression (OCCC): a 4 - year retrospective audit in a cardiothoracic specialist center-Royal Brompton and Harefield NHS Trust United Kingdom. **Resuscitation** 2002; 52: 269.
- 10- Wahba A, Gotz W, Birnbaum DE: Outcome of cardiopulmonary resuscitation following open heart surgery. **Scand Cardiovasc j** 1997;31:147.
- 11- Sanders AB, Ogle M, Ewy GA: Coronary perfusion pressure during cardiopulmonary resuscitation.**Am J Emerg Med** 1985; 3:11.
- 12- Weale FE, Rothwell-Jackson RL. The efficiency of cardiac massage. **Lancet** 1962; 1: 990.
- 13- Coletti RH, Hartien B, Gozdejewski S, et al. Origin of canine femoral pulses during standard CPR [abstract]. **Crit Care Med** 1983; 11: 218.
- 14 - Kern KB, Sanders AB ,Ewy GA. Open – chest cardiac massage after closed-chest compression in a canine model: when intervene?. **Resuscitation** 1987; 15:51.
- 15- Falk JL, Rackow EC, Weil MH. End tidal carbon dioxide concentration during cardiopulmonary resuscitation. **N Engl J Med** 1988; 318: 607.
- 16- Weil MH, Bisera J, Terevino RP. cardiac output and end tidal carbon dioxide. **Crit Care Med** 1985; 13: 907.
- 17- Garnett AR, Ornato JP, Gonzalez ER, et al. End tidal carbon dioxide measurement during cardiopulmonary resuscitation, **JAMA** 1987; 257: 512.
- 18- Ornato JP. Hemodynamic monitoring during CPR. **Ann Emerg Med** 1993; 22: 289.
- 19- Sanders AB, Kern KB, Otto CW et al. End – tidal carbon dioxide monitoring during cardiopulmonary resuscitation: A prognostic indicator for survival. **JAMA** 1989; 262: 1347.
- 20- Sanders AB, Ewy GA, Bragg S, et al: Expired pco<sub>2</sub> as a prognostic indicator of successful resuscitation from cardiac arrest. **Ann Emerg Med** 1985; 14: 948.