

:

:

BCG

(% /) (% /)

:

(% /) (% /)

/

/ PPD

(% /)

(% /)

% /

:

(% /) (% /) Cavitary (% /)

% /

% /

()

:

:

()

%

%

%

()

()

% Malawi ()

% /

.()

AFB

.()

M.T.B

.()

%

(.

(

(

(

%

%

(

.()

%

(/ %)

(/ %)

(/ %)

(/ %)

/ %

/ % BCG

/ %

/ % PPD

≤

(/ %)

(/ %)

(/ %)

BCG

PPD

(Pvalue 0.59)

% /

:

% /

% /

/ %

()

/ %

/ %

%

Malawi

% /

()

()

% /

()

References:

- 1- Corbett E L, Watt C y, Walker N, et al. The growing burden of tuberculosis: global trends and interactions with the HIV epidemic. *Arch Intern Med* 2003; 163:1009-1021.
- 2- Reichler M R, Reves R, Bur S, et al. Evaluations of investigations conducted to detect and prevent transmission of tuberculosis. *JAMA* 2002; 287:991-995.
- 3- Reichler M R, Taylor Z, Castro KG. Factors in tuberculosis contact investigations [letter]. *JAMA* 2002; 297-294.
- 4- Vidal R, Miravittles M, Cayla JA, Torella M, De Gracia J, Morell F. Increased risk of tuberculosis transmission in families with microepidemics. *Eur Resp J* 1997; 10:1327-1331.
- 5- Jouban C, Koulla-shiro S, Lekama Assiene T, Hagbe P. Tuberculosis screening of patient contacts in 1993 and 1994 in Yaounde, Cameroon. *Med Trop* 1996; 56:156-158.
- 6- Topley J, Maher D, Nyong'onya, Mbewe L. Transmission of tuberculosis to contacts of sputum positive adults in Malawi. *Arch Dis Child* 1996; 74:140-143.

World Health Organization Communicable Diseases. Global Tuberculosis Control. WHO Report 2000. Geneva, Switzerland: WHO, 2000.

- 7- Marks SM, Taylor Z, Qualls NL, Shrestha-Kuwahara RJ, Wilce MA, Nguyen CH. Outcomes of contact investigations of infectious tuberculosis patients. *Am J Respir Critic Care Med* 2000; 162:2033-2038.
- 8- Krishna Murthy VV, Nair SS, Gothi GD, Chakraborty A K. Incidence of tuberculosis among newly infected population and in relation to the duration of infected status. *Indian J Tuberc* 1976;33:3-7.
- 9- Ansari S, Thomas S, Campbell IA, Furness L, Evans MR. Refined tuberculosis control tracing in a low incidence area. *Respir Med* 1998; 92:1127-1131.
- 10- Noertyoyo K, Tam C M, Chan SL, Tan J, Chan-yeung M. Contact examination for tuberculosis in Hong Kong is useful. *Int J Tuberc Lung Dis* 2003; 6(1): 19-24.
- 12- Gessner BD, Weiss WN, Nolon CM. Risk factors for pediatric tuberculosis infection and disease after household exposure to adult index cases in Alaska. *J Pediatr* 1998;132:509-513.

PPD Induration	Percent(%)
0-5	2.9%
6-9.9	23.5%
10-14.9	35.3%
>15	38.2%

() PPD :

0-15 years	1
15-30 years	2
30-45 years	3
>65	1

:

XRay Finding	Contacts	P.T.B cases
Cavitary	3 (2.04%)	3
Infiltration	6 (4.08%)	3
Calcification	37 (25.17%)	1
Nodule	3 (2.04%)	2
Normal	51 (66.67%)	

: