

()

**مقایسه کموتاکسی در استوباکتری دی ازوتروفیکوم و ریزویوم
ملیلوتی در شرایط آزمایشگاه**

NaCl Zn⁺⁺ Mn⁺⁺ .

(%) NaCl Zn⁺⁺ Mn⁺⁺ EDTA

EDTA

:

Chemotaxis of Acetobacter Diazotroph and Rhizobium Meliloti to Roots Exudates

G. Emtiazi, Z. Etemadi and A. Sanjabi

Department of Biology, University of Isfahan

Abstract

Considering biological fertilizer for non host plant, the effect of different compounds and root exudates on chemotaxis of two strains isolated from sugarcane and meliloti was examined. The presence of 1mM EDTA increased the chemotaxis of Rhizobium to Malic acid while decreased principle attractant to Aspartate. The Rhizobium was most attracted to Amino acids while Acetobacter was attracted to Cytokine and Jibereline. The presence of Mn, Zn cationic and NaCl in appreciable concentration reduced the movement of Rhizobium to roots.

Keywords: Chemotaxis, Acetobacter, Rhizobium, Root, Symbiosis

l...

.()

.()

Tn5

.()

.()

.()

.()

()o

.()

.()

() Aza
0.6 K₂HPO₄ 0.2 K₂HPO₄]:
0.01 FeCl₃ 0.002 Na₂M₈O₄ 0.2 MgSO₄ 0.002 CaCl₂
/ KOH / .[
() / PH
Aza
Aza
)

() % / .() (/ pH
()
.
.

Grim ()
()
()
(EDTA / K₂HPO₄
1.5 600nm
(% / +) 14ml 1ml

l...

°C

/ /)
°C .(

.()

mm

.()

(μl)

.()



:

EDTA (%)

(%) Mg

()

NaCl (%) Zn⁺⁺ (%) Mn⁺⁺ (%)

Zn⁺⁺ Mn⁺⁺ EDTA

EDTA

l...

() EDTA

EDTA

EDTA.
NaCl Zn⁺⁺ Mn⁺⁺

NaCl Zn⁺⁺ Mn⁺⁺

EDTA

NaCl Zn Mn EDTA

%

:

--	--	--

/

(mm)	(mm)	
		/ gr :
+	+	
	+	
+	+	
+	+	/ gr
		/ gr :
+	+	
+	+	
	+	/ gr :
	+	
+	+	
+	+	
+		
		μL :
	+	(20 ppm)
		(10 ppm)
	+	(5 ppm)

-L
-L
-L

:

|

/...

(Mm)						
			*			EDTA % /
			*			EDTA % /
						EDTA %
			*			Mg % /
						Mg %
			*			Mn % /
						Mn %
			*			Zn % /
						Zn %
						NaCl %
						NaCl %
	+	+	+	+	+	
	-L		-L			
						EDTA %
						Mg %
						Mn %
						Zn %
						NaCl %
	+	+	+	+	+	
						(ppm)
						EDTA %
						Mn %
						Zn %
						NaCl %
						Mg %

:*

:

	-		-												
		+		+		+		++		+		+			EDTA %
		+		+		+		+		+		+			Mn %
		+		+		+		+		+		+			
		+		+				+		+		+			
+		+		+		+		+		+		+			

(mM)

:

%

b	a
c	a
D	b
E	F

(mM)

:

)

.(

%

/...

c	b	a	E	a	a	EDTA %
	E	E	D	c	b	Mg⁺⁺ %
	D	D	E	c	a	Mn %
	D	E	E	c	b	ZN %
a	E	E	E	c	b	NaCl %

.()

.()

.()

.()

% /

/

.()

(NaCl)

Mg⁺⁺ EDTA

()

NaCl Zn⁺⁺ Mn⁺⁺

EDTA

%

EDTA EDTA

pH EDTA
NaCl

()

()

1. Steenhoudt, O. *Azospirillum*, a free-living nitrogen-fixing bacterium closely associated with grasses: genetic, biochemical and ecological aspects. FEMS Microbiology Reviews. 24, 487-506, 2000.
2. Zhulin, I. B., Houwearts, D. A major chemotaxis gene cluster in a *Azospirillum brasilense* and relationships between chemotaxis operon in Alpha-proteobacteria. FEMS Microbiology Letters. 201, 61-67, 2002.
3. Hawes, M. C., and Smith, L. Y. Requirement for chemotaxis in pathogenicity of *Agrobacterium tumefaciens* on roots of soil. J. Bacteriol. 10, 5668-5671, 1989.

l...

4. Bergman, K., Hoffee, G. Hovestadt, M. Larosiliere, R., and Ronco, P. G. Physiology of behavioral mutants of *Rhizobium meliloti*, J. Bacteriol. 170, 3249-3254, 1988.
5. Barbour, W. M., Hattermann, D. R. and Stacey, G. Chemotaxis of *Bradyrhizobium japonicum* to soybean exudates. Appl. Envi. Micro. 57, 2635-2639, 1991.
6. Shuh-Chyung, S., and Liang-Ping, L. The transition of *Rhizobium fredii* lipopolysaccharides induced by soybean root exudation. J. gen. Micro. 40, 773-778, 1999.
7. Cappel, D., Chemotaxis by *Pseudomonas syringae* to tomato. Appl. Enviro. Micro. 54, 623-632, 1988.
8. Scher, F. Klocppor, J., and Singleton, I. Chemotaxis of *Flouressent Pseudomonas* to soybean seed exudates invitro and in soil. Can. J. Microbiol. 31, 570-574, 1985.
9. Soby, S., Patrick, K., and Kosuge, B. Chemotaxis of *Pseudomonas. Syringae*. Enviro. Microbiol. 57, 2918-2920, 1997.
10. Alef, K., and Nannipieri, P. Methods in applied soil microbiology and biochemistry. UK. Harcourt, Brace and Company, 134-141, 1995.
11. Grim, A. C., and Harwood, G. Chemotaxis of *Pseudomonas* to the polyaromatic hydrocarbon naphthalene. Appl. Env. Micro. 63, 4111-4115, 1997.
12. Stock, J., and Sandra, D. R. Chemotaxis In: Encyclopedia of Microbiology, Academic Press. 1, 772-779, 2000.
13. Barbour, W. M., Hattermann, D. R., and Stacey, G. Chemotaxis of *Bradyrhizobium Japonicum* to soybean Exudates. Apl. Enviro. Micro, 57. 2635-2639, 1991.
14. Soby, S., Kirkpatrick, B., and Kosuge, T. Chemotaxis of *Pseudomonas syringae* virulence mutant. Apl. Enviro. Micro. 57, 2918-2923, 1991.
15. Futamata, H., Sakai, M. Ozawa, H., and Urashima, Y. Chemotactic response to aminoacids of *fluorescent Pseudomonads* isolated from spinach roots grown in soils with different salinity levels, Soil, Sci, Plant nut. 44, 1-7, 2001.
16. Emtiazi, G., Etemadifar, Z., and Tavassoli, M. A novel nitrogen-fixing cellulytic bacterium associated with root of corn is a candidate for production of single cell protein. Biomass and Bioenergy (In press), 2003.