



**Introducing of the New Genera and Species of Fusulinacea in the
Permian Rocks, Shirgesht Area**

A. Taheri^{*} and H. Vaziri Moghadam^{}**

***Department of Geology, Technical University of Shahrood**
****Department of Geology, University of Isfahan**

Abstract

Fusulinacea are numerous and diverse in the Permian rocks of Iran. In Shirgesht area (Baghe-Vang section), the lower part of the Jamal Formation is composed of shale, marl and intercalation of biogenic limestones. The limestone beds contain numerous Schwagerinids, including the Bolorian species. In the upper part, with increasing water depth, the abundance of fusulinids decrease and only small fusulinids (family Ozawainellidae) are present. The textural rocks and accompany of sponge spicules and radiolaria indicate a relatively deeper marine environment for deposition of the sediments. Detailed study of schwagerinid of the basal section has led to the introduction of new genus of Iranella and Paraleeina. This genus is similar to Paraschwagerina, except the fact that Paraschwagerina lack cuniculi and are not found in sediments younger than Sakmarian.

Key words: Permian, Fusulinid, Iranella, Shirgesht area

(Ross,

.1985)

"

()

(Braiser,

.(Ross, 1995)

.1995a)

.(Ozawa, 1987)

.(Ross, 1967; Loeblich and Tappan, 1988)

.(Ross, 1995)

Archive of SID

(Ruttner et al., 1968)

()



:

(Geinitzina)

..... (Paleotextularia)

(Tetrataxis)

()



Arce

(

)

(

(

(


()

(

(

(

(



(
(
)
Chalaroschwagerina vulgarisformis
Chalaroschwgerina (
vulgarisiformis,
Paraschwagerina shuleri
Wiliams, 1963
Iranella bella
)
Iranella longa
(
Schwagerina hyperborea (Salter) (B47-B65)
.)

Iranella orbiculata

Iranella pauca

Rauser-Chernousova

()

Iranella bella n. sp.
(Plate 1:Fig.6-7)

Order Schwagerinidae Solovieva, 1985
Genus Iranella
Type- spesies: Iranella bella n. sp.

)Etymology. Bellus (lat.)- beautiful.

Material. 2 axial sections.

Paraschwagerina

() shuleri Wiliams

Iranella bella

Iranella longa
(Plate 2:Fig.1-3)

Iranella pauca
(Plate 1:Fig.3-5)

Etymology. Paucus (lat.)-little, few
Material. 7 axial section.

1974 *Schwagerina hyperborea* (Salter) sensu Kahler, p.
95-96, pl.1, fig.10.
Etymology. Longus (lat.)-long
Material. 5 axial sections.


Order Schwagerinidae Solovieva, 1985
Genus Paraleeina

(A, B)


Type-species: *Leeina postkraffti* (Leven, 1967).

Iranella orbiculata
(Plate 1:Fig.1-2)

Etymology. Orbicularis (lat.)-rounded.
Material. 3 axial sections.



A



B

A, B *Paraleeina edoensis* (Ozawa).X10

(Parafusulina)

(Skinnerella)

(Leeina)

)

()

(

)

(

()

()

Genus Genus *Iranella* n.gen.

()

Paraleeina n.gen.

)

(

()

5- F., Kahler, Iranische Fusuliniden. Jahrb. Geol. B.-A., Bd.117, s. 75-107, Wien. (1974).

6- A. R., Loeblich, and H., Tappan, Foraminiferal genera and their classification. 2 volume. Van Nostrand Reinhold Company. New – York. (1988).

7- T., Ozawa, Permian fusulinacean biogeographic provinces in Asia and their tectonic implications. In: Taira, A., Tashiro, M. (Eds.), Historical Biogeography and Plate Tectonic Evolution of Japan and Eastern Asia. Terra Sci. Publ.Co., Tokyo, pp. 45-63. (1987).

8- D. M., Rauser-Chernousova, F.R., Bensh, M.V., Vdovenko, N.B., Gibshman, E.Ja., Leven, O.A., Lipina, F.A., Reitlinger, M.N. Solovieva, and I.O. Chediya, Reference-book on the systematics of Paleozoic foraminifera (Endothyrida, Fusulinida), Publishing House Nauka, 207 p. Moscow. (1996).

3- M. D., Braiser, a, Fossil indicators of nutrient levels. (part1) Evolution and extinction in relation to oligotrophy. In: Bosence, D. W. J., and Allison, P. A., (eds), 1995, Marine palaeoenvironmental analysis from fossils, 133, Geological Society Special Publication. No. 83, pp. 133-150. (1995).

4- M. D, Braiser, b, Fossil indicators of nutrient levels. (part2) Eutrophication and climate change. In: Bosence, D. W. J., and Allison, P. A. (eds), 1995, Marine palaeoenvironmental Analysis from fossils, 133, Geological society Special Publication No. 83, pp. 113-132. (1995).

...

- 12- C.A. Ross, and J. R. P., Ross, Foraminiferal Zonation of Late Paleozoic depositional sequences. *Mar. Micropaleont.* V. 26, pp. 469-478. (1995).
- 13- J., Stocklin, Structural history and tectonics of Iran: A review. *Am. Assoc. Petrol. Geol. Bull.* V. 52, No, 7, pp. 1229- 1258. (1968).
- 14- J., Stocklin, Stratigraphic Lexicon of Iran. Part 1:Central, North and East Iran. *Geol. Surv, Iran, Rep.* 18, 338 p. (1971).
- .
- 9- C.A., Ross, Development of fusulinid (Foraminiferida) Faunal realms. *Jour. Paleont.* V, 41, pp. 1341-1354. (1967).
- 10- C. A., Ross, Paleobiogeography of fusulinacean foraminiefera. In: *Studies in benthic Foraminifera benthos' 90*, Sendai, 1990, Tokai University Press. pp. 23-31. (1992).
- 11- C. A., Ross, Permian Fusulinacean. In: Scholle, P. A., Peryt, T. M., and Ulmer- Scholle, D. S., (Eds) *The Permian of Northern Pangea, V1: Paleogeography, Paleoclimates, Stratigraphy.* Springer-Verlag. Berlin, Heidelberg. pp. 167-185. (1995).