


* ** * *
* **
Archive of SID
()
)
(

(1965)

Archive of SID



Microni⁶
ENCARTA 98
 WORLD ATLAS

Abdulsamad and Barbieri, 1999 .

.(Wilson, 1975; Ghos, 1977; Benjamini, 1979;

: O₂

(2004)

(1962)

()

(Plate

O₁

.()

1b)

%

(1997)

- O

: O₁


: O₃

.(Plate 1c)

.(Plate 1a)

3. Bioclastic Wackestone-Packstone
4. Orbitoidae, Bioclastic Packstone

1. Open Marine Microfacies
2. Pelagic Foraminifera (Wackestone-Packstone)



()

Hottinger (1997)

...

(Plate

(Plate 1d)

.1e)

(

)

(1965)

()

)

Pomar

(.B2 B1

Zhicheng et al. (1997) , (1991)

()

()

()

)

: B

(Zhicheng et al., 1997)

(/

:B1

: B3

%

(Plate

()

(. a,b,c)

.1f)

:B2

)

(Buonocunto et al , 1999


: L

-
- 4. Bioclastic, Intraclastic Grainstone
 - 5. Lagoon Microfacies

-
- 1. Shelf Margine Microfacies (Rudist Biostrome /Barrier)
 - 2. Rudist Boundstone
 - 3. Rudist Bioclastic Grainstone



a



b



c

b .

a

c .

L1

(Plate 2a)

1. Miliolida, Bioclastic Packstone

(Plate 2c)

(Hypersaline)

(Sub-haline)

(1975)

(Geel,2000)

T

T₁

(Reiss and Hottinger ,1984; Hottinger,

1997; Geel, 2000)

L₂

(Plate 2b)

()

Zhicheng

(1997)

)

(Geel, 2000, Romero et al., 2002

L₃

3. Tidal Flat Microfacies
4. Mudstone

1. Dasycladacea, Bioclastic Wackestone-Packstone
2. Peloid, Bioclastic Packstone

(Wilson, 1975; Flugle, 1982; Read, 1985; Carrozi, 1989; Jones Desrochers, 1992, Burchette and Wright, 1992 and Geel, 2000)

(Tucker and Wright, 1990;

Wright and Burchette, 1996,)

(01)

...

(1997)

Zhicheng,

(O2,O3,O4)

(Ross

(O1)

.and Skelton, 1993)

.(Kauffman and Johnson, 1988)

.(Irvin,1965)

(Outer Shelf)

(O2,O3,O4)

.(Flugel, 2004; Read, 1985; Tucker and Wright, 1990;
Walker and James, 1992)

.(Stanley, 1981)

(Haq et al.,1987)

.(Haq et al., 1987)

.(Buonocunto et al.1999)

L1,L2,L3)

(

Hottinger, Buonocunto et al., 1999)

(T1)

.(1997;

(/)


James (1997)

Archive of SID


)

(


Plate 1




a




b




c



d



e




f


Explanation of Plate 1

- a-Pelagic Foraminifera Wackestone-Packstone, X53
- b-Bioclastic Wackestone-Packstone, X53
- c-Orbitoidae, Bioclastic Packstone, X33
- d-Rudist, Bioclastic Packstone, X53
- e-Rudist, Bioclastic Grainstone, X33
- f-Bioclastic, Intraclastic Grainstone, X53


Plate 2




a



b



c



d

Explanation of Plate 2

a-Miliolida, Bioclastic Packstone, X53

b-Dasycladacea, Bioclastic Wackestone-Packstone, X33

c-Peloid Bioclastic Packstone, X53

d-Mudstone, X33

()

()

- ...
- ()
- ()
- ()
- 15- E. O., Abdulsamad, and R., Barbieri, Foraminiferal distribution and palaeo-ecological interpretation of the Eocene-Miocene carbonates at Al Jabal al Akhdar (Northeast Libya): *Journal of Micropalaeontology*, v.18, pp. 45-65; (1999).
- 16- C., Benjamini, Facies relationships in the Avedat Group (Eocene) in the Northern Noger, Israel: *Israel J. Earth Sciences*, v. 28, pp. 47-69; (1979).
- 17- C., Briand, A., Izart, D., Vaslet, D., Vachard, M., Makhlina, N., Goreva, T., Isakova, O., Kossovaya, and A., Jaroshenko, Stratigraphy and sequence stratigraphy of Moscovian and Gzhelian in the Moscow Basin: *Bull. Soc. Geol. France*, t. 169, no. 1, pp. 35-52; (1998).
- 18- F. P., Buonocunto, B., D'Argenio, V., Ferreir, and R., Sandulli, Orbital cyclostratigraphy and sequence stratigraphy of Upper Cretaceous platform carbonates at Monte sant ' Erasmo, Southern Apennines, Italy: *Cretaceous Research*, v. 20, pp. 81-95; (1999).
- 19- T. P., Burchette. and V. P., Wright, Carbonate ramp depositional system: *Sedimentary Geology*, v. 79, pp. 3-57; (1992).
- 11- A. V., Carozzi, Carbonate Rock Depositional Model: Prentice Hall, Newjersy, 604p; (1989).
- 20- R. J., Dunham, Classification of carbonate rock according to depositional texture. In: W. E. Ham (ed.,) *Classification of carbonate rocks, a symposium*. Am. Assoc. Petrol. Geol., Mem. 1, pp.108-121; (1962).
- 21- E., Flugel, *Microfacies of Carbonate Rocks*: Springer-Verlag, Berlin, 976p; (2004).

- compelex, Mallorca, Spain: *Sedimentology*, v. 38, pp. 243-270; (1991).
- 33- J. F., Read, Carbonate platform facies models: *Am. Assoc. Petrol. Geol. Bull.*, v. 69, no. 1, pp.1-21; (1985).
- 26- H. G., Reading, *Sedimentary Environment and Facies*,: Blackwell Sci. Pub., 615p; (1996).
- 34- Z., Reiss, and L., Hottinger, *The Gulf of Aqaba: Ecological Micropaleontology*, *Ecol. Stud.* 50, Springer, Berlin, 354p; (1984).
- 35- D.J., Ross, and P. W., Skelton, Rudist formation of the Cretaceous: a paleoecological, sedimentological and stratigraphical review. In V.P Wright (ed.), *Sedimentology Review*, No. 1: Blackwell Scientific Publication, Oxford, p. 73-91 ; (1993).
- 36- G. D. Jr., Stanley, Early history of scleractinian corals and its geological consequences: *Geology*, v. 9, p. 507-511; (1981).
- 37- M. E., Tucker, and V. P., Wright, *Carbonate Sedimentology*, Blackwell Sci. Pub. Oxford, 471p; (1990).
- 38- R. G., Walker, and N.P., James, (3rd edition) *Facies model: Response to sea level change*, Geological Association of Canada, Reprint Series 1,409p; (1992).
- 39- J. L., Wilson, *Carbonate facies in geologic history*: Spiringer-Verlag, NewYork, 471p; (1975).
- 40- V.P., Wright, and T.P, Burchette, Shallow-water carbonate environments: In: H.G.Reading (ed.) *Sedimentary environments: Processes, facies and stratigraphy*, pp.325-394; (1996).
- 41- Z., Zhicheng, H., Willenms, and Z., Binggao, Marine Cretaceous-Paleogene biofacies and ichnofacies in southern Tiber, China, and their sedimentary significance: *Marine Micropalaeontology*, v. 32, pp. 3-29; (1996).
- 22- T., Geel, Recognition of stratigraphic sequence in carbonate platform and slope deposits: Empirical models based on microfacies analysis of Plaeogene deposits in Southeastern Spain, *Palaeo., Palaeo., Palaeo.*, v. 155, pp. 211-238; (2000).
- 23- B. K., Ghosc, Palaeoecology of the Cenozoic reefal foraminifers and algae - A brief review: *Palaeo., Palaeo., Palaeo.*, v. 22, pp. 231-356; (1977).
- 24- B.U., Haq, J., Hardenbol, and P.R., Vail, Chronology of fluctuating sea level since the Triassic: *Science*, v. 235, pp. 1156-1167; (1987).
- 25- L., Hottinger, Shallow benthic foraminiferal assemblages as signals for depth of their deposition and their limitations: *Soc., Geol., Fr., Bull.*, v. 168, pp. 491-505; (1997).
- 26- M.L., Irvine, General theory of epiric clear water sedimentation. *Amer. Assoc. Petrol. Geol. Bull.*, v. 46, no.4, pp.445-459; (1965)
- 27- N.P., James, The cool-water carbonate depositional realm, In: James, N.P. and Clarke, J.A.D., (eds.), *Cool-water carbonates: SEPM Special publication*. 56, pp.1-20; (1997).
- 28- G. A., James, and J. G. Wynd, Stratigraphic nomenclature of Iranian oil consortium agree ment area: *A.A.P.G. Bulletin*, v.49, p. 2182-2245; (1965).
- 29- B., Jones, and A., Desrochers, Shallow platform carbonates: In: R.G. Walker and N.P. James (ed.) *Facies models: Response to sea-level*, pp.277-301; (1992).
- 30- E.G., Kauffman, and C.C., Johnson, The morphological and ecological evolution of Middle and Upper Cretaceous reef building rudists: *Palaios, Reefs issue*, v.3, P. 194-216; (1988).
- 31- Y., Lasemi, Carbonate microfacies and depositional environment of the Kinkaid Formation (Upper Mississippian) of the Illinois Basin: Ph.D. Thesis, University of Illinois, U. S. A., 139p; (1979).
- 32- L., Pomar, Reef geometric, erosion surface and high frequency sea level change, Upper Miocene reef