

( )

( )

( ) ( ) : ( )

Planktonic foraminifera mudstone-wackestone, Bioclast planktonic foraminifera wackestone-packstone, Planktonic foraminifera Heterostegina packstone, Bioclast Miogypsinid corallinacea packstone, Coral and corallinacea floatstone, Rotalia wackestone-packstone, Rotalia miliolid wackestone-packstone, Foraminifera (Rotalia, Meandropsina, Borelis, Archaias, Peneroplis) miliolid packstone-grainstone, Bioclast corallinacea miliolid packstone, Miliolid grainstone, Intraclast bioclast packstone, Bioclast wackestone-packstone and quartz mudstone.

) ( :

**Depositional Environment of the Upper Part of Asmari Formation (Burdigalian)  
Based on Microfacies and Paleoecology of Foraminifera, Southeast of the Laki  
Mountain, Southern Lordegan**

**S. Hakimzadeh, A. Seyrafian and H. Torabi**  
**Geology Department, the University of Isfahan**

**Abstract**

The upper Asmari Formation in southeast Laki Mountain comprises of 376 meters of thin, medium and thick beds of limestone interbedding with marly limestone. It is overlies by the Gachsaran formation. Based on field work and laboratory analysis, 13 microfacies related to basin (outer ramp), open marine (middle ramp), bar and lagoon (inner ramp) environments were recognized, as follows: Planktonic foraminifera mudstone-wackestone, Bioclast planktonic foraminifera wackestone-packstone, Planktonic foraminifera Heterostegina packstone, Bioclast Miogypsinid corallinacea packstone, Coral and corallinacea floatstone, Rotalia wackestone-packstone, Rotalia miliolid wackestone-packstone, Foraminifera (Rotalia, Meandropsina, Borelis, Archaias, Peneroplis) miliolid packstone-grainstone, Bioclast corallinacea miliolid packstone, Miliolid grainstone, Intraclast bioclast packstone, Bioclast wackestone-packstone and quartz mudstone. Paleoenvironment reconstruction based on the occurrence of foraminifera, related to microfacies and environmental factors (light, depth and water energy) reflects that deposition of the Upper Asmari Formation (Burdigalian) was in a carbonate ramp platform.

**Keywords:** Microfacies, Asmari Formation, Burdigalian, Foraminifera, Paleoecology

& Mayo, 1918)

(Richardson, 1924)

(Boech,

Lees & Richardson, 1929)

( )

(Thomas, 1948)

(Busk

Meandropsina iranica :


- Pseudotaberina malabarica (Henson) (Wynd, 1965)
- Dendritina Borelis pygmaea (Hanzawa) (Adams & Bourgeois, 1967)
- Peneroplis evolutus rangi (d'Orbigny) ( )
- Archaias kirkukensis (Henson) (Henson)
- Austrotrillina howchini (Schlumberger) ( )
- Amphistegina sp. Valvulinid sp. ( )
- (Adams& Bourgeois, Heterostegina sp. ( ))
- 1967; Loeblich & Tappan, 1998
- (Wynd, (Adams& Bourgeois, 1967)  
1965)
- (Dunham, 1965)
- (Embry & Klovan, 1972)
- (Carozzi, (Lasemi, 1972)  
1989)
- (Wilson, 1975)

Borelis melo (Fichtel  
& Moll) curdica (Reichel)

Globigerinoides      Glogigerina      ( )  
Basin ) : (Ba)  
. ( a ) : (Microfacies

: Ba1

(Planktonic      foraminifera      mudstone/  
wackestone)



.( )  
. ( )

Open ) : (O) : (Marine Microfacies  
 (Planktonic foraminifera Heterostegina  
 packstone) : (Hallock &  
 Glen, 1986)  
 (Beavington-penney & Racey,  
 2004)

Operculina Heterostegina : (Geel, 2000)

Amphistegina Elphidium Miogypsina Asterigerina : Ba2  
 Globorotalia Globigerina : (Bioclast planktonic foraminifera  
 wackestone/packstone)

Textularia : (c) : (Romero et al., 2002)

) : (Heterostegina : (Globigerina) : (Globigerinoides)

Operculina (Rasser et al., 2005) : (b) : (Romero et al., 2002)

Heterostegina Operculina : (Romero et al., 2002)

(Geel, 2000)

( d )

) Miogypsinid :

(Geel,

(

.2000)

:O2

(Bassi,

(Bioclast Miogypsinid corallinacea  
packstone)

.2005)


Miogypsina

Amphistegina

Miogypsinooides

(corallinacea)

Textularia



### توصیفات سنگ شناسی

تناوب مارن قرمز، کثیف و آبدرفت.

آهک متوسط لایه ای.

دو توالی از آهک متوسط تا ضخیم لایه به رنگ کرمی تا خاکستری، گاهًا توده ای، با میان لایه آهک مارنی در کف هر توالی، حاوی بریوزوثر و ساقه کربنوتید، هر توالی به سمت بالا ضخیم لایه شده و در مجموع ضخامت هر توالی به سمت بالا افزایش می یابد.

چهار توالی از تناوب آهک نازک، متوسط و ضخیم لایه، به رنگ کرمی مایل به خاکستری و خاکستری، در قاعده هر توالی آهک مارنی نودولار و آهک مارنی وجود دارد.

چند توالی از آهکهای نازک و متوسط لایه، به سمت بالا ضخیم لایه و توده ای، به رنگ کرمی تا خاکستری، حاوی آثار بیوکلاست.

چهار توالی از آهکهای نازک، متوسط و ضخیم لایه، گاهًا بیوکلاست دار با میان لایه آهک مارنی، به رنگ کرمی، کرمی مایل به صورتی، کرمی مایل به خاکستری و خاکستری، در برخی لایه ها تخلخل به صورت پراکنده وجود دارد.

چند توالی از تناوب آهکهای نازک، متوسط و ضخیم لایه، با میان لایه آهک مارنی، به رنگ کرمی مایل به خاکستری تا خاکستری، برخی لایه ها متخلخل.

چند توالی از تناوب آهکهای نازک، متوسط و ضخیم لایه به سمت بالا ضخیم لایه تو، با میان لایه های آهک مارنی، به رنگهای خاکستری مایل به قرمز، خاکستری مایل به صورتی، کرمی مایل به خاکستری تا خاکستری با تخلخل پراکنده.



(Rotalia ) ( Barrier ) : (B)  
 .( f ) : (Microfacies  
 Rotalia : (Coral and : B1  
 ( ) corallinacea floatstone)  
 .(Geel, 2000) ( )

Rotalia -  
 (Rotalia - : L2  
 miliolid wackestone/packstone)

Rotalia miliolid Discorbis Lagoon ) : (L)  
 .( g ) : (Microfacies  
 miliolid Rotalia : (L1  
 wackestone/packstone)  
 Rotalia

(Beavington.penney& Racey,  
 miliolid .2004)  
 Rotalia )

(Romero et al.,

.2002; Rasser et al., 2005)

Rotalia .

(Rasser

.(Geel, 2000)

.et al., 2005)

( )

:L3

.( )

Foraminifera (Rotalia, Meandropsina, Borelis, Archaias, Peneroplis) miliolid packstone/grainstone

( ) ( )

( )

miliolid.

(Bioclast corallinacea miliolid packstone)

## Meandropsina      Archaias      Peneroplis

Rotalia

b )

(Cosovic et al.,

2004)

,( a h )

(Rasser et al., 2005)


### (Peneroplids )


Peneroplid.


### (Miliolid grainstone)

L5


Borelis








a- Planktonic foraminifera mudstone/wackestone (Ba1), b- Bioclast planktonic foraminifera wackestone/packstone (Ba2), c- Planktonic foraminifera Heterostegina packstone (O1), d- Bioclast Miogypsinitid corallinacea packstone (O2), e- Coral and corallinacea floatstone (B1), f- Rotalia wackestone/packstone (L1), g- Rotalia miliolid wackestone/packstone (L2), h- Foraminifera (Rotalia, Archaias, Lepidocyclus) miliolid packstone/grainstone (L3). [D: Ditrupa, P: Planktonic foraminifera, Mio: Miogypsina, Bio: Bioclast, Co: Corallinacea, Het: Heterostegina, C: Coral, Ro: Rotalia, Mi: Miliolid, L: Lepidocyclus, Ar: Archaias; All scale bars are 0.25 mm in length]



a- Foraminifera (Borelis, Peneroplis) miliolid packstone/grainstone (L3), b- Bioclast Corallinacea miliolid packstone (L4), c- Miliolid grainstone (L5), d- Intraclast bioclast packstone (L6), e- Bioclast wackestone/packstone (L7), f- quartz mudstone (L8) [Bo: Borelis, Bio: Bioclast, Mi: Miliolid, Pe: Peneroplis, Co: Corallinacea, In: Intraclast; All scale bars are 0.25 mm in length]

( )  
Amphistegina  
(O2) Miogypsina  
)  
( L8 L1)  
Borelis Peneroplis Archaias)  
(miliolid  
(Operculina Heterostegina)  
(O1)  
Operculina  
(Allen et al.,  
2001)  
Ba1)  
(  
(Ba2



(Geel, 2000)


(Rasser et al., 2005)

( )  
 ( )  
 (Beavington-penney & Racey,  
 .2004)  
 (Beavington-penney & Racey,  
 .2004)  
 (Rasser  
 Amphistegina .et al., 2005)  
 Borelis pygmaea ) Borelis  
 .( ) ( )  
 Heterostegina



(Sample no. H50, X40)

(Sample no.



**Amphistegina** (a)

**Borelis** (b)

S94, X40)

Amphistegina

( )



a



b

افزایش نور و انرژی

Amphistegina

(Sample no. H21, X40)

(Sample no. S93, X40)

(a

(b

Heterostegina

(Meandropsina )

(Miliolid Borelis )

(Amphistegina )

)

(Operculina Heterostegina

کاهش ارزی



a

[فرا]یش عمیق



b

Heterostegina

(Sample no.S91, X40)

Heterostegina

(a

(Sample no.H21, X25)

Heterostegina

(b

( )

- 7- Beavington-penney, S. J. and A. Racey, Ecology of extant Nummulitids and other larger benthic foraminifera, applications in palaeoenvironmental analysis: Earth-Science,v. 67,p. 219-265, 2004.
- 8- Busk, H. G.and H. T. Mayo, Some notes on the geology of the Persian oilfields. Journal Institute Petroleum Technology 5, 5-26, 1918.
- 9- Carozzi, A. V., Carbonate rocks depositional model: Pentice Hall New Jersy, 604p, 1989.
- 10- Cosovic, V., K. Crobne, and A. Moro, Palaeoenvironmental model for Eocene foraminiferal limestones of the Adriatic carbonate platform (Istrian Peninsula): Facies,v. 50, p. 61-75, 2004.
- 11- Dunham R. J., Classification of carbonate rocks according to their depositional texture. AAPG Mem. 1, p. 108–121, 1962.
- 12- Embry, A. F. and E. J. Klovan, Absolute water limits of late Devonian paleoecological zones: Geol. Rdsch, 1972.
- 13- Geel, T., Recognition of stratigraphic sequences in carbonate platform and slope deposits, empirical models based on microfacies analysis of Paleogene deposits in Southeastern Spain: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 155, p. 211-238, 2000.
- 14- Hallock, P., and E. C. Glenn, Larger foraminifera: a tool for paleo-environmental analysis of Cenozoic carbonate depositional facies, In Beavington-Penney, S. J. and A. Racey (Eds.), Ecology of extant Nummulitids and other larger benthic foraminifera, applications in palaeoenvironmental analysis: Earth-Science, v. 67, p. 219-265, 1986.
- 15- Lasemi, Y., Carbonate microfacies and depositional environment of the Kinkoid Formation (Upper Mississippian) of the Illinios basin: PhD. Thesis., University of Illinios, U.S.A., 139p, 1979.
- 16- Lees, G. M., and F. D. F. Richardson, The geology of the oil field belt of S. W. Iran: Geol. Mag. Iraq., v. 67, no. 3, p. 227-252, 1940.
- 4- Adams, T. D. and F. Bourgeois, Asmari biostratigraphy: Geological and Exploration, IOOC Report, V. 1074, Unpublished, 1967.
- 5- Allen, P. A., P. M., Burgess, J., Galewsky, and H. D., Sinclair, Flexural-eustation numerical model for drowning of the Eocene perialpine carbonate ramp and implications for Alpine geodynamics: Bulletin, v. 113, no. 8, p. 1052-1066, 2001.
- 6- Bassi, D., Larger foraminiferal and Coralline algal facies in an Upper Eocene storm-influenced, shallow water carbonate platform (Colli Beici, Vrth - Eastern Italy): Palaeogeography, Palaeoclimatology, Palaeoecology, v. 226, p. 17-35, 2005.

- 23- Seyrafian, A., Microfacies and depositional environment of the Asmari Formation at the Dehdez area: Carbonates and Evaporites, v. 15, no. 2, p. 121-130, 2000.
- 24- Thomas, A. N, The Asmari Limestone of southwest Iran: AIOC Report, 730 p. Unpublished, 1946.
- 25- Vaziri-Moghaddam, H., M. Kimiagari and A. Taheri, Depositional environment and sequence stratigraphy of the Oligocene-Miocene Asmari Formation in SW Iran: Facies, v.52, p.41-51, 2006.
- 26- Wilson, J. L., Carbonate facies in geology history: New York, Springer – Verlag, 471 p, 1975.
- 27- Wynd, J. G., Biofacies of the Iranian oil consortium agreement area: IOOC Report, v. 1082, 40 Plates, 80 p. Unpublished, 1965.
- 17- Loeblich, A. R., and H., Tappan, Foraminiferal Genera and their Classification: Van Nostrand Reinhold, New York, 970 p, 1998.
- 18- Pomar, L., Type of carbonate platforms, a genetic approach: Basin research, v. 13, p.313-334, 2001b.
- 19- Rasser, M. W., C. Scheibner, and M. Mutti, A palaeoenvironmental standard section for early Ilerdian tropical carbonate factories (Corbieres, France, Pyrenees, Spain): Facies, v. 51, p. 217-232, 2005.
- 20- Romero, J. E. Caus, and J. Rosell, A model for the palaeoenvironmental distribution of larger foraminifera based on late Middle Eocene deposits on the margin of the South Pyranean basin (NE Spain): Palaeogeography, Palaeoclimatology, Palaeoecology, v. 179, p. 43-56, 2002.
- 21- Seyrafian, A., H. Vaziri, and H. Torabi, Biostratigraphy of the Asmari Formation, Burujen area, Iran: J. Sci., I. R. Iran, v. 7, no. 1, p. 31-48, 1996.
- 22- Seyrafian, A. and A. Hamedani, Microfacies and depositional environment of the Upper Asmari Formation (Burdigalian), north-central Zagros Basin, Iran: N. Jb. Geol. Palaont. Abh., v. 210, p. 129 –141, 1998.