

(Acipenser persicus)

*

..

(Acipenser persicus)

(p< /)

:

[]

)

(

()

[]

(*Acipenser baeri*)

()

[]

()

[]

[]

[]

()

()

[]

[]

(*Seriola quinqueradiatoy*)

()

(*Acipenser persicus*)

cm

g

[]

()

L

[]

:

3. Devitsina

1. Kohbara
2. Hidaka

[]

:

(

cm* *

s

cm

()

()

.()

s

.()

()

()

.[]

.()


s

[]

()

t



‘ ‘ ‘ ‘ ‘



S



()



S

(p < /))

(

) .()

(

.()

()

() .(p < /)


.(p > /)

(p < /)

.()

)

(



()
 (%)
 () (%)

							← ()
/					*	**	↓
/							
/							
/	*						
/							
/	**		*				
/							
/							
/						*	
/							
/		*					
/	*	**			**		
/						*	
/						*	
/							
/		*					
/						*	
/							
/	*	/	**	/	*	**	/
/							

p< / ** p< / *

(p < /)

(p > /)

(p < /)

(p < /)

[] ()

:

()

s ()

()

() (p < /)

()

(p < /)

()

(p < /)

()

(p < /)

(p > /)

Fundulus heteroclitus

[]

(*Acipenser gueldenstadti*)
 (A.medirostris) (A.baeri)
 [] (Huso huso) (A.stellatus)

[]

[]

(*Menidia menidia*)
 (*Pseudopleuronectes americanus*)

- [1] Kuzmin S., Mironov S., Vostroushkin D., Shutov V.; «Behavioral responses to various chemical incentives in hybrid Beluga×Russian Sturgeon (*Huso huso*×*Acipenser gueldenstaedtii*) fry»; *J. Appl. Ichthyology*; 1999; 15: 233-236.
- [2] Kasumyan A. O., Prokopova O. M.; «Taste preferences and the dynamics behavioral taste response in the Tench, *Tinca tinca* (Cyprinidae)»; *J. of Ichthyology*; 2001; 41: 640-653.
- [3] Kohbara j., Hidaka I., Morishita T., Miyajima T.; «Gustatory and olfactory sensitivity to extracts of jack mackerel Kasumyan A. O., Doving K. B.; Taste preferences in fishes; *Fish and Fisheries*; 2003; (4): 289-347.
- [4] Muscle in young yellowtail *Seriola quinqueradiata*»; *Aquaculture*; 2000; 181: 127-140.
- [5] Hidaka I., Kohbara J., Araki T., Morishita T., Miyajima T., Shimizu S., Kuriyama I.; «Identification of feeding stimulants from a jack mackerel muscle extract for young yellowtail *Seriola quinqueradiata*»; *Aquaculture*; 2000; 181: 115-126.
- [6] Kasumyan A. O., Taufik L. R.; «Behavior reaction of juvenile Sturgeons (Acipenseridae) to amino acids»; *J. of Ichthyology*; 1994; 34: 90-102.
- [7] Devitsina G. V.; «Morphology of primary projections of chemosensory systems and some aspects of their interaction in the brain of Acipenseridae»; *J. of Ichthyology*; 2000; 40: 64-74.

() []

- []
- [10] Kasumyan A.O.; Olfactory sensitivity of the sturgeons to free amino acids; *Biophysics*; 1994; 39, 519-522.
- [11] Kasumyan A. O., Kazhlaev A. A.; «Formation of searching behavioral reaction and olfactory sensitivity to food chemical signals during ontogeny of sturgeons (Acipenseridae»); *Journal of Ichthyology*; 1993; 33, 51-65.
- [12] William E. S. C., James C. N. III., Richard A. G., Charles D. D.; «Stimulants of feeding behavior in fish: analysis of tissues of diverse marine organisms»); *Reference, Biol.Bull.*; 1996; 190: 146-160.
- [13] Mearns K. J.; «Sensitivity of brown trout (*Salmo trutta* L.) and Atlantic salmon (*Salmo salar* L.) fry to amino acids at the start of exogenous feeding»); *Aquaculture*; 1986; 55, 191-200.
- [14] Devitsina G. V.; «On the chemosensory and tactile provision of feeding behavior in Codfishes of the White Sea»); *J. of Ichthyology*; 1997; 37: 87-92.
- [15] Kasumyan A. O., Devitsina G. V.; «The effect of olfactory deprivation on chemosensory sensitive and the state of taste receptors of Acipenserids»); *J. of Ichthyology*; 1997; 37: 786-798.
- [16] Kasumyan A. O., Marusov E. A.; «Behavioral responses of intact and chronically anosmiated Minnow *Phoxinus phoxinus* (Cyprinidae) to free amino acids»); *J. of Ichthyology*; 2003; 43: 528-539.