

## *Streblospio gynobranchiata*

( )

(Spionidae)

\*

) *Streblospio gynobranchiata*

m

(

± m

/ ± / m

± m

/ g/m ± / m

*Streblospio gynobranchiata* :

. [ ]

. [ ]

Spionidae

*S. gynobranchiata*

(> cm)

,*Streblospio* [ ]


[ ]

[ ]

*Streblospio*

[ ]

m [ ]  
[ ]  
km  
m  
% ( )  
°C  
pH  
[ ]  
km  
( )



[ ]

Van Veen

(p < / )

cm

cm

[ ]

[ ]

( )

%

cm

[ ]

( )

[ ]

[ ]

mm

/

( m )

( cm )

[ ]

°

(



°)

( )

[ ]

SPSS

( )



*S. gynobranchiata*

/ **	/ *	in/m	
/ **	/ **	g/m	
/ **	/ **	in/m	
/ **	/ **	g/m	
/ **	/ **	in/m	
/ **	/ **	g/m	
/ **	/ **	in/m	
/ **	/ **	g/m	

/ \*\*

/ \*

(p < / )

( )

( )

*S. gynobranchiata*

/	/	/	/	F
/ **	/ **	/	/ *	P
/	/	/	/	F
/ **	/ **	/ **	/ *	P

/ \*\*

/ \*

*S. gynobranchiata*

<b>m</b>	<b>m</b>	<b>m</b>	
/ ± / a	± / a	/ ± / b	
/ ± / a	/ ± / b	/ ± c	
/ ± / a	/ ± / b	/ ± / c	
/ ± / a	/ ± / b	/ ± / c	

*S. gynobranchiata*

<b>m</b>	<b>m</b>	<b>m</b>	
/ ± / a	/ ± / a	/ ± / b	
/ ± / a	/ ± / b	/ ± / c	
/ ± / a	/ ± / b	/ ± / c	
/ ± / a	/ ± / b	/ ± / c	

( $p < /$  )

( )

( )

*S. gynobranchiata*

<b>m</b>	<b>m</b>	<b>m</b>	
/	/	/	F
/ **	/ **	/ **	P
<b>m</b>	<b>m</b>	<b>m</b>	
/	/	/	F
/ **	/ **	/ **	P

/ \*\*

*S. gynobranchiata*

/ ± / d	/ ± / a	/ ± / c	/ ± / b	<b>m</b>
/ ± / d	/ ± / a	/ ± / b	± / c	<b>m</b>
/ ± / d	/ ± / b	/ ± / a	/ ± / c	<b>m</b>

*S. gynobranchiata*

/ ± / d	/ ± / a	/ ± / c	/ ± / b	<b>m</b>
/ ± / d	/ ± / a	/ ± / b	/ ± / c	<b>m</b>
/ ± / b	/ ± / a	/ ± / a	/ ± / b	<b>m</b>

[ ] ( ) / g

*S. gynobranchiata*

( ) / g

/ /

/ °C / /

% /

[ ]

( / ) [ ]

(*Streblospio*)

[ ]

( )

[ ]

*S. gynobranchiata*

[ ]

)

(

)

*S. gynobranchiata*

(

/ g

( )

/ g

)

/ g/m

( / g

( )

/ g

)

*S. shrubsolii S. benedicti*

( / g

[ ]

[ ]

*S. benedicti*

*S. shrubsolii S. benedicti*

( ) / g /

[ ]

[ ]

Biscay

*S. benedicti*

Danny Eibye – Jacobsen

*Streblospio gynobranchiata*

[ ]

*S. gynobranchiata*

[ ]

- [1] Williams J. D.; A new species of *Polydora* (Polychaeta: Spionidae) from the Indo-West Pacific and first record of host hermit crab egg predation by a commensal polydorid worm. *Zoological Journal of the Linnean Society* 129: 537- 548. [ ]
- [2] Pearse V., Pearse J., Buchsbaum M., Buchsbaum R.; *Living Invertebrates*. Blackwell Scientific Publication and Box Wood Press; 1987; pp. 847.
- [3] Sarda R., Martin D.; Population of *streblospio* (Polychaeta:Spionidae) In temperate zone: dsemography and production; *J. mar. boil. Ass.U.K*; 1993; 73: 769 - 584.
- [4] Rossi F., Lardicci C.; Role of nutritive value of sediment in regulating population dynamics of the deposit-feeding polychaeta *streblospio shrubsolii*; *Marine Biology*; 2002; 140: 1129 - 1138.
- [5] Lardicci C., Ceccherelli G., Rossi F.; *Streblospio shrubsolii* (Polychaeta: Spionidae): temporal fluctuation in size and reproduction activity; *Cah Biol Marl*; 1997; 38: 207 - 214.
- [6] Rice A. S., Levin L. A.; *Streblospio gynobranchiata*, a new spionid polychaete species (Annelida: Polychaeta) from Florida and the Gulf of Mexico with an analysis of phylogenetic relationships within the genus *Streblospio*; *Proceeding of the Biological Society of Washington*; 1998; 111: 694- 707.
- [8] Grigrovich I. A., Therriault T .W., MacIsaac H. J.; History of aquatic invertebrate invasions in the Caspian Sea; *Biological Invasions*; 2003; 5:103- 115. [ ]
- [10] Mistri M., Fano E. A., Ghion F., Rossi R.; Disturbance and community patern of Polychaetes Inhabiting Valle Magnavacca (Valli di Comacchio, Northern Adriatic Sea, Italy); *Mrine Ecology*; 2002; 23(1): 31- 49.
- [11] Tolosa I., Mora S. D., Sheikholeslami M. R., Villeneuve J. P., Bartocci J., Cattini C.; Liphatic and aromatic hydrocarbons in coastal Caspian Sea sediments; *Marine Pollution Bulletin*; 2004; 48: 44-60.
- [12] Muniz P., Pires A. M. S.; Polychaete association in a subtropical environment (Sao Sebastiao Channel, Brazil): A structureal analysis; *Marine Ecology*; 2000; 21 (2): 145 – 160.



- [13] Cinar M.E., Ergen Z., Dagli E., Petersen M.E.; Alien species of spionid polychaetes (*Streblospio gynobranchiata* and *Polydora cornuta*) in Izmir Bay, eastern Mediterranean; *Journal of the Marine Biological Association of the United Kingdom*; 2005; 85: 821-827.
- [14] Abrantes A., Pinto F., Moreira M. H.; Ecology of polychaete *Nereis diversicolor*, in the Canal de Mira (Ria de Aveiro, Portugal): Population dynamics, production and oogenic cycle. *Acta Oecology*; 1999; 20(4): 267-283.
- [15] . . . [ ]
- [16] . . . [ ]
- [17] Nybakken J. W.; *Marine Biology and ecological approach*; Harper Collins College; pp. 445.
- [18] Rozali Othman M., Samat A., Hoo L. S.; The effect of bed sediment quality on distribution of macrobenthos in Labu River system and selected site in Langat River, Malasia; *Online Journal of Biological Science*; 2 (1): 32 – 34.
- [19] Garcia -Arberas L., Rallo A.; Population dynamics and production of *Streblospio benedicti* (Polychaeta) in a non-polluted estuary on the Basque coast (Gulf of Biscay); *SCI. MAR*; 2004; 68 (2): 193- 203.
- [20] Essink K., Dekker R.; General pattern in invasion ecology tested in the Dutch Wadden Sea: the case of a brackish-marine polychaetous worm; *Biological Invasions*; 2002; 4: 359- 368.