



تاریخ دریافت ۸۳/۱۰/۱۶ پذیرش مقاله ۸۴/۰۱/۲۵

CAH :

CAH :

% / :

II % CAH III % / II :

( $p < /$  ). III % II % :

( ) :

( ) :

( )

RA ( )

( )

RA

RA

( )

:

b a ) SI (cm<sup>2</sup>) = a<sup>cm</sup> × b<sup>cm</sup>

(

( )

SI

(grades)

( )

CAH

:

G0= - cm<sup>2</sup>, G1= - cm<sup>2</sup>, G2= - cm<sup>2</sup>,

G3= - cm<sup>2</sup>, G4= - cm<sup>2</sup>, G5 ≥ cm<sup>2</sup>

:

( )

( ) SI

Mean±SD

( )

( )

Logical GE ) Real-Time

/ MHZ (EUB Hitachi

MHZ

± cm<sup>2</sup>

cm<sup>2</sup> (p < / ) / ± / cm<sup>2</sup>

% /

± /

) (

II

( ) (

<sup>4</sup> Grades

<sup>1</sup> Nishiya

<sup>2</sup> Chronic Active Hepatitis - (CAH)

<sup>3</sup> Blind

C.T scan

$\pm \text{ cm}^2$  SI

$\pm \text{ cm}^2$  SI

SI

IV,III

( ) SI

( )

MRI

( )

( )

MRI ,C.T Scan

C.T Scan

( )

( )

SI

%

( $p < /$  ) .

%

II

%

II

II

( $p = /$  )

III % /

III %

III

( $p < /$  )

%

II % I %

III

II I

SI  
 $\pm / \text{cm}^2$                        $/ \pm / \text{cm}^2$   
 SI  
 I  
 $\text{cm}^2$      $\text{cm}^2$     SI  
 I  
 $\text{cm}^2$      $\text{cm}^2$   
 a    b a  
 b     $\text{cm}^2$   
 )    %    II    %    I    %  
 (    SI    II     $\pm \text{cm}^2$   
 CAH    ( )  
     SI ( )  
     SI     $\pm \text{cm}^2$   
 )    I  
 (...    I

:

	O = (SI ≤ cm <sup>2</sup> )		I = (SI = - cm <sup>2</sup> )		II = (SI = Z - cm <sup>2</sup> )		III = (SI = - cm <sup>2</sup> )		IV = (SI = - cm <sup>2</sup> )		V = (SL ≥ cm <sup>2</sup> )			
													Mean ± SD	
													$\pm / \text{cm}^2$	
			/				/						$\pm \text{cm}^2$	
C.A.H													$, \pm / \text{cm}^2$	

**References:**

1. Cutran k: Pathologic Basis of Disease. 5<sup>th</sup> Ed, New York, Mosby, 1994: 747-753.
2. Denhis LK et al: Harrisons Principles of internal Medicine. 15<sup>th</sup> Ed, New York, McGraw-Hil, 2001: 362-365, 669, 714-726, 1761.
3. Picardi M et al: Measurment of spleen by ultrasound in patients with thrombocytosis: a prospective study. Blood, 2002, 99(11): 4228-4230.
4. Ishibashi H, Higuchi N,shimamura R: Sonographic assessment and grading of spleen size. J clin ult, 1994, Jan; 19(1): 21-5.
5. MCGahan JP, Goldberg BB: Diagnostic ultrasound. 1<sup>st</sup> Ed, Philadelphia, Lippincott-Raren, 1998: 776-779.
6. Patrick CF, Giles WS: Margulies Alimentary tract Radiology. 5<sup>th</sup> Ed, <sup>st</sup> Louis, Mosby, 1994: 28-34, 1759-1762, 1771-1774.
7. Al-Salem AH et al: Sonographic Assessment of spleen size in Saudi patients with sickle cell disease. Ann Saudi med, 1998,18(3): 214-220.
8. Nishiya K, Hisakawa N and co-woorkers: Enlarged spleen detected by abdominal ultrasonography in patients with RA. Ann Rheu dis, 2000, 59: 750(a).
9. Konus Ol, Ozdemir A, Akkaya A: Normal liver, spleen, and kidney dimensions in children: evaluation with sonography. Am J of Roe, 1998, 171: 1693-1698.
10. Megremis SD, vlachonikolis IG, Tsilimigaki AM: Spleen length in childhood with us: normal values based on age, sex, and somatometric parameters. Rad, 2004, 231: 129-134.
11. Okumura Y et al: Differentiation of mononucleosis from, hepatitis by sonographic measurement of spleen size. JCU, 1987, Jun 15(5): 3-6.
12. Sutton D: Textbook of-Radiology and imaging. 7<sup>th</sup> Ed, London, churchill livingstone, 2003: 746-749, 758 -7759, 775-777.
13. MCDonald MN, Shoemakerk: Splenic in -der in thalasemic. Hematologic Budap, 1995: 26(4): 205-209(ABS).
14. Meschan I: An Atlas of Anatomy Basic to Radiology. 3<sup>th</sup> Ed, New York, Saunders. 1975: 982-987.
15. Yetter EM, Acosta KB, Olsom MC, Blundell K: Estimating splenic volume: sonographic Measurements correlated with Helical CT determination. AJR, 2003, 181: 1615-1620.