

Tarbiat Moallem University, 20th Seminar on Algebra,
2-3 Ordibehesht, 1388 (Apr. 22-23, 2009) pp 14-15

ON THE CHARACTERIZATION OF SIMPLE GROUPS $B_n(q)$
AND $C_n(q)$

NEDA AHANJIDEH* AND ALI IRANMANESH

Department of Mathematics
Tarbiat Modares University
P.O.Box: 14115-137, Tehran, Iran
iranmana@modares.ac.ir

ABSTRACT. Let G be a finite group and $\pi(G)$ be the set of prime divisors of the order of G . For $t \in \pi(G)$ denote by $n_t(G)$ the order of a normalizer of t - Sylow subgroup of G and put $n(G) = \{n_t(G) \mid t \in \pi(G)\}$. In this talk, we discuss about an answer to the following problem for the simple groups of Lie type B_n, C_n :

Let L be a finite non-abelian simple group and G be a finite group with $n(L) = n(G)$. Is it true that $L \cong G$?

Also, we discuss about difference between orders of the solvable subgroups of the non-isomorphic simple groups $B_n(q)$ and $C_n(q)$.

1. INTRODUCTION

Characterization by orders of Sylow normalizers has first been considered by Bi in 1992 (see [2]). It is known that if G is $A_n(q)$, ${}^2A_n(q)$, $C_2(q)$, ${}^2D_n(q)$, alternating group, Mathieu simple groups, Janko groups and $Sz(2^{2m+1})$, then G is characterizable by orders of Sylow normalizers. In this talk, we discuss that if $n = 2$ or $q \not\equiv \pm 1 \pmod{8}$, then $B_n(q)$ and $C_n(q)$ are characterizable by orders of Sylow normalizers and otherwise, $B_n(q)$ and $C_n(q)$ are 2-recognizable by orders of Sylow normalizers.

For a finite group G , let $\text{Ord}(\mathcal{S}_{sol}(G))$ be the set of orders of its solvable subgroups. The following conjecture was proposed by S. Abe and N. Iiyori [1]: Let G be a finite group and S be a non-abelian simple group. Then $G \cong S$ if and only if $\text{Ord}(\mathcal{S}_{sol}(G)) = \text{Ord}(\mathcal{S}_{sol}(S))$.

It was proved that if S is a simple group and G is a finite group such that $\text{Ord}(\mathcal{S}_{sol}(G)) = \text{Ord}(\mathcal{S}_{sol}(S))$, then $G \cong S$ or $\{G, S\} = \{B_n(q), C_n(q)\}$, where $n \geq 3$ and q is an odd prime power (see [3]). The purpose of this talk is to prove that the $\text{Ord}(\mathcal{S}_{sol}(B_n(q)))$ and $\text{Ord}(\mathcal{S}_{sol}(C_n(q)))$ are distinct.

2000 Mathematics Subject Classification: 20D06, 20D20, 20G40, 20C33.

keywords and phrases: Sylow subgroup, Classical groups, Simple group of Lie type, Characterization, Irreducible linear group, solvable subgroups, Hall subgroup, Fitting subgroup.

2. MAIN RESULTS

Theorem 2.1. *Let $S_{n,q} \in \{B_n(q), C_n(q)\}$ and let G be any finite group such that $n(S_{n,q}) = n(G)$. If $q \not\equiv \pm 1 \pmod{8}$, then G is isomorphic to $S_{n,q}$. Otherwise, G is isomorphic to $B_n(q)$ or $C_n(q)$.*

Theorem 2.2. *Let q be an odd prime power and $n \geq 3$. If $S \in \{B_n(q), C_n(q)\}$, then there are infinite pairs $\{(n, q)\}$ such that*

$$\text{Ord}(\mathcal{S}_{\text{sol}}(B_n(q))) \neq \text{Ord}(\mathcal{S}_{\text{sol}}(C_n(q))).$$

Theorem 2.3. *Let q be an odd prime power and $n \geq 3$. For the infinite pairs $\{(n, q)\}$, simple groups $B_n(q)$ and $C_n(q)$ are characterizable by orders of solvable subgroups.*

REFERENCES

- [1] S. Abe and N. Iiyori, *A generalization of prime graphs of finite groups*, Hokkaido Mathematical Journal **29** (2) (2000), 391-407.
- [2] J. Bi, *A characterization of $L_n(q)$ by the normalizers' orders of their Sylow subgroups*, Acta Math. Sinica (New Ser.) **11** (3) (1995), 300-306.
- [3] K. Denecke, X. Li and J. Bi, *A characterization of finite simple groups by the orders of solvable subgroups*, Science in China Series A: Mathematics **50** (5) (2007), 715-726.