

Tarbiat Moallem University, 20<sup>th</sup> Seminar on Algebra,  
2-3 Ordibehesht, 1388 (Apr. 22-23, 2009) pp 14-15

ON THE CHARACTERIZATION OF SIMPLE GROUPS  $B_n(q)$   
AND  $C_n(q)$

NEDA AHANJIDEH\* AND ALI IRANMANESH

Department of Mathematics  
Tarbiat Modares University  
P.O.Box: 14115-137, Tehran, Iran  
iranmana@modares.ac.ir

**ABSTRACT.** Let  $G$  be a finite group and  $\pi(G)$  be the set of prime divisors of the order of  $G$ . For  $t \in \pi(G)$  denote by  $n_t(G)$  the order of a normalizer of  $t$ - Sylow subgroup of  $G$  and put  $n(G) = \{n_t(G) \mid t \in \pi(G)\}$ . In this talk, we discuss about an answer to the following problem for the simple groups of Lie type  $B_n, C_n$ :

Let  $L$  be a finite non-abelian simple group and  $G$  be a finite group with  $n(L) = n(G)$ . Is it true that  $L \cong G$ ?

Also, we discuss about difference between orders of the solvable subgroups of the non-isomorphic simple groups  $B_n(q)$  and  $C_n(q)$ .

1. INTRODUCTION

Characterization by orders of Sylow normalizers has first been considered by Bi in 1992 (see [2]). It is known that if  $G$  is  $A_n(q)$ ,  ${}^2A_n(q)$ ,  $C_2(q)$ ,  ${}^2D_n(q)$ , alternating group, Mathieu simple groups, Janko groups and  $Sz(2^{2m+1})$ , then  $G$  is characterizable by orders of Sylow normalizers. In this talk, we discuss that if  $n = 2$  or  $q \not\equiv \pm 1 \pmod{8}$ , then  $B_n(q)$  and  $C_n(q)$  are characterizable by orders of Sylow normalizers and otherwise,  $B_n(q)$  and  $C_n(q)$  are 2-recognizable by orders of Sylow normalizers.

For a finite group  $G$ , let  $\text{Ord}(\mathcal{S}_{\text{sol}}(G))$  be the set of orders of its solvable subgroups. The following conjecture was proposed by S. Abe and N. Iiyori [1]: Let  $G$  be a finite group and  $S$  be a non-abelian simple group. Then  $G \cong S$  if and only if  $\text{Ord}(\mathcal{S}_{\text{sol}}(G)) = \text{Ord}(\mathcal{S}_{\text{sol}}(S))$ .

It was proved that if  $S$  is a simple group and  $G$  is a finite group such that  $\text{Ord}(\mathcal{S}_{\text{sol}}(G)) = \text{Ord}(\mathcal{S}_{\text{sol}}(S))$ , then  $G \cong S$  or  $\{G, S\} = \{B_n(q), C_n(q)\}$ , where  $n \geq 3$  and  $q$  is an odd prime power (see [3]). The purpose of this talk is to prove that the  $\text{Ord}(\mathcal{S}_{\text{sol}}(B_n(q)))$  and  $\text{Ord}(\mathcal{S}_{\text{sol}}(C_n(q)))$  are distinct.

---

**2000 Mathematics Subject Classification:** 20D06, 20D20, 20G40, 20C33.

**keywords and phrases:** Sylow subgroup, Classical groups, Simple group of Lie type, Characterization, Irreducible linear group, solvable subgroups, Hall subgroup, Fitting subgroup.

## 2. MAIN RESULTS

**Theorem 2.1.** *Let  $S_{n,q} \in \{B_n(q), C_n(q)\}$  and let  $G$  be any finite group such that  $n(S_{n,q}) = n(G)$ . If  $q \not\equiv \pm 1 \pmod{8}$ , then  $G$  is isomorphic to  $S_{n,q}$ . Otherwise,  $G$  is isomorphic to  $B_n(q)$  or  $C_n(q)$ .*

**Theorem 2.2.** *Let  $q$  be an odd prime power and  $n \geq 3$ . If  $S \in \{B_n(q), C_n(q)\}$ , then there are infinite pairs  $\{(n, q)\}$  such that*

$$\text{Ord}(S_{\text{sol}}(B_n(q))) \neq \text{Ord}(S_{\text{sol}}(C_n(q))).$$

**Theorem 2.3.** *Let  $q$  be an odd prime power and  $n \geq 3$ . For the infinite pairs  $\{(n, q)\}$ , simple groups  $B_n(q)$  and  $C_n(q)$  are characterizable by orders of solvable subgroups.*

## REFERENCES

- [1] S. Abe and N. Iiyori, *A generalization of prime graphs of finite groups*, Hokkaido Mathematical Journal **29** (2) (2000), 391-407.
- [2] J. Bi, *A characterization of  $L_n(q)$  by the normalizers' orders of their Sylow subgroups*, Acta Math. Sinica (New Ser.) **11** (3) (1995), 300-306.
- [3] K. Denecke, X. Li and J. Bi, *A characterization of finite simple groups by the orders of solvable subgroups*, Science in China Series A: Mathematics **50** (5) (2007), 715-726.