

Tarbiat Moallem University, 20th Seminar on Algebra,
2-3 Ordibehesht, 1388 (Apr. 22-23, 2009) pp 55-57

RELATIVE FLATNESS AND ASYMPTOTIC BEHAVIOUR

F. DOROSTKAR

Department of Mathematics
Faculty of Science
University of Guilan
dorostkar@guilan.ac.ir
(Joint work with H. Ansari-Toroghy)

ABSTRACT. Let R be a commutative ring and let I be an ideal of R . Let M be a Noetherian R -modules. Let F be an R -module which is flat relative to M . Suppose M' is a submodule of M and let $T = F \otimes_R -$. Then it is shown that the sequences of sets

$Ass_R(T(M)/I^n(T(M')))$ and $Ass_R(I^n(T(M))/I^n(T(M')))$, $n \in \mathbf{N}$
are ultimately constant.

1. INTRODUCTION

Throughout of this paper R will denote a commutative ring (with a nonzero identity), and \mathbf{N} is the set of all positive integers.

Let I be an ideal of R and M be a Noetherian R -module. If M' is a submodule of M it follows from [3] that both sequences of sets $Ass_R(M/I^n M')$ and $Ass_R(I^n M/I^n M')$, $n \in \mathbf{N}$ are ultimately constant.

Let R be a commutative Noetherian ring and let I be an ideal of R . Further assume that F is a flat R -module. Set $T = F \otimes_R -$. In [4], S. Yassemi showed that the $Ass_R(T(M))$ can be specified in terms of $Ass_R(M)$ and $Coass_R(F)$. Also it is shown that the sequences of sets

$$Ass_R(T(M/I^n M)) \text{ and } Ass_R(T(I^n M/I^{n+1} M)), n \in \mathbf{N}$$

are ultimately constant.

Now let M be an R -module and let the zero submodule of M have a primary decomposition. Let F be an R -module which is flat relative to M . Set $E = E(\bigoplus_{P \in Max(R)} R/P)$ and $F^\vee = Hom_R(F, E)$. In this paper we will generalize the results mentioned in last paragraph by showing that $M \otimes_R F$ has a primary decomposition and its weakly associated primes can be specified (see 2.1) in terms of $W.Ass_R(M)$ and $Ass_R(F^\vee)$ under an additional property. When R is a quasi-semi local ring, $W.Ass_R(M \otimes_R F)$ is specified (see 2.4) in

2000 Mathematics Subject Classification: 13D45, 13D07 .

keywords and phrases: Relative flatness, associated primes, weakly associated primes, attached primes.

terms of $W.Ass_R(M)$ and $Coass_R(F)$. Also when R is a commutative Noetherian ring, $Ass_R(M \otimes_R F)$ can be specified (see 2.3, 2.6) in terms of $Ass_R(M)$ and $Coass_R(F)$ without any restriction. Finally it is shown that if M is a Noetherian R -module then for every submodule M' of M , the sequences of sets

$$Ass_R((M \otimes_R F)/I^n(M' \otimes_R F)), n \in \mathbf{N}$$

and

$$Ass_R(I^n(M \otimes_R F)/I^n(M' \otimes_R F)), n \in \mathbf{N}$$

are ultimately constant (see 2.8). We recall that F is flat relative to M (or F is M -flat) if and only if for any submodule N of M , the homomorphism $F \otimes_R N \rightarrow F \otimes_R M$ is monic (see [1]).

2. MAIN RESULTS

Theorem 2.1. *Let M be an R -module and let F be an R -module which is flat relative to M . Further assume that the zero submodule of M has a primary decomposition and $Ass_R(F^\vee) = W.Ass_R(F^\vee)$, where $F^\vee = Hom_R(F, E)$ and $E = E(\bigoplus_{P \in Max(R)} R/P)$. Then $M \otimes_R F$ has a primary decomposition and we have*

$$W.Ass_R(M \otimes_R F) = \{P \in W.Ass_R(M) : P \subseteq Q \text{ for some } Q \in W.Ass_R(F^\vee)\}.$$

Corollary 2.2. *Let the situation be as in 2.1 and let F be flat R -module. Then we have*

$$W.Ass_R(M \otimes_R F) = \{P \in W.Ass_R(M) : P \subseteq Q \text{ for some } Q \in W.Ass_R(F^\vee)\}.$$

Corollary 2.3. *Let R be a commutative Noetherian ring and let M be an R -module. Let F be an R -module which is flat relative to M . Further assume that the zero submodule of M has a primary decomposition. Then $M \otimes_R F$ has a primary decomposition and we have*

$$Ass_R(M \otimes_R F) = \{P \in Ass_R(M) : P \subseteq Q \text{ for some } Q \in Coass_R(F)\}.$$

Corollary 2.4. *Let R be a semi- quasi local ring and let M be an R -module with the property that its zero submodule has a primary decomposition. Further assume that F is an R -module which is flat relative to M and that $W.Coass_R(F) = Coass_R(F)$ (This is true, for example, when each element of $W.cass_R(M)$ is finitely generated by [6], (2.4)). Then we have*

$$W.Ass_R(M \otimes_R F) = \{P \in W.Ass_R(M) : P \subseteq Q \text{ for some } Q \in Coass_R(F)\}.$$

Corollary 2.5. *Let M be an R -module and let F be an R -module which is flat relative to M . Further assume that the zero submodule of M has a primary decomposition and $Ass_R(F^\vee) = W.Ass_R(F^\vee)$, where $E = E(\bigoplus_{P \in Max(R)} R/P)$*

and $F^\vee = Hom_R(F, E)$. Then $(M \otimes_R F) \neq 0$ if and only if there exists $P \in W.Ass_R(M)$ such that $P \subseteq Q$ for some $Q \in Ass_R(F^\vee)$. Further,

$$W.Ass_R(M \otimes_R F) = \bigcup_{P \in W.Ass_R(M)} W.Ass_R(F/PF).$$

Theorem 2.6. *Let R be a commutative Noetherian ring and let M be an R -module. Let F be an R -module which is flat relative to M . Then we have*

$$\text{Ass}_R(M \otimes_R F) = \{P \in \text{Ass}_R(M) : P \subseteq Q \text{ for some } Q \in \text{Coass}_R(F)\}.$$

Corollary 2.7. *Let R be a commutative Noetherian ring let M be an R -module. let F be an R -module which is flat relative to M . Then we have*

$$\text{Ass}_R(M \otimes_R F) = \bigcup_{P \in \text{Ass}_R(M)} \text{Ass}_R(F/PF).$$

Theorem 2.8. *Let R be a commutative Noetherian ring and let M be a Noetherian R -module. Suppose F is an R -module which is flat relative to M . Let M' be a submodule of M and let $T = F \otimes_R -$. Then for an ideal I of R , the sequence of sets*

$$\text{Ass}_R(T(M)/I^n(T(M'))) \text{ and } \text{Ass}_R(I^n(T(M))/I^n(T(M'))), n \in \mathbf{N}$$

are ultimately constant. If we denote the ultimate constant value of the above sequences by T_1 and T_2 , then we have

$$T_1 = \{P \in \text{As}^*(I, M', M) : P \subseteq Q \text{ for some } Q \in \text{Coass}_R(F)\}$$

and

$$T_2 = \{P \in \text{Bs}^*(I, M', M) : P \subseteq Q \text{ for some } Q \in \text{Coass}_R(F)\}$$

Theorem 2.9. *Let M be a Noetherian R -module and suppose that F is an R -module which is flat relative to M . Let M' be a submodule of M and let $T = F \otimes_R -$. Then for an ideal I of R , the sequence of sets*

$$\text{Ass}_R(T(M)/I^n(T(M'))) \text{ and } \text{Ass}_R(I^n(T(M))/I^n(T(M'))), n \in \mathbf{N}$$

are ultimately constant.

REFERENCES

- [1] F.W. Anderson and Kent R. Fuller, *Rings and Categories of Modules* (second edition), Springer-Verlag New York, 1992.
- [2] H. Ansari-Toroghy, *Relative injectivity and secondary representation*, Southeast Asian Bulletin of Math., **28** (2004), 989-998.
- [3] D. E. Rush, *Asymptotic primes and integral closure in modules*, Quart.J. Math. Oxford (2) **43** (1992), 477-499.
- [4] S. Yassemi, *Coassociated primes*, Communication in Algebra, **23**(1995), 1473-1498.
- [5] S. Yassemi, *Coassociated primes of modules over commutative rings*, Math Scand. **80** (1997), 175-187.
- [6] S. Yassemi, *Weakly associated primes under change of rings*, communication in Algebra, (6) **26** (1998) , 2007-2018 .
- [7] S. Yassemi, *Weakly associated primes filtration*, Acta Math. Hunger **92** (2001), 179-183.