Tarbiat Moallem University, 20th Seminar on Algebra 2-3 Ordibehesht, 1388 (Apr. 22-23, 2009) pp 139-140

CENTRAL AUTOMORPHISMS OF SEMIDIRECT PRODUCT OF FINITE GROUPS

HAMID MOUSAVI AND AMIR SHOMALI

Department of Mathematics University of Tabriz Tabriz, Iran hmousavi@tabrizu.ac.ir a_shomali@tabrizu.ac.ir

ABSTRACT. In this paper we find the order and structure of $\operatorname{Aut}_c(G)$ for $G = K \rtimes H$ by imposing some conditions on H and K. As a result, we show that if M is a non-normal maximal subgroup of a solvable group G, such that all central automorphisms of G fix $Core_G(M)$ pointwise, then $\operatorname{Aut}_c(G)$, is meta-abelian.

1. INTRODUCTION

An automorphism σ of a group G is central if σ commutes with every automorphism in Inn(G), the group of inner automorphisms of G, or equivalently, if $g^{-1}\sigma(g)$ lies in the center Z(G) of G, for all g in G. The central automorphisms fix the commutator subgroup G' of G pointwise, and form a normal subgroup, denoted by $\text{Aut}_c(G)$, of the full automorphism group Aut(G). The group of central automorphisms of a finite group G is of great importance in the investigation of Aut(G), and has been studied by several authors (see, for example, [1-6]).

In [1] Adney and Yen has shown that if G is a finite purely non-abelian group then $|\operatorname{Aut}_c(G)| = |\operatorname{Hom}(G/G', Z(G))|$. Suppose $G = K \times H$ where K is purely non-abelian and H an abelian subgroup of G. In [5] Jamali and Jafari introduced some subgroups for $\operatorname{Aut}_c(G)$ to find order and structure of $\operatorname{Aut}_c(G)$.

In this paper we find the order and structure of $\operatorname{Aut}_c(G)$ for $G = K \rtimes H$ by imposing some conditions on H and K. As a result, we show that if M is a non-normal maximal subgroup of a solvable group G, such that all central automorphisms of G fix $Core_G(M)$ pointwise, then $\operatorname{Aut}_c(G)$, is meta-abelian.

In this talk, G is a finite group, π_H and π_K are the projection maps from G into H and K respectively and σ_H , σ_K are restriction of σ on H and K

²⁰⁰⁰ Mathematics Subject Classification: 20D15, 20D45.

keywords and phrases: Central Automorphism, Solvable group, Semidirect group.

respectively. Also we set:

$$\begin{split} R &= \{ \pi_{\scriptscriptstyle H} \sigma_{\scriptscriptstyle H} | \sigma \in \operatorname{Aut}_c(G) \} \quad S = \{ \pi_{\scriptscriptstyle K} \sigma_{\scriptscriptstyle K} | \sigma \in \operatorname{Aut}_c(G) \} \\ T &= \{ \pi_{\scriptscriptstyle K} \sigma_{\scriptscriptstyle H} | \sigma \in \operatorname{Aut}_c(G) \} \quad U = \{ \pi_{\scriptscriptstyle H} \sigma_{\scriptscriptstyle K} | \sigma \in \operatorname{Aut}_c(G) \} \end{split}$$

2. Main results

Theorem 2.1. Let $G = K \rtimes H$ be a semidirect product, such that (|H|, |K|) = 1. then $\operatorname{Aut}_c(G) \cong R \times S$. Furthermore, if Z(H) acts trivially on K then $\operatorname{Aut}_c(G) \cong \operatorname{Aut}_c(H) \times S$. In particular if $G = K \times H$, then $\operatorname{Aut}_c(G) \cong \operatorname{Aut}_c(K)$.

Corollary 2.2. Let G be a solvable group. If M is a non-normal maximal subgroup of index p^t such that $p \nmid |\frac{M}{C}|$ and (|M/C|, |C|) = 1, where $C = Core_G(M)$ then we can write $G = K \rtimes H$, for which (|K|, |H|) = 1 and $H \leq M$. Hence $\operatorname{Aut}_c(G) \cong R \times S$.

Theorem 2.3. Let $\frac{G}{N} = \frac{K}{N} \rtimes \frac{H}{N}$ be a semidirect product and the central automorphisms fix the subgroup N of G pointwise and $(|\frac{H}{N}|, |\frac{K}{N}|) = 1$. then $\operatorname{Aut}_c(G) \cong R \times S$.

Theorem 2.4. Let $G = K \rtimes H$ be a semidirect product, such that $C_K(H) = 1$. Then $\operatorname{Aut}_c(G) \cong R \rtimes U$.

Theorem 2.5. Let $\frac{G}{N} = \frac{K}{N} \rtimes \frac{H}{N}$ be a semidirect product and the central automorphisms fix the subgroup N of G pointwise and $\mathcal{C}_{K/N}(H/N) = 1$. then $\operatorname{Aut}_c(G) \cong R \rtimes U'$ where $U' = \{f_\sigma \mid f_\sigma \in \operatorname{Hom}(K, Z(G) \cap H), \sigma \in \operatorname{Aut}_c(G)\}.$

Corollary 2.6. Let G be a solvable group with non-normal maximal subgroup M. If all central automorphisms fix $Core_G(M)$ pointwise, then $Aut_c(G) \cong R$ and so is abelian.

Theorem 2.7. Let $G = K \rtimes H$ be a semidirect product, where K is purely non-abelian and Z(H) acts trivially on K. then $\operatorname{Aut}_c(G) \cong RTUS$.

References

- J. E. Adney and T.Yen, Automorphisms of a p-groups, Illinois J. math. 9 (1965), 137-143.
- [2] M. J. Curran and D. J. McCaughan, Central automorphisms that are almost inner, Comm. Algebra, 29(5) (2001), 2081-2087.
- [3] B. Huppert, Endliche Gruppen, Vol. 1, Springer-Verlag, (1967).
- M. H. Jafari, Elementary abelian p-group as central automorphism groups, Comm. Algebra, 34 (2006), 601-607.
- [5] M. H. Jafari and A. R. Jamali, On the occurrence of some finite groups in the central automorphism group of finite groups, *Math. Proc. Royal Irish Acad.*, **106A(2)** (2006), 139-148.
- [6] A. R. Jamali and H. Mousavi, On the central automorphism groups of finite p-groups, Algebra Colloq., 9(1) (2002), 7-14.

140