

ID :	10724
Themes :	قلب و عروق
Title :	Comparison of beta oxidation of fatty acid enzymes activity in diabetic and normal rat hearts and effect of ranolazine
Authors :	1-Hajar Khazraei, 2-Masome Akmali, 3-Hossein Mirkhani
Address :	1,3-Zand street, Faculty of medicine, Department of pharmacology, Shiraz, Iran 2-Zand street, Faculty of medicine, Department of Biochemistry, Shiraz, Iran Khazraee@sums.ac.ir
Abstract :	Introduction: Clinical and experimental evidence suggests that diabetes increased rates of fatty acid oxidation in the myocardium. Ranolazine is a new antianginal drug that affects as an inhibitor of fatty acid beta-oxidation in hearts. The objective of this study was to examine the effect of different concentrations of ranolazine on cardiac fatty acid oxidation in streptozocin (STZ)-diabetic rats. Method : After 8 week of STZ-injection, rat hearts were isolated and homogenated. The activities of 3-hydroxy acyl CoA dehydrogenase, hydratase and thiolase related to trifunctional protein were determined by spectrophotometric methods in rat hearts. Results: The activity of 3-hydroxy acyl CoA dehydrogenase, and hydratase was increased but 3-ketoacyl-CoA thiolase activity is decreased in diabetic hearts compared to normal ones. Ranolazine (10 µmol/L) partially inhibited enzyme complex activity such as hydratase and 3-ketoacyl-CoA thiolase in rat heart.But there is not any significant effect on 3-hydroxy acyl CoA dehydrogenase. Conclusion: Ranolazine in therapeutic concentration (10 micro molar) has partial inhibitory effects on fatty acid beta-oxidation.
Keywords :	ranolazine, cardiac fatty acid oxidation, diabetes, 3-KAT