Talk



Higher-order asymptotic formula for the eigenvalues of Sturm-Liouville ...

Higher-order asymptotic formula for the eigenvalues of Sturm-Liouville problem with indefinite weight function in the case of y(a) = y'(b) = 0, y'(a) = y(b) = 0

F. Dastmalchi Saei

Department of Mathematics, Tabriz Branch, Islamic Azad University, Tabriz, Iran

Abstract

In this paper, we investigate the asymptotic behavior of the differential equation

 $y'' + (\lambda r(x) - q(x))y = 0,$ $0 \le x \le 1,$

where [0, 1] contains a finite number of zeros of r(x), λ is a real parameter and the function q(x) is bounded and integrable in [0, 1]. Using a technique used previously in [7], we derive the higher-order asymptotic distribution of the positive eigenvalues associated with this equation for the SLP (in the boundary condition of, y(0) = y'(1) = 0 or y'(0) = y(1) = 0).

Keywords: Sturm-Liouville problem, eigenvalues, asymptotic

Mathematics Subject Classification: 34L20

1 Introduction

We study the indefinite Sturm-Liouville spectral problem

$$y'' + (\lambda \ r(x) - q(x))y = 0, \qquad a \le x \le b$$
(1)

$$y'(a) = y(b) = 0 \lor y(a) = y'(b) = 0$$

defined on the interval [a, b] where λ is a real parameter, r(x), q(x) are real and integrable on [a, b]; moreover,

$$\int_{a}^{b} \sqrt{r_{+}(t)} dt > 0, \quad where \quad r_{+}(x) = max\{r(x), 0\}.$$
⁽²⁾

It follows from [2] that the spectrum of this problem is discrete and has no finite accumulation points; moreover, only finitely many eigenvalues lie the outside the real and imaginary axes. In what follows, we shall assume that λ is a positive parameter. In [2] it was shown that the asymptotics of the eigenvalues is of the form

$$\lambda_n \sim \frac{n\pi}{\int_a^b \sqrt{r_+(t)} dt}.$$

Our goal is to refine the asymptotics under the additional assumptions of smoothness of the functions r(x) and q(x). In addition, we assume that r(x) has a finite number of zeros, which are called

Talk



k Higher-order asymptotic formula for the eigenvalues of Sturm-Liouville ...

turning points.

The outline of our paper is as follows. First, we find the asymptotics of eigenvalues for one turning point. Next, using a technique previously in [7], we derive the higher order asymptotic distribution of the positive eigenvalues in the case of two turning points. Finally, in the case of an arbitrary finite number of turning points it can be reduced to the two cases discussed above.

2. The case of one turning point

First, consider the case

$$r(x) = (x - x_{\nu})^{l_{\nu}} h(x), \qquad h(x) > 0.$$

To simplify the formulas, we assume that x varies on the closed interval with endpoints a and b, where r(a) < 0 and r(b) > 0. The turning point x_{ν} lies between a and b. We distinguish four different types of turning points:

$$T_{\nu} = \begin{cases} I & \text{if } l_{\nu} \text{ is even and } r(x) < 0 \text{ in } [a, b] \\ II & \text{if } l_{\nu} \text{ is even and } r(x) > 0 \text{ in } [a, b] \\ III & \text{if } l_{\nu} \text{ is odd and } r(x) < 0 \text{ in } [x_{\nu}, b] \\ IV & \text{if } l_{\nu} \text{ is odd and } r(x) > 0 \text{ in } [x_{\nu}, b] \end{cases}$$

is called of type x_{ν} . By Langer's transformation we can make zero of r(x) the origin. To be specific, let us define the Langer's transformation $\xi(x)$ for different type of TP. For a turning point of Type I :

$$\xi_I(x) = \begin{cases} -\{\int_x^{x_\nu} (-r)^{1/2}(t)dt\}^{\frac{2}{\ell+2}} & x \le x_\nu \\ -\{\int_{x_\nu}^x (-r(t))^{1/2}dt\}^{\frac{2}{\ell+2}} & x_\nu \le x \end{cases}$$

For a turning point of Type II :

$$\xi_{II}(x) = \begin{cases} \left\{ \int_{x}^{x_{\nu}} r^{1/2}(t) dt \right\}^{\frac{2}{\ell+2}} & x \le x_{\nu} \\ \left\{ \int_{x_{\nu}}^{x} r(t)^{1/2} dt \right\}^{\frac{2}{\ell+2}} & x_{\nu} \le x. \end{cases}$$

For a turning point of Type III :

$$\xi_{III}(x) = \begin{cases} \{\int_x^{x_\nu} r^{1/2}(t)dt\}^{\frac{2}{\ell+2}} & x \le x_\nu\\ -\{\int_{x_\nu}^x (-r(t))^{1/2}dt\}^{\frac{2}{\ell+2}} & x_\nu \le x \end{cases}$$

For a turning point of Type IV :

$$\xi_{IV}(x) = \begin{cases} -\{\int_x^{x_{\nu}} (-r(t))^{1/2} dt\}^{\frac{2}{\ell+2}} & x \le x_{\nu} \\ \{\int_{x_{\nu}}^x (r(t))^{1/2} dt\}^{\frac{2}{\ell+2}} & x_{\nu} \le x. \end{cases}$$

From [9] we rewrite showing the connection between the argument of complex valued solution of (14) in the interval containing one of the turning point say, x_{ν} , and the argument of complex valued solution of Sturm-Liouville equation with one turning point in x = 0 in the same interval. In fact the following result illustrates a crucial relationship between a general problem (14) with a turning point at x_{ν} to a transformed problem in which is mapped to x = 0. We show that such a transformation preserves the argument of any fixed complex valued solution.

Theorem 1.1. Let z be a strictly complex -valued solution of the differential equations

$$y'' + (\rho^2 r(x) - q(x))y = 0, \qquad x \in [0, 1]$$
(3)

and W be a solution of

$$W^{''} + (u^2(-1)^{M_\nu}\xi^{\ell_\nu} - R_\nu(\xi))W = 0, \qquad \xi \in [c,d]$$
(4)



562-570

then on the interval $[x_{\nu} - \epsilon, x_{\nu} + \epsilon]$

$$argW(\xi(x)) = \arg z,$$

where $r(x) = \prod_{j=1}^{n} (x - x_j)^{l_j} \phi_0(x)$ and

$$R_{\nu}(\xi) = \left(\frac{dx}{d\xi}\right)^{1/2} \frac{d^2}{d\xi^2} \left\{\frac{1}{\left(\frac{dx}{d\xi}\right)^{1/2}}\right\} + \left(\frac{dx}{d\xi}\right)^2 q(x(\xi)).$$

 M_k = the number of turning of type (III) or (IV) in $(x_k, 1)$, or one can see that $(-1)^{M_k} = (-1)^{l_n+\dots+l_{k-1}}$, c < 0 < d, $u^2 = \frac{(\ell_{\nu}+2)^2}{4}\rho^2$, the transformation $\xi(x)$ is Langer's transformation.

Proof: For proof see [9].

3. The main result

We begin by consolidating some results from [5,9] for completeness. For a complex-valued solution $\Omega(x,\lambda)$, of

$$y'' + \lambda x^{\alpha} y = 0, \qquad (E_0$$

we form the logarithmic derivative $r_0(x,\lambda) = \Omega'(x,\lambda)/\Omega(x,\lambda)$, a quantity that exists for each $x \in [a,b]$ since the real and imaginary parts of Ω are linearly independent solution of (E_0) . The quantity $r_1(x,\lambda)$ is defined by setting

$$r_1(x,\lambda) = -\int_x^b q(t)e^{2\int_x^t r_0(s,\lambda)ds}dt,$$

while the $r_n(x, \lambda)$ are defined recursively (for $n \ge 1$) by

$$r_{n+1}(x,\lambda) = \int_x^b r_n^2(x,\lambda) \exp(2\sum_{l=0}^n \int_x^t r_l(s,\lambda) ds) dt$$

It follows (cf.[4]) that the function

$$r(x,\lambda) = \sum_{n=0}^{\infty} r_n(x,\lambda) := S(x,\lambda) + iT(x,\lambda)$$

is a series solution (in x) of the Riccati equation

$$\upsilon' = q - \lambda x^{\alpha} - \upsilon^2$$

from which one can reconstruct solutions of (14) with Neumann condition y'(a) = y'(b) = 0 via the following result:

Theorem 1.2. (see Harris-Talarico[4]) There exists λ_0 such that any real valued solution of

$$y'' + (\lambda x^{\alpha} - q(x))y = 0 \tag{5}$$

 $can \ be \ expressed \ as$:

$$Z(x,\lambda) = c_1 e^{\int_a^x S(t,\lambda)dt} \cos(c_2 + \int_a^x T(t,\lambda)dt)$$

for $x \in [a, b]$ (a < 0 < b) and $|\lambda| \ge \lambda_0$ where $c_1, c_2 \in \Re$. If $Z(., \lambda)$ satisfies

$$y(a)\cos\gamma + y'(a)\sin\gamma = 0 \tag{6}$$

then

$$c_{2} =: c_{2}^{a} = \frac{\pi}{2} \qquad \qquad if \quad \gamma = 0$$
$$= \arctan(\frac{1}{T(a,\lambda)}S(a,\lambda) + \cot\gamma) \qquad \qquad if\gamma \neq 0 \qquad (7)$$

www.SID.ir



Talk Higher-order asymptotic formula for the eigenvalues of Sturm-Liouville ... pp. 562-570

 $Similarly, if Z \ satisfies$

$$y(b)\cos\beta + y'(b)\sin\beta = 0 \tag{8}$$

then

$$c_{2} =: c_{2}^{b} = n\pi + \frac{\pi}{2} \qquad \qquad if \quad \beta = 0$$
$$= n\pi + \arctan(\frac{1}{T(b,\lambda)}S(b,\lambda) + \cot\beta) \qquad if \quad \beta \neq 0 \qquad \qquad (9)$$

for all integer n.

It follows from (7) and (9) that the eigenvalues of (7),(6) and (8), i.e., our problem (14), are the values of λ for which

$$c_{2}^{a} + \int_{a}^{b} T(t,\lambda)dt = c_{2}^{b}$$
 (10)

We see from [3], that the asymptotic distribution of the eigenvalues of (7), (6) and (8) is therefore determined by the following transcendental equation:

$$n\pi + \arctan(\frac{1}{T(b,\lambda)}S(b,\lambda) + \cot\beta) = \int_{a}^{b} T(t,\lambda)dt + \arctan(\frac{1}{T(a,\lambda)}S(a,\lambda) + \cot\gamma)$$

$$= \Im \int_{a}^{b} r(t,\lambda)dt + \arctan(\frac{1}{T(b,\lambda)}S(b,\lambda) + \cot\gamma)$$

$$= \Im(\int_{a}^{b} r_{0}(t,\lambda)dt + \int_{a}^{b} r_{1}(t,\lambda)dt + \cdots)$$

$$+ \arctan(\frac{1}{T(a,\lambda)}S(a,\lambda) + \cot\gamma)$$
(11)
$$= \arg\Omega(b,\lambda) - \arg\Omega(a,\lambda) - \frac{\pi}{2k}\int_{0}^{b} xq(x)J_{\nu}^{2}(k^{-1}\lambda^{1/2}x^{k})dx + \cdots$$

$$+ \arctan(\frac{1}{T(a,\lambda)}S(a,\lambda) + \cot\gamma).$$

Note that we use the following result from [8],

$$\Im \int_{a}^{b} r(t,\lambda)dt = \arg \Omega(b,\lambda) - \arg \Omega(a,\lambda) - \frac{\pi}{2k} \int_{0}^{b} xq(x) J_{\nu}^{2}(k^{-1}\lambda^{1/2}x^{k})dx$$

By applying the above relation to approximate eigenvalues in the case of $\gamma = 0, \beta = \frac{\pi}{2}$ or $\gamma = \frac{\pi}{2}, \beta = 0.$

Theorem 1.3. Consider the differential equation (1) on [a, b] under condition (2). Then the positive eigenvalues admit the following asymptotic representation: Case1: $\gamma = 0, \beta = \frac{\pi}{2}$ (a) Let x_{ν} be of type IV. Then

$$\sqrt{\lambda_n} = \frac{n\pi - \frac{\pi}{4}}{\int_a^b \sqrt{r_+(t)}dt} - \frac{1}{n\pi} \left(\frac{4(\nu - 1)^2 - 1}{8\int_a^b \sqrt{r_+(t)}dt} - \frac{1}{2}H(b)\right) + O\left(\frac{1}{n^2}\right)$$

where

$$H(b) = \int_{x_{\nu}}^{b} \left(\frac{q(x)}{\tilde{r}(x)} - \frac{1}{\tilde{r}^{3/4}} \frac{d^{2}}{dx^{2}} (\tilde{r}^{-1/4})\right) \frac{\tilde{r}}{r^{\frac{1}{2}}} dx,$$



and

$$\tilde{r} = (\frac{d\xi}{dx})^2 = \frac{4r(x)}{(l+2)^2(\xi(x))^l}.$$

(b) Let x_{ν} be of type III. Then

$$\sqrt{\lambda_n} = \frac{n\pi + \frac{\pi}{4}}{\int_a^b \sqrt{r_+(t)}dt} - \frac{1}{n\pi} (\frac{4(\nu-1)^2 - 1}{8\int_a^b \sqrt{r_+(t)}dt} - \frac{1}{2}H(a)) + O(\frac{1}{n^2})$$

where

$$H(a) = \int_{a}^{x_{\nu}} \left(\frac{q(x)}{\tilde{r}(x)} - \frac{1}{\tilde{r}^{3/4}} \frac{d^{2}}{dx^{2}} (\tilde{r}^{-1/4})\right) \frac{\tilde{r}}{r^{\frac{1}{2}}} dx,$$

and

$$\tilde{r} = \left(\frac{d\xi}{dx}\right)^2 = \frac{4r(x)}{(l+2)^2(\xi(x))^l}.$$

(c) Let x_{ν} be of type II. Then

$$\sqrt{\lambda_n} = \frac{n\pi}{\int_a^b \sqrt{r(t)}dt} - \frac{1}{n\pi} \{ \frac{4(\nu-1)^2 - 1}{8\int_a^{x_\nu} \sqrt{r(t)}dt} + \{ \frac{4(\nu-1)^2 - 1}{8\int_{x_\nu}^b \sqrt{r(t)}dt} - \frac{1}{2}H(a) - \frac{1}{2}H(b) \} + O(\frac{1}{n^2})$$

where H(a) and H(b) are defined above. Case1: $\gamma = 0, \beta = \frac{\pi}{2}$ (a) Let x_{ν} be of type IV. Then

$$\sqrt{\lambda_n} = \frac{n\pi - \frac{\pi}{4}}{\int_a^b \sqrt{r_+(t)}dt} - \frac{1}{n\pi} \left(\frac{4(\nu - 1)^2 - 1}{8\int_a^b \sqrt{r_+(t)}dt} - \frac{1}{2}H(b)\right) + O(\frac{1}{n^2})$$

where

$$H(b) = \int_{x_{\nu}}^{b} \left(\frac{q(x)}{\tilde{r}(x)} - \frac{1}{\tilde{r}^{3/4}} \frac{d^{2}}{dx^{2}} (\tilde{r}^{-1/4})\right) \frac{\tilde{r}}{r^{\frac{1}{2}}} dx,$$

and

$$\tilde{r} = (\frac{d\xi}{dx})^2 = \frac{4r(x)}{(l+2)^2(\xi(x))^l}.$$

(b) Let x_{ν} be of type III. Then

$$\sqrt{\lambda_n} = \frac{n\pi + \frac{\pi}{4}}{\int_a^b \sqrt{r_+(t)}dt} - \frac{1}{n\pi} \left(\frac{4(\nu-1)^2 - 1}{8\int_a^b \sqrt{r_+(t)}dt} - \frac{1}{2}H(a)\right) + O(\frac{1}{n^2})$$

where

$$H(a) = \int_{a}^{x_{\nu}} \left(\frac{q(x)}{\tilde{r}(x)} - \frac{1}{\tilde{r}^{3/4}} \frac{d^{2}}{dx^{2}} (\tilde{r}^{-1/4})\right) \frac{\tilde{r}}{r^{\frac{1}{2}}} dx,$$

and

$$\tilde{r} = \left(\frac{d\xi}{dx}\right)^2 = \frac{4r(x)}{(l+2)^2(\xi(x))^l}.$$

www.SID.ir



Talk

Higher-order asymptotic formula for the eigenvalues of Sturm-Liouville ... pp.

(c) Let x_{ν} be of type II. Then

$$\sqrt{\lambda_n} = \frac{n\pi}{\int_a^b \sqrt{r(t)}dt} - \frac{1}{n\pi} \{ \frac{4(\nu-1)^2 - 1}{8\int_a^{x_\nu} \sqrt{r(t)}dt} + \{ \frac{4(\nu-1)^2 - 1}{8\int_{x_\nu}^b \sqrt{r(t)}dt} - \frac{1}{2}H(a) - \frac{1}{2}H(b) \} + O(\frac{1}{n^2}) \}$$

where H(a) and H(b) are defined above.

Proof: Let us proof only case (1*a*), from [14] for $\alpha = 4n \pm 1$ and x < 0 we know that $\Omega(x, \lambda) = e^{i\pi/2} |x|^{1/2} H_{\nu}^{(1)}(e^{ik\pi}\rho |x|^k)$, and from [1, (9.2.7)], Hankel's expansion the form

$$H_{\nu}^{(1)}(z) = \sqrt{\frac{2}{\pi z}} e^{i(z - \frac{1}{2}\nu\pi - \frac{\pi}{4})} M(z), \quad (-\pi < \arg z < \pi),$$

where M(z) is bounded away from the origin and $M(z) \to 1$ as $|z| \to \infty$ in the sector.

Now since $\Omega(x,\lambda) = i\rho^{-\nu}e^{-ik\pi\nu}(e^{ik\pi}\rho|x|^k)^{\nu}H_{\nu}^{(1)}(e^{ik\pi}\rho|x|^k)$, where $k = \frac{\alpha+2}{2}, \nu = \frac{1}{\alpha+2}$ and $k\nu = \frac{1}{2}$, by using ([1],(9.1.30)) we find $\Omega'(x,\lambda) = ike^{ik\pi}\rho|x|^{k-3/2}xH_{\nu-1}^{(1)}(e^{ik\pi}\rho|x|^k)$. Therefore

$$\begin{split} &\frac{\Omega'(x,\lambda)}{\Omega(x,\lambda)} = k\rho e^{ik\pi}x \mid x \mid^{k-2} \frac{H_{\nu-1}^{(1)}(z)}{H_{\nu}^{(1)}(z)} \\ &= k\rho e^{ik\pi}x |x|^{k-2}(i + \frac{8\nu - 4}{8z} + O(\frac{1}{z^2})), \qquad z = e^{ik\pi}\rho |x|^k. \end{split}$$

Then

$$\Re \frac{\Omega'(x,\lambda)}{\Omega(x,\lambda)} = k\rho x |x|^{k-2} - k \frac{8\nu - 4}{8|x|} + O(\frac{1}{z^2}).$$

and

$$\Im \frac{\Omega'(x,\lambda)}{\Omega(x,\lambda)} = 0.$$

On the other hand,

$$\frac{S(a,\lambda)}{T(a,\lambda)} = \frac{\Re r_0(a,\lambda) + \Re r_1(a,\lambda) + \cdots}{\Im r_0(a,\lambda) + \Im r_1(a,\lambda) + \cdots} = \frac{\Re \frac{\Omega'(a,\lambda)}{\Omega(a,\lambda)} + \Re\{-\int_a^b q(t)e^{2\int_a^t r_0(s,\lambda)ds}dt\} + \cdots}{\Im \frac{\Omega'(a,\lambda)}{\Omega(a,\lambda)} + \Im\{-\int_a^b q(t)e^{2\int_a^t r_0(s,\lambda)ds}dt\} + \cdots}$$

Now we have:

$$\begin{split} \int_{a}^{b} q(t)e^{2\int_{a}^{t}r_{0}(s,\lambda)ds}dt &= \int_{a}^{0}q(t)e^{2\int_{a}^{t}r_{0}(s,\lambda)ds}dt + \int_{0}^{b}q(t)e^{2\int_{a}^{0}r_{0}(s,\lambda)ds+2\int_{0}^{t}r_{0}(s,\lambda)ds}dt \\ &= \frac{1}{|\Omega(a,\lambda)|^{2}}\int_{a}^{0}q(t)|\Omega(t,\lambda)|^{2}e^{2i\{\arg\Omega(t,\lambda)-\arg\Omega(a,\lambda)\}}dt \\ &+ \int_{0}^{b}q(t)|\frac{\Omega(0,\lambda)}{\Omega(a,\lambda)}|^{2}e^{2i\{\arg\Omega(0,\lambda)-\arg\Omega(a,\lambda)\}} \times |\frac{\Omega(t,\lambda)}{\Omega(0,\lambda)}|^{2}e^{2i\{\arg\Omega(t,\lambda)-\arg\Omega(0,\lambda)\}}dt \\ &= \frac{e^{-2i\arg\Omega(a,\lambda)}}{|\Omega(a,\lambda)|^{2}}\{\int_{a}^{0}q(t)|\Omega(t,\lambda)|^{2}e^{2i\arg\Omega(t,\lambda)}dt + \int_{0}^{b}q(t)|\Omega(t,\lambda)|^{2}e^{2i\arg\Omega(t,\lambda)}dt\} = O(1) \end{split}$$

Similarly for $\alpha = 4n \mp 1$ and x > 0 we know that $\Omega(x, \lambda) = x^{1/2} H_{\nu}^{(1)}(\rho x^k)$. Now since $\Omega(x, \lambda) = \rho^{-\nu}(\rho x^k)^{\nu} H_{\nu}^{(1)}(\rho x^k)$, by using ([1],(9.1.30)) we find $\Omega'(x, \lambda) = kx^{k-1/2} \rho H_{\nu-1}^{(1)}(\rho x^k)$. Therefore

$$\frac{\Omega'(x,\lambda)}{\Omega(x,\lambda)} = k\rho x^{k-1} \frac{H_{\nu-1}^{(1)}(z)}{H_{\nu}^{(1)}(z)} = k\rho x^{k-1} (i + \frac{8\nu - 4}{8z} + O(\frac{1}{z^2})).$$



Talk Higher-order asymptotic formula for the eigenvalues of Sturm-Liouville ... pp.

Then

$$\Re \frac{\Omega'(x,\lambda)}{\Omega(x,\lambda)} = k\rho x^{k-1} \frac{8\nu - 4}{8\rho x^k} = k \frac{8\nu - 4}{8x} + O(\frac{1}{z^2}),$$

and

$$\Im \frac{\Omega'(x,\lambda)}{\Omega(x,\lambda)} = k\rho x^{k-1} + O(\frac{1}{z^2}),$$

whence

$$\Re \frac{\Omega'(b,\lambda)}{\Omega(b,\lambda)} = k \frac{8\nu-4}{8b} + O(\frac{1}{z^2}),$$

and

$$\Im \frac{\Omega'(b,\lambda)}{\Omega(b,\lambda)} = k\rho b^{k-1} = \sqrt{\lambda} b^{k-1} + O(\frac{1}{z^2}),$$

also

$$\Re r_1(b,\lambda) = \Im r_1(b,\lambda) = 0.$$

By substitution in (11) and using :

$$\arctan x = x - \frac{x^3}{3} + \cdots,$$

we have (the particulars of the calculations are omitted):

$$\sqrt{\lambda_n} = \frac{n\pi - \frac{\pi}{4}}{\int_a^b \sqrt{r_+(t)}dt} - \frac{1}{n\pi} (\frac{4(\nu-1)^2 - 1}{8\int_a^b \sqrt{r(t)}dt} - \frac{1}{2}H(b)) + O(\frac{1}{n^2}).$$

2 Main Result

The cases of two and n turning points

From now then, without losing generalization, we suppose that the coefficients q(x) and r(x) satisfy:

(i) r(x) is real and has in [0, 1] n zeros x_ν of order $l_\nu \in \mathbb{N}$,1 $\leq \nu \leq n$ where $0 < x_1 < x_2 < \ldots < x_n < 1$.

(ii) The function $\phi_0: I \to R - \{0\}, x \to r(x) \prod_{\nu=1}^n (x - x_\nu)^{-l_\nu}$ is twice continuously differentiable. (iii) q(x) is bounded and integrable in I.

We shall use the symbol $\Omega_{IV}(\xi, u)$ to signify the complex-valued solution of

$$W^{''} + u^2 \xi^{l_v} W = 0,$$

where ξ is corresponding Langer's transformation of turning point of type IV. We will use the symbols $\Omega_I(\xi, u), \Omega_{II}(\xi, u)$ and $\Omega_{III}(\xi, u)$ in similar case.

Now we can derive the following results on the distribution of the eigenvalues of (14) with Neumann boundary condition .

We consider only the following case:

1.a $T_1 = IV, T_2 = III.$

We suppose that the weight function r(x) has in [0,1] two zeros x_1 and x_2 where x_1 of type *IV* and x_3 of type *III*. By (11) the distribution of positive eigenvalue satisfies :



562-570

 $\operatorname{pp.}$

$$n\pi = \Im \int_{0}^{1} \frac{y'}{y} dx = \Im \left(\int_{0}^{\alpha_{12}} \frac{y'}{y} dx + \int_{\alpha_{12}}^{1} \frac{y'}{y} dx \right) + \arctan\left(\frac{S(0,\lambda)}{T(0,\lambda)}\right) - \arctan\left(\frac{S(1,\lambda)}{T(1,\lambda)}\right)$$

$$= \rho \int_{0}^{1} \sqrt{r_{+}(t)} dt + \frac{\pi}{2} - \frac{1}{4\rho} \left(\frac{8\nu_{1} - 4}{\int_{x_{1}}^{x_{2}} \sqrt{r(t)} dt} + \frac{8\nu_{2} - 4}{\int_{x_{1}}^{x_{2}} \sqrt{r(t)} dt} \right) + \frac{1}{4\rho} \left(\frac{4\nu_{1}^{2} - 1}{\int_{x_{1}}^{x_{2}} \sqrt{r(t)} dt} + \frac{4\nu_{2}^{2} - 1}{\int_{x_{1}}^{x_{2}} \sqrt{r(t)} dt} \right)$$

$$- \frac{1}{2u} P(x_{1}, x_{2}) + O\left(\frac{1}{u^{2}}\right), \qquad (12)$$

where $\alpha_{12} \in (x_1, x_2)$ is such that $\int_{x_1}^{\alpha_{12}} \sqrt{r(t)} dt = \int_{\alpha_{12}}^{x_2} \sqrt{r(t)} dt$ (the existence of α_{12} follows by Intermediate Value Theorem), and

$$P(x_1, x_2) = P_{IV}(x_1, \alpha_{12}) + P_{III}(x_2, \alpha_{12}) = \int_{x_1}^{\alpha_{12}} E_{IV}(x) dx + \int_{\alpha_{12}}^{x_2} E_{III}(x) dx,$$

$$P_{IV}(x_1,\alpha_{12}) = \int_0^{d_{12}} \frac{R_{IV}(\xi)}{\xi^{\frac{1}{2}}} d\xi = \int_{x_1}^{\alpha_{12}} (\frac{q(x)}{\tilde{r}(x)} - \frac{1}{\tilde{r}^{3/4}} \frac{d^2}{dx^2} (\tilde{r}^{-1/4})) \frac{\tilde{r}}{r^{\frac{1}{2}}} dx := \int_{x_1}^{\alpha_{12}} E_{IV}(x) dx,$$

and

$$\tilde{r} = \left(\frac{d\xi}{dx}\right)^2 = \frac{4r(x)}{(l+2)^2(\xi(x))^l}, \xi(\alpha_{12}) = d_{12}, \xi(x_1) = 0, \xi(0) = c_{12}$$
$$P_{III}(x_2, \alpha_{12}) = \int_{\alpha_{12}}^{x_2} \left(\frac{q(x)}{\tilde{r}(x)} - \frac{1}{\tilde{r}^{3/4}} \frac{d^2}{dx^2} (\tilde{r}^{-1/4})\right) \frac{\tilde{r}}{r^{\frac{1}{2}}} dx := \int_{\alpha_{12}}^{x_2} E_{III}(x) dx.$$

By inversion, we get

$$\rho_n = \frac{n\pi - \frac{\pi}{2}}{\int_0^1 \sqrt{r_+(t)}dt} - \frac{1}{n\pi} \{ \frac{[4(\nu_1 - 1)^2 - 1] + [4(\nu_2 - 1)^2 - 1]}{4\int_{x_1}^{x_2} \sqrt{r(t)}dt} - \frac{1}{2}P(x_1, x_2) \} + O(\frac{1}{n^2}).$$

By applying the same method and using theorem (1),(2) we get:

$$\begin{split} n\pi &= \rho \int_{0}^{1} \sqrt{r_{+}(t)} dt - \frac{(n-1)\pi}{2} + \frac{n\pi}{2} + \frac{1}{4\rho} (\frac{4(\nu_{1}-1)^{2}-1}{\int_{x_{1}}^{x_{2}} \sqrt{r(t)} dt} + \frac{4(\nu_{2}-1)^{2}-1}{\int_{x_{2}}^{x_{3}} \sqrt{r(t)} dt} + \frac{4(\nu_{2}-1)^{2}-1}{\int_{x_{2}}^{x_{3}} \sqrt{r(t)} dt} + \frac{4(\nu_{3}-1)^{2}-1}{\int_{x_{3}}^{x_{4}} \sqrt{r(t)} dt} + \cdots + \frac{4(\nu_{n-1}-1)^{2}-1}{\int_{x_{n-2}}^{x_{n-1}} \sqrt{r(t)} dt} + \frac{4(\nu_{n-1}-1)^{2}-1}{\int_{x_{n-1}}^{x_{n}} \sqrt{r(t)} dt} \\ &+ \frac{4\nu_{n}^{2}-1}{\int_{x_{n-1}}^{x_{n}} \sqrt{r(t)} dt}) - \frac{1}{2u} P(x_{1}, x_{2}, \cdots, x_{n}) + O(\frac{1}{u^{2}}), \end{split}$$

where

$$P(x_1, x_2, \cdots, x_n) = \int_{x_1}^{\alpha_{12}} E_{IV}(x) dx + \sum_{i=1}^{n-2} \int_{\alpha_{i(i+1)}}^{x_{i+1}} E_{II}(x) + \sum_{i=1}^{n-2} \int_{x_{i+1}}^{\alpha_{i+1(i+2)}} E_{II}(x) + \int_{\alpha_{(n-1)n}}^{1} E_{III}(x) dx$$

and by inversion:



Talk Higher-order asymptotic formula for the eigenvalues of Sturm-Liouville ...

$$\rho_n = \frac{n\pi - \frac{\pi}{2}}{\int_0^1 \sqrt{r_+(x)} dx} - \frac{1}{n\pi} \left[\frac{[4(\nu_1 - 1)^2 - 1] + [4(\nu_2 - 1)^2 - 1]}{4\int_{x_1}^{x_2} \sqrt{r(t)} dt} + \frac{[4(\nu_2 - 1)^2 - 1] + [4(\nu_3 - 1)^2 - 1]}{4\int_{x_2}^{x_3} \sqrt{r(t)} dt} + \frac{[4(\nu_{n-1} - 1)^2 - 1] + (4(\nu_n - 1)^2 - 1)}{4\int_{x_{n-1}}^{x_n} \sqrt{r(t)} dt} - \frac{1}{2} P(x_1, x_2, x_3, ..., x_n) \right] + O(\frac{1}{n^2}).$$
(12)

REMARK. Note that the reader can obtain asymptotic distribution of eigenvalues in different types of (TP) by consideration of combination of the above cases.

References

- M. Abramowitz and J.A. Stegun, Hand book of Mathematical Function, Appl. Math. Ser., no. 55, U.S. Govt. Printing office, Washington, D.C., 1964.
- [2] F.V. Atkinson and A. Mingarelli, Asymptotics of the number of zeros and the eigenvalues of general weighted Sturm-Liouville problems, Journal der reinen und angewandten Mathematik, 395(1986)380-93.
- [3] W. Eberhard, G. Freiling and A. Schneider, Connection formulae for second- order differential equations with a complex parameter and having an arbitrary number of turningn points, Math. Nachr, 85(1994)205-229.
- [4] W. Eberhard, G. Freiling and A. Schneider, On the distribution of the eigenvalues of a class of indefinite eigenvalues problems. J. Diff. and Integral Equ, 3(1990)1167-1179.
- [5] W. Eberhard and G. Freiling, The distribution of the eigenvalues for second-order eigenvalues problems in the presence of an arbitrary number of turning points, Results in Mathematics. Birkhäuser Verlag, 21(1992)24-24.
- [6] G. Freiling and V. Yorko, Inverse Sturm-liouville problems and their applications, Nova Science Publishers, Inc, 2001.
- [7] B.J. Harris and S.T. Talarico, On the eigenvalues of second-order linear differential equations with fractional transition points, Mathematical Proceedings of the Royal Irish Academy, 99A(1)(1999)29-38.
- [8] A. Jodayree Akbarfam and A. Mingarelli, Higher-order asymptotic distribution of the eigenvalues of nondefinite Strurm -Liouville problems with one turning point, J. Com. App. Math, 149(2002)423-437.
- [9] A. Jodayree Akbarfam and A. Mingarelli, Higher-order asymptotic of the eigenvalues of Strurm -Liouville problems with a turning point of arbitrary order, Cana. App. math. Quart, 12(2004)35-60.
- [10] R.E. Langer, The boundary problem associated with a Differential equation in which the cofficient of the parameter changes sign. Transact. Amer. Math. Soc, 31(1929)1-24.
- [11] A. Leung, Distribution of eigenvalues in the presence of higher order turning points, Transact. Amer. Math. Soc, 229(1977)11-135.
- [12] A.B. Mingarelli, Asymptotic distribution of eigenvalues of non-definit Sturm-Liouville problems in Ordinary differential equations and operators, W.N.Everitt and R.T.Lewis(eds), Lecture Notes in Math. 1032, Springerverlag, berlin, 1032(1983)375-383.
- [13] F. W. J. Olver, Asymptotics and Special Functions. Acad. Press, New York 1974.
- [14] S.T. Talarico, The asymptotic form of eigenvalues for indefinite Sturm-Liouville problems, Unpublished Ph.D. thesis, Northern Illinois University, 1995.

Email:dastmalchi@iaut.ac.ir