

The production of new potential secondary salt tolerant *Tritipyrum* introgression genotypes (NPSSTIGs) and characterization the substitution lines (5D/5E^b) in 6x wheat × 6x primary *tritipyrum* progenies (F₂, F₃) by cytology and molecular cytogenetic methods

Zohre Pourfereidouni¹, Hossein Shahsevand hassani^{2,3,4} and Amin Baghizadeh⁵

¹Agricultural Biotechnology M.SC student of Shahid Bahonar University of Kerman and International Center for Science, High Technology and Environmental Science, Kerman, Iran.

Zohre.pourfereidouni@gmail.com

²College of Agriculture, Shahid Bahonar University of Kerman, Kerman, Iran

³International Center for Science, High Technology and Environmental Science, Kerman, Iran

⁴College of agriculture, Shiraz University, Shiraz, Iran.

Hassani@mail.uk.ac.ir

⁵International Center for Science, High Technology and Environmental Science, Kerman, Iran.

Amin_4156@yahoo.com

Abstract

Intergeneric hybridization is one way to transferring desirable gene/s into field crops. In order to improve primary *Tritipyrum* lines by D genome of bread wheat (*Triticum aestivum*, 2n=6x=42, AABBDD). The various 167 crosses between 6x bread wheat cultivars (AABBDD) and primary *Tritipyrum* lines (2n=6x=42, AABBE^bE^b) was carried out and 49 F₁ seeds was produced. The chromosomal counting in 291 mitotic cells of selfed progenies including, 44 F₂/F₃ plants and 19 F₂/F₃ plants (217 meiotic cells) of new potential secondary salt tolerant *Tritipyrum* introgression plants (NPSSTTIPs) by classical cytology indicated the aneuploidy reduction in comparison with aneuploidy amount F₁ progeny of 6x wheat and primary *tritipyrum* crosses. The presence of single chromosome/s in F₁ NPSSTTIPs plants showed no chromosomal homeology between D and E^b chromosomes. The assessment of E^b chromosomes in mitotic spreads of individual F₃ NPSSTTIPs plants derived from bread wheat (cv:omid) × primary *Tritipyrum* (line: St/b) and bread wheat (cv: navid) × primary *Tritipyrum* (line: (Ka/b)(Cr/b)), respectively, by in situ hybridization, using genomic DNA of *pseudoroegneria stipifolia* species (2n=2x=14, SS) as probe, labeled with Biotin-11-dUTP, indicated 1-6 single E^b chromosome in these two progenies. This probe was recognized as a powerful tool for selecting the chromosomal constitution of NPSSTTIP plants containing 14'A, 14'B, 12'D, 2'E^b chromosomes in F₃ segregation generations from 6x primary *Tritipyrum* × bread wheat hybridization. If the 5E^b chromosome of primary *Tritipyrum* substitutes with 5D chromosome of bread wheat, then the substitution (5D/5E^b) lines will be obtained.

Key words: primary *tritipyrum*, secondary *tritipyrum*, intergeneric hybridization, *Pseudoroegneria stipifolia* genomic probe, genomic in situ hybridization (GISH).