

nursing in pediatric's accident

Mohammadreza mostafaei^{*1}, fatemeh hassandost¹

1- Instructor, Qazvin of medical sciences university, faculty nursing midwifery, Qazvin, Iran.

*-correspond author: Mohammadreza mostafaei. email: raar144@yahoo.com

Introduction: The child is able to perform actions such as running, jumping and driving a tricycle will be investigated to explore your surroundings without the presence of adults. Any tension between parents and children influenced her behavior and could lead to injury. Most fall injuries due to emergency department visits from birth to 12 months of age and every 1/5 minutes injury to an infant. Shaken baby syndrome Shaken Baby Syndrome (SBS)That is, shaken baby syndrome, a type of behavior in infants and small children are more serious.

Methods: this research is a cross-section literature review based on resources

Results: Severe shaking causes the brain in the skull and rotated to create a force that can not break blood vessels and neurons. Many symptoms such as vomiting, irritability, poor feeding and lethargy, malaise often seen as a normal stage of infancy and childhood. In relatively severe type of seizure disorder symptoms may apnea, bradycardia, and level of consciousness or death.

Conclusion: the rapid growth of sensory and motor denounces the environment is very curious. Parents and caregivers should be aware of the dangers that they can have sufficient Mrtbathdyd. Therefore, parents should provide training in prenatal guidance. However, despite the events in the period of infancy is high.

Key words: Children, Accident Nursing

